

A simple way to deliver vacuum therapy: the tube-in-tube endoluminal vacuum therapy modification

A patient underwent endoscopic submucosal dissection of the middle esophagus to treat a superficial squamous cell carcinoma (T1a). He was admitted to the emergency room 2 weeks later with signs of sepsis. Computed tomography (CT) scan showed a mediastinal abscess, and an endoscopic evaluation revealed esophageal perforation at the resection site communicating with the mediastinal abscess. We decided to manage this situation with vacuum therapy [1–3]. However, a small esophageal perforation orifice and a much larger mediastinal abscess raised the question of how to position the sponge into the mediastinum. Inspired by the experience of pediatric surgeons, we assembled an aspiration device using a 12-Fr Levin tube within a 20-Fr Levin tube. The vacuum pump was attached to the inner tube, and the outer tube functioned to prevent aspiration biopsies or clogging. No sponge was used, and the aspiration device was easily positioned inside the mediastinal abscess by sliding it over a guidewire (► **Video 1**).

As the patient's clinical status improved, the conservative approach was maintained, and three other endoscopic examinations were performed, showing progressive cavity size reduction associated with granulation tissue formation. The fourth endoscopy showed complete cavity closure. Balloon dilation was needed to treat the resulting esophageal stricture.

We hypothesized that this technical modification to deliver negative pressure with a tube inside another tube may render endoluminal vacuum therapy easier, potentially reducing the number of sessions because there is no need to replace sponges. Further studies are needed.

Endoscopy_UCTN_Code_CPL_1AH_2AG



► **Video 1** Endoluminal vacuum therapy using a tube-in-tube approach to treat mediastinal abscess and esophageal perforation 2 weeks after endoscopic submucosal dissection to treat superficial squamous cell carcinoma.

Competing interests

The authors declare that they have no conflict of interest.

The authors

Marcelo Lima^{1,2}, Gustavo Lima¹, Caterina Pennacchi¹, Rodrigo Scomparin¹, Gustavo de Paulo¹, Bruno Martins¹, Fauze Maluf-Filho^{1,2}

- 1 São Paulo University Faculty of Medicine, Cancer Institute, Endoscopy Division, São Paulo, Brazil
- 2 Hospital Beneficência Portuguesa de São Paulo, Division of Endoscopy, São Paulo, Brazil

Corresponding author

Marcelo Lima, MD

Sao Paulo University Faculty of Medicine, Cancer Institute, Endoscopy Division, Rue Doutor Arnaldo, 251, São Paulo 01246-903, Brazil
Fax: +55-11-3892-2296
marcelosimasdelima@gmail.com

References

- [1] Wedemeyer J, Schneider A, Manns MP et al. Endoscopic vacuum-assisted closure of upper intestinal anastomotic leaks. *Gastrointestinal Endosc* 2008; 67: 708–711
- [2] Bludau M, Hölscher AH, Herbold T et al. Management of upper intestinal leaks using an endoscopic vacuum-assisted closure system (E-VAC). *Surg Endosc* 2014; 28: 896–901
- [3] Newton NJ, Sharrock A, Rickard R et al. Systematic review of the use of endo-luminal topical negative pressure in oesophageal leaks and perforations. *Dis Esophagus* 2017; 30: 1–5

Bibliography

Endoscopy 2021; 53: E317

DOI 10.1055/a-1270-6582

ISSN 0013-726X

published online 19.10.2020

© 2020, Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany