




Brisk bleeding after gastric lesion biopsy – possible needle tract seeding after endoscopic ultrasound-guided fine-needle biopsy of a pancreatic metastasis from renal cell carcinoma




► **Fig. 1** Color-coded duplex endoscopic ultrasound (EUS) prior to EUS-guided fine-needle biopsy showed a hypoechoic mass with increased vascularity in the pancreatic body.



► **Video 1** Endoscopic ultrasound (EUS)-guided biopsy and bleeding management in a case of possible needle tract seeding after EUS-guided fine-needle biopsy of pancreatic metastasis from renal cell carcinoma.



► **Fig. 2** Narrow-band imaging of the mucosal lesion on the posterior gastric wall presented an aberrant vascular pattern.



► **Fig. 3** Gastroscopic image of the arterial bleeding after forceps biopsy.


A 62-year-old patient had undergone left-sided partial nephrectomy due to renal cell carcinoma. The postoperative tumor classification was pT1a, pNx, L0, V0, G2, R0.

The patient presented 6 years later with abdominal pain and unintended weight loss. Computed tomography (CT) imaging indicated multiple pancreatic lesions, therefore an endoscopic ultrasound (EUS)-guided transgastric fine-needle biopsy was performed using the Procore 19G needle (Cook Medical, Limerick, Ireland) (► **Fig. 1**). Recurrence of the renal cell carcinoma was diagnosed. The tumor board decided for a pancreatotomy with splenectomy as there were no further metastases.


The patient was readmitted 6 months later because of neck swelling. The CT scan revealed a nodular goiter and a pneumomediastinum of unknown origin. Subsequent bronchoscopy and gastroscopy excluded perforation as the cause of the pneumomediastinum, which remained unclear. However, a mucosal

lesion presenting an aberrant vascular pattern was detected on the posterior wall of the gastric body (► **Fig. 2**). Forceps biopsy led to arterial bleeding (► **Fig. 3**). An over-the-scope-clip (OTSC; Ovesco, Tübingen, Germany) was applied to control the bleeding (► **Video 1**). Histological examination showed a renal cell carcinoma underneath the gastric mucosa (► **Fig. 4**). Since the location of the gastric lesion corresponded to the fine-needle biopsy site, it was most likely the procedure had caused needle tract seeding to the gastric wall. Because fine-needle biopsy of the nodular goiter also revealed metastases of the renal cell carcinoma, a thyroidectomy and gastric wedge-resection were performed (► **Fig. 5**).

Pancreatic metastases are rare, with a reported incidence varying from 1.6% to 11% [1]. The most common metastasis to the pancreas is renal cell carcinoma [2]. EUS-guided fine-needle biopsy is considered a safe technique with few adverse events. However, needle tract seeding,



► **Fig. 4** Histological examination of the gastric biopsy revealed a clear cell renal cell carcinoma growing underneath the gastric mucosa.



► **Fig. 5** Intraoperative image during the gastric wedge resection showed the renal cell carcinoma metastasis caused by needle tract seeding.

although uncommon, is a serious adverse event that may impair patient's outcome [3]. Considering the associated risk, EUS-guided fine-needle biopsy should be carried out only when the results obtained are useful for therapeutic decision-making [4], and the needle tract line should be placed within the surgical resection margins.

Competing interests

The authors declare that they have no conflict of interest.

The authors

Simone Freund¹, **Tina Schaller²**, **Claus Schöler³**, **Helmut Messmann¹**, **Stefan K. Gölder¹**

- 1 Department of Gastroenterology, University Hospital Augsburg, Germany
- 2 Institute of Pathology, University Hospital Augsburg, Germany
- 3 Department of General, Visceral and Transplantation Surgery, University Hospital Augsburg, Germany

Corresponding author

Simone Freund, MD

Department of Gastroenterology, University Hospital Augsburg, Stenglinstr. 2, 86156 Augsburg, Germany
simone.freund@uk-augsburg.de

References

- [1] Adsay NV, Andea A, Basturk O et al. Secondary tumors of the pancreas: an analysis of a surgical and autopsy database and review of the literature. *Virchows Archiv* 2004; 444: 527–535
- [2] Smith AL, Odronic SI, Springer BS et al. Solid tumor metastases to the pancreas diagnosed by FNA: a single-institution experience and review of the literature. *Cancer Cytopathol* 2015; 123: 347–355
- [3] Minaga K, Takenaka M, Katanuma A et al. Needle tract seeding: an overlooked rare complication of endoscopic ultrasound-guided fine-needle aspiration. *Oncology* 2017; 93: 107–112
- [4] Fujii LL, Levy MJ. Basic techniques in endoscopic ultrasound-guided fine needle aspiration for solid lesions: adverse events and avoiding them. *Endosc Ultrasound* 2014; 3: 35–45

Bibliography

Endoscopy 2022; 54: E380–E381

DOI 10.1055/a-1541-7061

ISSN 0013-726X

published online 9.8.2021

© 2021. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany