


Endoscopic findings of paraduodenal or groove pancreatitis

Paraduodenal pancreatitis is an uncommon type of focal chronic pancreatitis characterized by scarring of the pancreaticoduodenal space. The possible pathogenesis includes disconnection or impairment of communication between the ducts of Santorini and Wirsung [1]. The clinical presentation of paraduodenal pancreatitis can be similar to that of pancreatic head carcinoma, so its diagnosis and treatment is usually a challenge. Other previous entities with similar clinicopathological features, such as duodenal cystic dystrophy, duodenal wall cyst, or groove pancreatitis, should no longer be used [2, 3].


The clinical presentation of paraduodenal pancreatitis is secondary to duodenal stenosis and comprises recurrent episodes of abdominal pain, nausea, and vomiting. Conservative medical treatment is the standard of care, but surgery accounts for 12.8%–19.5% of pancreaticoduodenectomies performed for chronic pancreatitis [3]. This occurs because of misdiagnosis as pancreatic cancer and failure of medical treatment. Herein, we present a case of paraduodenal pancreatitis with its unique and illustrative endoscopic findings.

A 49-year-old man with a past history of chronic alcohol consumption presented with epigastric abdominal pain, vomiting, reduced oral intake, and diarrhea. He had a history of admissions for acute alcoholic hepatitis and Wernicke–Korsakow encephalopathy. His physical examination revealed abdominal tenderness, while no abdominal mass was detected. Laboratory tests revealed macrocytic anemia, hyperamylasemia ($>3\times$ normal), and cholestasis, along with low pancreatic fecal elastase levels. A computed tomography (CT) scan revealed cystic thickening of the duodenal wall, with duodenal stenosis (**► Fig. 1**).

Upper gastrointestinal endoscopy was performed to rule out portal hyperten-



► Fig. 1 Computed tomography (CT) scan showing cystic thickening of the duodenal wall, with duodenal stenosis, fibrous tissue within the pancreaticoduodenal groove, and swelling of the pancreatic head, with a 30-mm hypodense cystic lesion communicating with a slightly dilated pancreatic duct.



► Video 1 Upper gastrointestinal endoscopy showing a congested, edematous, and erythematous mucosa in the second part of duodenum, around the major papilla.

sion; neither varices or portal hypertension gastropathy were found. However, a duodenal stricture, allowing passage of the scope, was observed. The duodenal mucosa was congested, edematous, and erythematous in the second part of duodenum, around the major papilla (**► Video 1**). Conservative management with fasting and parenteral nutrition for 4 days was successful. Pancreatic enzyme replacement therapy was initiated and




the patient was discharged 7 days after the endoscopy.

Endoscopy_UCTN_Code_CCL_1AB_2AG

Competing interests

The authors declare that they have no conflict of interest.

The authors

Pedro López-Muñoz , **Vicente Lorenzo-Zúñiga** , **Noelia Alonso-Lázaro**, **María García-Campos**, **Lidia Argüello**, **Marco Bustamante-Balén** , **Vicente Pons-Beltrán**

Gastrointestinal Endoscopy Unit, Department of Gastroenterology and Hepatology, Hospital Universitari i Politècnic La Fe/IIS La Fe, Valencia, Spain

Corresponding author

Pedro López-Muñoz, MD

Gastrointestinal Endoscopy Unit,
Department of Gastroenterology and
Hepatology, Hospital Universitari i Politècnic
La Fe/Instituto Investigación Sanitaria La Fe,
Avenida de Fernando Abril Martorell, 106,
46026 Valencia, Spain
pedro.lopez8928@gmail.com

References

- [1] Chantarojanasiri T, Isayama H, Nakai Y et al. Groove pancreatitis: endoscopic treatment via the minor papilla and duct of Santorini morphology. *Gut Liver* 2018; 12: 208–213
- [2] Arora A, Dev A, Mukund A et al. Paraduodenal pancreatitis. *Clin Radiol* 2014; 69: 299–306
- [3] Adsay NV, Zamboni G. Paraduodenal pancreatitis: a clinico-pathologically distinct entity unifying “cystic dystrophy of heterotopic pancreas”, “para-duodenal wall cyst”, and “groove pancreatitis”. *Semin Diagn Pathol* 2004; 21: 247–254

Bibliography

Endoscopy 2022; 54: E735–E736

DOI 10.1055/a-1773-0499

ISSN 0013-726X

published online 10.3.2022

© 2022. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>