


Successful resection of a 22-cm subcircumferential rectal lesion treated by endoscopic submucosal dissection with a mucosal-anchor-assisted multiple-tunnel method


Colorectal endoscopic submucosal dissection (ESD) has become widespread since the development of devices and traction or tunnel methods, but ESD for large lesions remains challenging. In particular, when ESD using the tunnel method is used on huge lesions, if the mucosa and submucosa remaining after tunnel creation are dissected, the dissected parts will accumulate on the oral side. This makes the final dissection difficult because the advantage of gravity is lost. We report a case of ESD of a large lesion using a modified tunnel method with a mucosal anchor.

A 78-year-old man with chronic diarrhea underwent colonoscopy, which identified a subcircumferential laterally spreading tumor occupying nearly the entire rectum (► Fig. 1, ► Fig. 2). Because magnifying endoscopic examination provided no clear evidence of a submucosal invasive carcinoma, ESD was performed (► Video 1). After three tunnels had been created, the anal-side mucosa was left in place as an anchor to maintain good traction until the final dissection. The submucosal tunnels allowed easy identification of the dissection line, and a mucosal anchor prevented the lesion from gathering on the oral side (► Fig. 3). Finally, the remaining mucosal anchors were removed, and ESD was completed without complications (► Fig. 4). Histopathology indicated a well-differentiated intramucosal adenocarcinoma, 220×150 mm in size, with negative resection margins and no lymphovascular invasion. At 1 and 2 months later, stenosis was observed at the ESD site, requiring three endoscopic balloon dilations (► Fig. 5). At the 6-month follow-up, the patient was asymptomatic.


There are few case reports of ESD for lesions larger than 20 cm, and those that exist all describe circumferential lesions [1, 2]. In terms of stenosis, it is desirable to preserve as much mucosa as possible



► Fig. 1 Endoscopic view of a subcircumferential laterally spreading tumor in the rectum of a 78-year-old man.



► Fig. 2 Chromoendoscopic view with indigo carmine dye.




► Video 1 Subcircumferential rectal lesion treated by endoscopic submucosal dissection using a mucosal-anchor-assisted multiple-tunnel method.

[3]; however, this may be more difficult technically for subcircumferential than for circumferential lesions because peripheral incisions are required. In this case, ESD was completed safely with effective traction on the lesion by creating tunnels and leaving a mucosal anchor.


Endoscopy_UCTN_Code_TTT_1AQ_2AD

Competing interests


The authors declare that they have no conflict of interest.



► **Fig. 3** Endoscopic view after creation of three tunnels with a mucosal anchor (arrows).



► **Fig. 4** Ulcer floor after endoscopic sub-mucosal dissection (ESD).



► **Fig. 5** Stenosis that occurred 2 months after ESD.

The authors

Hideyuki Chiba **Hiroki Kuwabara** **Keiji Yamada**, **Naoya Okada**, **Jun Arimoto**, **Michiko Nakaoka**

Department of Gastroenterology, Omori Red Cross Hospital, Tokyo, Japan

Corresponding author

Hideyuki Chiba, MD, PhD

Department of Gastroenterology,
Omori Red Cross Hospital, 4-30-1 Chuo,
Ota-ku, Tokyo 143-8527, Japan
h.chiba04@gmail.com

References

- [1] Ohara Y, Toyonaga T, Tsubouchi E et al. Clinical course after endoscopic submucosal dissection in the rectum leaving a circumferential mucosal defect of 26 cm in length. *Endoscopy* 2016; 48: E4–E5
- [2] Takahashi H, Hayashi Y, Sunada K et al. Complete resection of a 225-mm circumferential rectosigmoid intramucosal carcinoma by endoscopic submucosal dissection. *Endoscopy* 2016; 48: E161–E162
- [3] Hayashi T, Kudo SE, Miyachi H et al. Management and risk factor of stenosis after endoscopic submucosal dissection for colorectal neoplasms. *Gastrointest Endosc* 2017; 86: 358–369

Bibliography

Endoscopy 2022; 54: E746–E747

DOI 10.1055/a-1778-3448

ISSN 0013-726X

published online 17.3.2022

© 2022. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>