


## Underwater endoloop-assisted endoscopic resection for colorectal pedunculated polyps




Detachable snares, such as endoloops, are effective for the removal of pedunculated colorectal polyps; however, they are often difficult to apply [1–3]. The underwater method and related techniques using instruments with waterjet functions are increasingly being developed and applied for the resection of colonic polyps [4, 5]. Here, we demonstrate the use of the underwater method to facilitate endoloop-assisted polypectomy in four patients with pedunculated colonic polyps (► **Video 1**).

The first case involved a 25-mm polyp with a thick stalk and a large villous head. First, we attempted to maneuver the endoloop in gas, but this was difficult because the loop clung to the rough surface of the polyp head. However, the head floated and its surface turned smooth in water, which allowed the endoloop to pass over it more easily (► **Fig. 1**). We also used the waterjet to assist in this process. En bloc resection was performed. The second polyp did not have a large head, but it was floppy. Therefore, a method similar to that used in case #1 was performed to remove the lesion (► **Fig. 2**).

In the third case, a large pedunculated lipoma was suspected in a patient with abdominal pain, and we believed that this lesion might have caused intermittent obstructive symptoms. The lesion had a very long stalk with a 25-mm head. As in the previous cases, manipulation of the endoloop around the head of the lesion, which was very floppy, did not result in adequate application of the endoloop. However, as expected, the lesion floated upon instillation of water to fill the lumen as fat has a lower density than water. The endoloop was then easily applied to the stalk base (► **Fig. 3**). In the case of the



► **Video 1** Four cases of underwater endoloop-assisted snare resection for pedunculated polyps are shown.




► **Fig. 1** Endoscopic images showing: **a** underwater application of an endoloop around a pedunculated polyp; **b** subsequent underwater snaring of the polyp.

fourth polyp, we demonstrated that the snaring process was easier and more precise in water than in gas.


Endoscopy\_UCTN\_Code\_TTT\_1AQ\_2AZ

### Acknowledgments

We would like to thank Drs. Rei Higashiura, Masaki Kawabata, and Yusuke Seiki for their assistance in clinical practice, and would like to thank Editage ([www.editage.com](http://www.editage.com)) for English language editing.



► **Fig. 2** Endoscopic images showing: **a** a pedunculated polyp that floated in water; **b** application of an endoloop, which was performed in water.



► **Fig. 3** Endoscopic image showing underwater application of an endoloop around a pedunculated lipoma.

## Competing interests

The authors declare that they have no conflict of interest.

## The authors

**Shunsuke Yamamoto**<sup>1</sup>  **Adolfo Parra-Blanco**<sup>2</sup>

- 1 Department of Gastroenterology and Hepatology, National Hospital Organization Osaka National Hospital, Osaka, Japan
- 2 NHR Nottingham Biomedical Research Centre, Department of Gastroenterology, Nottingham University Hospitals NHS Trust and University of Nottingham, Nottingham, UK

## Corresponding author

**Shunsuke Yamamoto, MD, PhD**  
2-1-14 Houenzaka, Chuo-ku, Osaka,  
540-0006, Japan  
shun0515suke@gmail.com

## References

- [1] Iishi H, Tatsuta M, Narahara H et al. Endoscopic resection of large pedunculated colorectal polyps using a detachable snare. *Gastrointest Endosc* 1996; 44: 594–597
- [2] Kouklakis G, Mpoumpoumaris A, Gatopoulou A et al. Endoscopic resection of large pedunculated colonic polyps and risk of post polypectomy bleeding with adrenaline injection versus endoloop and hemoclip: a prospective, randomized study. *Surg Endosc* 2009; 23: 2732–2737
- [3] Matsushita M, Hajiro K, Takakuwa H et al. Ineffective use of a detachable snare for colonoscopic polypectomy of large polyps. *Gastrointest Endosc* 1998; 47: 496–499
- [4] Binmoeller KF, Weiler F, Shah J et al. 'Underwater' EMR without submucosal injection for large sessile colorectal polyps (with video). *Gastrointest Endosc* 2012; 75: 1086–1091
- [5] Yahagi N, Nishizawa T, Sasaki M et al. Water pressure method for duodenal endoscopic submucosal dissection. *Endoscopy* 2017; 49: E227–E228

## Bibliography

*Endoscopy* 2022; 54: E835–E836  
DOI 10.1055/a-1824-5056  
ISSN 0013-726X  
published online 13.5.2022  
© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)  
Georg Thieme Verlag KG, Rüdigerstraße 14,  
70469 Stuttgart, Germany



## ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at  
<https://mc.manuscriptcentral.com/e-videos>