



## Direct retroperitoneal necrosectomy through an insufficient jejunal blind stump after gastrectomy for necrotizing pancreatitis




A 53-year-old patient with a history of alcohol abuse presented with acute severe epigastric pain. Computed tomography (CT) showed signs of acute pancreatitis with a 9-cm measuring walled-off pancreatic necrosis (WOPN) in the pancreatic tail, with broad-based contact to the greater curvature of the stomach. Initial gastroscopy revealed severe ischemic gastric wall necrosis without signs of perforation (► **Fig. 1**). An electrocautery-enhanced lumen-apposing metal stent (LAMS; 15×10 mm) was implanted transgastrically under endoscopic ultrasound guidance, to enable direct necrosectomy (► **Fig. 2**). Because of a suspicion of splenic infarction, another CT scan was performed; this showed free air collections in the upper abdomen, with urgent suspicion of gastric wall perforation in the area of the ischemic gastric wall. A gastrectomy was done, with reconstruction by esophagojejunostomy and Roux-en-Y anastomosis. Gastroscopy 6 days postoperatively revealed an anastomotic leak at the esophagojejunostomy. Endoscopic vacuum therapy was started with changes every 3–4 days. At 16 days postoperatively, a second endoscopic vacuum sponge was inserted into a newly occurring insufficiency in the jejunal blind stump that accessed the 5-cm necrotic pancreatic cavity. At 3 weeks later, after complete healing of the esophagojejunostomy anastomosis, vacuum therapy was ended, and a LAMS (20×16 mm) was implanted in the jejunal blind stump providing access for necrosectomy of the WOPN (► **Video 1**). After five extensive endoscopic necrosectomies, the stent was removed. The patient was free of infection up to that time and was discharged from the hospital. At follow-up gastroscopy 1 week later, the jejunal blind stump had healed except for a 6-mm blind-ending fistula without secretion.



► **Video 1** Direct retroperitoneal necrosectomy through an insufficient jejunal blind stump, after gastrectomy for necrotizing pancreatitis.



► **Fig. 1** Gastroscopy after transgastric implantation of a lumen-apposing metal stent (LAMS), with the intention of draining a walled-off pancreatic necrosis collection. White arrow, distal flange of the LAMS; black arrow, severe gastric wall necrosis.



► **Fig. 2** Endosonographic ultrasound showing a 10.4×6.6-cm walled-off pancreatic necrosis (WOPN) cavity with inhomogeneous content.

Acute pancreatitis is a common disease with an unpredictable course and a wide range of severity [1,2]. This case highlights the difficulty in managing the potential complications and describes how a secondary post-surgical complication enabled an unusual approach for endoscopic treatment.

Endoscopy\_UCTN\_Code\_TTT\_1AR\_2AI

### Competing interests

The authors declare that they have no conflict of interest.

## The authors

Antonia Gantschnigg<sup>1</sup> , Franz Singhartinger<sup>1</sup> , Iris Mühlbacher<sup>1</sup>, Oliver Koch<sup>1</sup>, Klaus Emmanuel<sup>1</sup>, Josef Holzinger<sup>1</sup>

<sup>1</sup> Department of General, Visceral and Thoracic Surgery, Paracelsus Medical University/ Salzburger Landeskliniken (SALK), Salzburg, Austria

## Corresponding author

**Antonia Gantschnigg, MD**

Department of General, Visceral and Thoracic Surgery, University Hospital Salzburg, Müllner Hauptstraße 48, 5020 Salzburg, Austria  
a.gantschnigg@salk.at

## References

- [1] Boxhoorn L, Voermans RP, Bouwense SA et al. Acute pancreatitis. *Lancet* 2020; 396: 726–734
- [2] Zerem E. Treatment of severe acute pancreatitis and its complications. *World J Gastroenterol* 2014; 20: 13879–13892

## Bibliography

*Endoscopy* 2022; 54: E939–E940

DOI 10.1055/a-1882-4603

ISSN 0013-726X

published online 14.7.2022

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



## ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is an open access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>