



Cap-suction underwater endoscopic mucosal resection for en bloc resection of nongranular pseudodepressed colonic lesion: a novel technique when conventional snaring is not possible

OPEN
ACCESS


Colorectal lesions with a nongranular depressed component are difficult to snare and have a high risk of submucosal invasion; thus, en bloc resection of these lesions is mandatory, usually by endoscopic submucosal dissection (ESD) [1, 2]. Underwater endoscopic mucosal resection (UEMR), described by Binmoeller in 2012 [3], allows endoscopic resection without prior submucosal injection, as the colonic lesion “floats” in a lumen filled with water along with the mucosa and submucosa. Endoscopic ultrasound has shown that when the lumen is filled with water, the muscularis propria retains a circular configuration and does not follow the involutions of the mucosa and submucosa [3]. Cap-assisted EMR using a straight distal attachment has been described to resect polyps not easily amenable to standard EMR [4, 5]. Herein, we present the case of a 16 mm, nongranular, 0-IIa + IIc “u”-shaped lesion, which was partially hidden by a fold and difficult to face, located in the sigmoid colon (► Fig. 1, ► Fig. 2).



► Fig. 1 Nongranular 0-IIa + IIc “u”-shaped lesion, which was difficult to face.




► Fig. 2 Evaluation under white-light imaging.




► Fig. 3 Japan NBI Expert Team type 2b pattern under evaluation with narrow-band imaging and near focus.



► Fig. 4 Pit pattern VI (severe) under evaluation with crystal violet and near focus.





Video 1 Cap-suction underwater endoscopic mucosal resection. **a** Previous attempt. **b** After conic-cap suction.

Evaluation under narrow-band imaging (NBI) with near focus showed Japan NBI Expert Team (JNET) type 2b (► **Fig. 3**) and chromoendoscopy showed pit pattern Vi severe-type (► **Fig. 4**).

En bloc “classic” UEMR was attempted but it was not possible to snare the lesion. Thereafter, with the lumen filled with water, a conic cap (ST hood DH-30CR; Fujifilm Europe, Düsseldorf, Germany) was used to gently aspirate the 0-IIc component, facilitating the infolding of the lesion into the gravity-free underwater environment; the lesion was then easily snared and resected by underwater EMR (► **Fig. 5**, ► **Video 1**).

As UEMR is a reversible technique because there is no injection and no deformity of the working space, we recommend trying this approach first for en bloc resection of nongranular 0-IIc lesions without overt features of malignancy, as it is a fast and cheap technique compared with ESD or endoscopic full-thickness resection. If classic UEMR does not seem feasible, conic cap aspiration before UEMR may be helpful for resection of flat depressed lesions.

Endoscopy_UCTN_Code_TTT_1AQ_2AD

Acknowledgments

We want to thank Dr. Maria Pellise for her support in the writing of this video

Competing Interest

The authors declare that they have no conflict of interest.

The authors

Hugo Uchima¹, **Anna Calm**¹, **Juan Colán-Hernández**¹, **Raquel López**², **Caterina Fumagalli**², **Alba Hernández**², **Vicente Moreno**¹

- 1 Endoscopy Unit, Gastroenterology Department, Hospital Universitari Germans Trias i Pujol, Badalona, Barcelona, Spain
- 2 Pathology Department, Hospital Universitari Germans Trias i Pujol, Badalona, Barcelona, Spain

Corresponding author

Hugo Uchima, MD
Gastrointestinal Endoscopy, Hospital Universitari Germans Trias i Pujol, Carretera de Canyet S/N, Badalona, Barcelona 08916, Spain
huchima.germanstrias@gencat.cat

References

- [1] Yamada M, Saito Y, Sakamoto T et al. Endoscopic predictors of deep submucosal invasion in colorectal laterally spreading tumors. *Endoscopy* 2016; 48: 456–464

- [2] Burgess NG, Hourigan LF, Zanati SA et al. Risk stratification for covert invasive cancer among patients referred for colonic endoscopic mucosal resection: a large multicenter cohort. *Gastroenterology* 2017; 153: 732–742
- [3] Binmoeller KF, Weilert F, Shah J et al. “Underwater” EMR without submucosal injection for large sessile colorectal polyps (with video). *Gastrointest Endosc* 2012; 75: 1086–1091
- [4] Kashani A, Lo SK, Jamil LH. Cap-assisted endoscopic mucosal resection is highly effective for nonpedunculated colorectal lesions. *J Clin Gastroenterol* 2016; 50: 163–168
- [5] Conio M, Bianchi S, Repici A et al. Cap-assisted endoscopic mucosal resection for colorectal polyps. *Dis Colon Rectum* 2010; 53: 919–927

Bibliography

Endoscopy 2023; 55: E106–E107
DOI 10.1055/a-1948-2054
ISSN 0013-726X
published online 14.10.2022
© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>