

Management of multiple esophageal leaks with an ultra-large fully covered metallic stent after aborted peroral endoscopic myotomy

OPEN
ACCESS




► **Fig. 1** Chest computed tomography with contrast, demonstrating periesophageal fluid and pleural effusion after peroral endoscopic myotomy.

Aborted peroral endoscopic myotomy (POEM) is rare, and submucosal fibrosis is the most common direct cause of technical failure [1,2]. Age, disease duration, sigmoid esophagus, and prior interventions are the main risk factors for submucosal fibrosis [2,3].


We present the case of an 85-year-old woman with type 2 achalasia previously treated with posterior approach POEM, in whom symptoms reappeared 13 months after the procedure, severely affecting her nutritional status and quality of life. A repeat POEM procedure with an anterior approach was intended.

During the procedure, a sigmoid esophagus with poor mucosal lifting and fibrotic adhesion between mucosal and muscle layers prevented the creation of the tunnel. After a second failed incision, the procedure was aborted. A type II mucosal injury occurred, which was closed promptly with endoscopic through-the-scope clips. However, frank perforation became evident soon after the procedure (► **Fig. 1**, ► **Video 1**). The patient developed sepsis and was transferred to the intensive care unit with mechanical ventilation.

A decision was taken to insert a ultra-large esophageal stent (Niti-S Mega Stent; TaeWoong Medical, Gyeonggi-do, South Korea) (► **Fig. 2**, ► **Fig. 3**, ► **Video 1**). Inotropic support was with-



► **Video 1** Use of an ultra-large stent for management of multiple esophageal leaks after peroral endoscopic myotomy.



► **Fig. 2** Fluoroscopic confirmation of stent placement.



► **Fig. 3** Esophagogram after stent placement; no leaks were observed.

drawn after 24 hours and the patient was extubated. Enteral nutrition was introduced at Day 3, and antibiotics were given for 14 days without any complications regarding the stent. Inadvertent mucosotomy and esophageal leak are the most common early complications of POEM. Endoscopic clips

are the first management approach [1, 4]. Given the extensive fibrosis, multiple incisions, and suspicion of unnoticed injuries, insertion of an ultra-large stent was chosen.


The Mega stent is an ultra-large fully covered metallic stent, tailored for the management of post-bariatric surgery leaks. Its length and shape allow large leaks to be covered with decreased risk of migration that may prove useful in complications such as multiple leaks or extensive mucosal injury, as in the current case [5].

Endoscopy_UCTN_Code_CPL_1AH_2AJ

Competing interests

The authors declare that they have no conflict of interest.

The authors

Ana Lorena Madrigal Méndez¹ , **Daniela Grant**¹, **Viviana Hernández**¹, **Kenneth Ernest-Suárez**^{1,2}, **Jorge Vargas-Madrigal**³, **Luis Diego Arguedas**¹, **Álvaro Villalobos**¹

- 1 Gastroenterology, Caja Costarricense de Seguro Social, San José, Costa Rica
- 2 University of Costa Rica, Faculty of Sciences, San José, Costa Rica
- 3 Gastroenterology Department, Hospital Enrique Baltodano Briceño, Liberia, Costa Rica

Corresponding author

Ana Lorena Madrigal Méndez, MD
Department of Gastroenterology, San Juan de Dios Hospital, Paseo Colón, San José, 10102, Costa Rica
mumacr@gmail.com

References

- [1] Zhang YQ, Yao LQ, Meidong X et al. Early diagnosis and management of esophageal leakage after peroral endoscopic myotomy for achalasia. *Turkish J Gastroenterol* 2016; 27: 97–102
- [2] Wu QN, Xu XY, Zhang XC et al. Submucosal fibrosis in achalasia patients is a rare cause of aborted peroral endoscopic myotomy procedures. *Endoscopy* 2017; 49: 736–744
- [3] Wang Y, Liu ZQ, Xu MD et al. Clinical and endoscopic predictors for intraprocedural mucosal injury during per-oral endoscopic myotomy. *Gastrointest Endosc* 2019; 89: 769–778
- [4] Haito-Chavez Y, Inoue H, Beard KW et al. Comprehensive analysis of adverse events associated with per oral endoscopic myotomy in 1826 patients: an international multicenter study. *Am J Gastroenterol* 2017; 112: 1267–1276
- [5] Shehab HM, Hakky SM, Gawdat KA. An endoscopic strategy combining Mega stents and over-the-scope clips for the management of post-bariatric surgery leaks and fistulas. *Obes Surg* 2016; 26: 941–948

Bibliography

Endoscopy 2023; 55: E478–E479
DOI 10.1055/a-2018-4213
ISSN 0013-726X
© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply, discounts and waivers acc. to HINARI are available.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>