# Snare-tip soft coagulation to treat esophageal stent epithelial hyperplasia





**Fig.1** Application of snare-tip soft coagulation (STSC) to epithelial hyperplasia. **a**, **b** Proximal and distal ends of the partially covered metal stent demonstrating epithelial hyperplasia. **c**, **d** Ablated epithelial hyperplasia following STSC. **e**, **f** Post-STSC and stent-in-stent therapy revealing near complete resolution of epithelial hyperplasia.

A 65-year-old man with a spontaneous esophageal perforation (Boerhaave syndrome) was managed with a partially covered metal stent (23 mm wide × 12.5 mm long). After an indwelling time of 8 weeks, endoscopic removal was attempted. Significant epithelial hyperplasia was noted at the proximal and distal ends of the stent (> Fig. 1 a, b). Despite treatment with argon plasma coagulation (APC) combined with stent-in-stent insertion [1], attempts at removal failed. We proceeded to utilize a dedicated 10-mm hot snare to perform snare-tip soft coaqulation (STSC; Effect 4, 80 Watts; ERBE VIO300D) (Erbe Elektromedizin, Tübingen, Germany) to ablate the epithelial hyperplasia (> Video 1). A uniform field of ablated tissue was obtained (> Fig. 1 c, d). To promote sloughing of the ablated tissue, a fully covered metal

stent was placed within the pre-existing stent. After 2 weeks, the inner stent was removed, revealing near-complete clearance of the epithelial hyperplasia (**> Fig. 1 e, f**). After a small amount of additional STSC at the proximal margin, stent removal was easily accomplished with rat-toothed forceps.

While STSC was developed as a measure to reduce recurrent adenoma post colorectal endoscopic mucosal resection [2], its favorable properties render STSC invaluable in other settings. For example, when compared to other modalities such as APC, direct contact with the snare tip enables uniform delivery of energy to complete destruction of the target tissue [3]. Furthermore, a peak voltage of 190 V results in desiccation. The ensuing lack of carbonization and charring facilitates reliable energy delivery. Lastly, because resistance to current flow increases exponentially as tissue is desiccated, excessive current flow is limited. Subsequent termination of energy transfer has the potential to prevent deep tissue injury [4]. While placement of the inner stent may have contributed to tissue destruction, we believe that STSC is a safe, costeffective, and efficacious way to treat esophageal stent epithelial hyperplasia.

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AZ

#### **Competing interests**

Michael J. Bourke: Research Support: Olympus, Cook Medical, Boston Scientific. The remaining authors have no conflicts of interest to disclose.



**Video 1** Snare-tip soft coagulation to treat esophageal stent epithelial hyperplasia.

#### The authors

### Sunil Gupta<sup>1,2</sup>, Anthony Whitfield<sup>1,2</sup>, Andrew Tang<sup>1</sup>, Eric Y. T. Lee<sup>1</sup>, Stephen J. Williams<sup>1</sup>, Nicholas G. Burgess<sup>1,2</sup>, Michael J. Bourke<sup>1,2</sup>

- 1 Westmead Hospital, Department of Gastroenterology and Hepatology, Sydney, Australia
- 2 University of Sydney, Westmead Clinical School, Sydney, Australia

#### Corresponding author

#### Michael J. Bourke, MD

University of Sydney, Department of Medicine, Westmead Hospital, Department of Gastroenterology and Hepatology, Suite 106a 151-155 Hawkesbury Road, Sydney, New South Wales, Australia 2145 Fax: +61-2-9845-5637 michael@citywestgastro.com.au

#### References

- [1] Fiocca F, Cereatti F, Antypas P et al. Argon plasma coagulation: a less-expensive alternative to the "stent-in-stent" technique for removal of embedded partially covered esophageal stents. Gastrointest Endosc 2016; 83: 453
- [2] Sidhu M, Shahidi N, Gupta S et al. Outcomes of thermal ablation of the mucosal defect margin after endoscopic mucosal resection: a prospective, international, multicenter trial of 1000 large nonpedunculated colorectal polyps. Gastroenterology 2021; 161: 163–170
- [3] Whitfield AM, Burgess NG, Bahin FF et al. Histopathological effects of electrosurgical interventions in an in vivo porcine model of colonic endoscopic mucosal resection. Gut 2022; 71: 864–870
- [4] Morris ML, Tucker RD, Baron TH et al. Electrosurgery in gastrointestinal endoscopy: principles to practice. Am J Gastroenterol 2009; 104: 1563–1574

## Bibliography

Endoscopy 2023; 55: E572–E573 DOI 10.1055/a-2050-7431 ISSN 0013-726X © 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https:// creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



### ENDOSCOPY E-VIDEOS https://eref.thieme.de/e-videos



Endoscopy E-Videos is an open access online section, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply, discounts and wavers acc. to HINARI are available.

This section has its own submission website at https://mc.manuscriptcentral.com/e-videos