# Successful endoscopic closure of a large esophageal perforation using the purse-string technique





**Video 1** Endoscopic closure of a large esophageal perforation.

Endoscopy has been increasingly utilized to manage esophageal perforations [1–3].

We present in this case a successful endoscopic closure of a large esophageal perforation using the purse-string technique [4–5] (**> Video 1**).

A 79-year-old woman with multiple comorbidities underwent a savary dilation of esophageal stricture at an outside institution. She presented with septic shock, lactic acidosis, and respiratory failure.

A chest x-ray revealed pneumomediastinum and a right-sided pneumothorax. A chest tube was inserted. She was resuscitated with intravenous fluids, bicarbonate, antibiotics, and three vasopressors. A chest computed tomography (CT) revealed pneumomediastinum and improved pneumothorax after chest tube insertion without mediastinal fluid collections. She was deemed a poor candidate for surgery.

Endoscopy revealed a large mid-esophageal perforation (**> Fig. 1**). Endoscopic suturing was challenging due to the narrow space. After a running suture was torn, we decided to switch to the pursestring technique.

An endoloop catheter and a clip were advanced through the two channels of a double-channel endoscope. The endoloop was fixed to the margins of the perforation by clips. The endoloop was then tightened approximating the margins of the defect (▶ Fig. 2) and deployed. A residual defect was seen proximally and a second endoloop was tightened and deployed similarly.

A through-the-scope esophageal stent (18 mm×149 mm) was deployed covering the site of perforation (> Fig. 3). The proximal end of the stent was sutured to the esophageal wall to prevent migration. Five days after the procedure, she was extubated and weaned off vasopressors. Two weeks after the procedure, chest CT showed a remarkable decrease in the pneumomediastinum.

She was weaned off oxygen, and the chest tube was removed. Enteral feeding was started through a jejunal extension of a preexisting gastrostomy tube.



**Fig. 1** A large full-thickness esophageal perforation (red arrow).



► **Fig.2** Endoscopic view of the stringpurse technique after tightening the endoloop.

Three weeks after the procedure, an endoscopy was performed. The esophageal stent was retrieved. The perforation site had healed (**> Fig. 4**). Contrast injection demonstrated complete sealing of the esophageal wall (**> Fig. 5**).

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AI

### **Competing interests**

The authors declare that they have no conflict of interest.



► Fig. 3 A fully covered esophageal stent deployed after the primary closure of the defect.



► Fig.4 Endoscopic view after 3 weeks, demonstrating closure of the defect.



► Fig.5 Contrast injection after esophageal stent removal showing complete sealing of the esophageal wall with no extravasation of contrast.

## The authors

#### Michael Lajin<sup>1</sup> Naser Khan<sup>2</sup>, Fateh Bazerbachi<sup>3</sup>

- 1 Sharp Healthcare, San Diego, California, USA
- 2 Javon Bea Hospital-Rockton, Rockford, Illinois, USA
- 3 CentraCare Health System, St. Cloud, Minnesota, USA

#### Corresponding author

#### Michael Lajin, MD

Sharp Healthcare, 8860 Center Drive Suite 330, La Mesa, California 91942, USA mlajin@yahoo.com

#### References

- Gomez-Esquivel R, Raju GS. Endoscopic closure of acute esophageal perforations. Curr Gastroenterol Rep 2013; 15: 1–5
- [2] Avila N, Schwartz A, Tarnasky P et al. Urgent esophageal repair with endoscopic suturing. VideoGIE 2018; 3: 77–78
- [3] Pech O. latrogenic Perforation of an advanced esophageal carcinoma treated with placement of a self-expanding metal stent. Video J Encycl GI Endosc 2013; 1: 81–82
- [4] Ryu JY, Park BK, Kim WS et al. Endoscopic closure of iatrogenic colon perforation using dual-channel endoscope with an endoloop and clips: methods and feasibility data (with videos). Surg Endosc 2019; 33: 1342–1348
- [5] Li B, Chen D, Wan X. Endoscopic pursestring suture technique: An effective remedy for a large iatrogenic rectal perforation. Dig Endosc 2019; 31: e9–e10

#### **Bibliography**

Endoscopy 2023; 55: E715–E716 DOI 10.1055/a-2078-0761 ISSN 0013-726X © 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



# ENDOSCOPY E-VIDEOS https://eref.thieme.de/e-videos



*E-Videos* is an open access online section of the journal *Endoscopy*, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high-quality video and are published with a Creative Commons CC-BY license. Endoscopy E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: https:// www.research4life.org/access/eligibility/).

This section has its own submission website at https://mc.manuscriptcentral.com/e-videos