# Gastric antral defect closure after endoscopic submucosal dissection with the reopenable clip-over-the-line method using a multibending scope



Mucosal defect closure after gastric endoscopic submucosal dissection (ESD) is challenging because of the thick mucosal and muscle layers [1]; it is particularly difficult in the gastric antrum. When a conventional endoscope is pushed distally to access the antrum, the greater curvature is stretched. This widens the mucosal defect and stretches the muscle layer, making closure of the defect difficult (> Fig. 1). Herein, we present a multibending scope technique that allows antral defect closure using the reopenable clip-over-the-line method (ROLM) to avoid stretching the muscle layer [2]. A 60-mm mucosal defect was left after qastric ESD in our patient (► Fig. 2; ▶ Video 1). A multibending scope (GIF-2TQ260M; Olympus) was used with the second (top) wheel adjusted to a slightly upward position. Consequently, it was possible to confirm the anal side of the defect edge endoscopically, without stretching the muscle layer. Defect closure was performed using the ROLM, a defect closure technique using a line and reopenable clip (SureClip 8 mm, 16 mm; MicroTech, Nanjing, China) [3]. First, a clip with a line secured to the tooth was placed at the anal edge of the defect through the working channel. The line was passed through a hole in the tooth of the next reopenable clip, before it was also introduced into the working channel. This clip was placed on the contralateral defect edge and nearby muscle layer, with the line tightened over the closed clip before it was deployed. The same steps were repeated to gradually close the defect. Two ROLMs were required for complete defect closure in this case, and the remaining line was cut with a new loop cutter (FS-5L-1; Olympus) [4]. The patient was discharged, without experiencing any adverse events.



▶ Fig. 1 Comparison of mucosal defect closure in the gastric antrum using: **a**, **b** a conventional upper gastrointestinal scope, which stretches the muscle layer of the defect owing to the stretching of the greater curvature; **c**, **d** a multibending scope, which touches the greater curvature only slightly and does not stretch the muscle layer of the defect.

In conclusion, a multibending scope is effective for closing antral mucosal defects because it prevents stretching of the gastric muscle layer.

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AG

### **Competing interests**

The authors declare that they have no conflict of interest.

#### The authors

Tatsuma Nomura<sup>1,2</sup> Shinya Sugimoto<sup>1</sup>, Taishi Temma<sup>1</sup>, Jun Oyamada<sup>1</sup>, Keiichi Ito<sup>2</sup>, Akira Kamei<sup>1</sup>, Noriya Uedo<sup>3</sup>

- 1 Department of Gastroenterology, Ise Red Cross Hospital, Ise, Mie, Japan
- 2 Department of Gastroenterology, Mie Prefectural Shima Hospital, Shima, Mie, Japan
- 3 Department of Gastrointestinal Oncology, Osaka International Cancer Institute, Osaka, Japan



▶ Fig. 2 Endoscopic images of defect closure after endoscopic submucosal dissection (ESD) in the gastric antrum using a multibending scope showing: a appearance of the 60-mm mucosal defect after ESD with a conventional upper gastrointestinal scope; b the same defect viewed with a multibending scope with degassing; c placement of the first clip with a line; d placement of a clip with a line previously threaded through the tooth on the contralateral mucosal defect edge and muscle layer; e defect closure, with the muscle layer folded over, after the placement of further clips threaded onto the line on the contralateral defect edge and muscle layer; f complete closure of the mucosal defect.



**Video 1** Defect closure after endoscopic submucosal dissection in the gastric antrum with the reopenable clip-over-the-line method using a multibending scope.

## Corresponding author

#### Tatsuma Nomura, MD

Department of Gastroenterology, Ise Red Cross Hospital, 1-471-2 Funae, Ise, Mie 516-8512, Japan m06076tn@icloud.com

## References

- Nomura T, Sugimoto S, Temma T et al. Suturing techniques with endoscopic clips and special devices after endoscopic resection. Dig Endosc 2023; 35: 287–301
- [2] Nomura T, Sugimoto S, Temma T et al. Clipline closure with the reopenable clip over line method for a large mucosal defect after gastric endoscopic submucosal dissection. Endoscopy 2022; 54: E1–E2

- [3] Nomura T, Sugimoto S, Temma T et al. Reopenable clip-over-the-line method for closing large mucosal defects following gastric endoscopic submucosal dissection: prospective feasibility study. Dig Endosc 2022. doi:10.1111/den.14466
- [4] Nomura T, Sugimoto S, Oyamada J et al. Application of a new loop cutter for nylon lines and the reopenable clip-over-line method for large defect closure after duodenal endoscopic submucosal dissection. Endoscopy 2023; 55: E503–E504

## Bibliography

Endoscopy 2023; 55: E866–E867 DOI 10.1055/a-2107-2474 ISSN 0013-726X

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



## ENDOSCOPY E-VIDEOS https://eref.thieme.de/e-videos



E-Videos is an open access online section of the journal Endoscopy, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high-quality video and are published with a Creative Commons CC-BY license. Endoscopy E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: https:// www.research4Life.org/access/eligibility/).

This section has its own submission website at

https://mc.manuscriptcentral.com/e-videos