A loop-assisted inversion technique for easy removal of a gastric stromal tumor in the fundus





▶ Fig.1 Computed tomography image showing a stromal tumor (about 2×1 cm) at the gastric fundus.

A 54-year-old woman presented with a stromal tumor (approximately 2×1cm) in the gastric fundus (> Fig. 1 and > Fig. 2a). After it had been marked and submucosal injection performed under endoscopic guidance, an electrosurgical knife was used to make a circular incision (> Video 1). This was challenging because of the difficult approach and the high risk of perforation, with an IT knife being used to make the incision (> Fig. **2b**). A clip-anchored loop was fixed 1 cm from the incised wound (> Fig. 2 c). A snare was then used to trap the incised mucosa and lift it, with the loop ring being slowly tightened (> Fig. 2d). After

the snare was released, inversion of the tumor was observed (▶ Fig.2e and ▶ Fig.3). Next, an electrosurgical knife was used to cut and expose the tumor margins, and a snare was then used to trap the tumor base and perform electrosurgical excision (▶ Fig.2f). After the excision, the clean inverted wound was sutured using clips (▶ Fig.2g). Finally, the loop was released. The resected specimen was an intact tumor measuring approximately 2×1 cm (▶ Fig.4). At follow-up 1 month later, a flat wound with a residual loop was observed (▶ Fig.5).

Endoscopic full-thickness resection (EFTR) is regularly used to treat gastric



▶ Fig.2 Schematic showing the stages involved in the procedure: **a** a tumor is present in the gastric fundus; **b** circumferential incision is performed after sufficient submucosal injection; **c** a loop is placed encircling the tumor and anchored by clips fixed 1 cm away from the incision; **d** a snare is used to entrap the incised mucosa and pull it towards the cardia, with the loop slowly tightened as the snare is pulled; **e** a protrusion that includes the tumor and normal tissue is seen after tightening of the loop and release of the snare; **f** an electrosurgical knife is used to cut and expose the tumor margins, then a snare placed around the margins of the tumor is used to trap and completely resect it by thermal snare cutting; **g** the internal wound surface is sutured, forming the double suture in combination with the loop ligation.







Fig.4 Photograph of the resected intact tumor.

stromal tumors, is considered safe, and has a clinical outcome equivalent to surgery [1]. Gastric fundal tumors are associated with a high risk of perforation [2]. If perforation occurs, infection, intraperitoneal implantation metastasis, and postoperative bleeding of the serosal surface are potential concerns [3]. Several methods have been recommended for the management of unavoidable perforations [4]. We used a clip to fix the loop around the tumor and a snare to invert it. Double-suture techniques involving loops and clips are safe, easy, and quick.

Endoscopy_UCTN_Code_TTT_1AO_2AG

Funding

the Project of Bureau of Science & Technology Nanchong City 22SXQT0401



► Fig. 5 Endoscopic appearance at follow-up showing a flat wound closed by clips with the residual loop still in place after 1 month.

Competing interests

The authors declare that they have no conflict of interest.

The authors

Zhang Tao ⁰, Long Chen, Jie Liu, Yi Ming Peng, Feng Ying Lin, Liang Sun, Jian Chen

Department of Gastroenterology, Nanchong Central Hospital, The Second Clinical Medical College, North Sichuan Medical College, Nanchong City, Sichuan, China

Corresponding author

Zhang Tao, MD

Department of Gastroenterology, Nanchong Central Hospital, The Second Clinical Medical



Fig.3 Endoscopic image showing the inverted tumor.

College, North Sichuan Medical College, Nanchong City, Sichuan, 637000, China 305514271@qq.com

References

- [1] Liu S, Zhou X, Yao YX et al. Resection of the gastric submucosal tumor (G-SMT) originating from the muscularis propria layer: comparison of efficacy, patients' tolerability, and clinical outcomes between endoscopic fullthickness resection and surgical resection. Surg Endosc 2020; 34: 4053–4064
- [2] Li L, Wang F, Wang B et al. Endoscopic submucosal dissection of gastric fundus subepithelial tumors originating from the muscularis propria. Exp Ther Med 2013; 6: 391– 395
- [3] Chen Q, Yu M, Lei Y et al. Efficacy and safety of endoscopic submucosal dissection for large gastric stromal tumors. Clin Res Hepatol Gastroenterol 2020; 44: 90–100
- [4] Lee JH, Kedia P, Stavropoulos SN et al. AGA clinical practice update on endoscopic management of perforation in gastrointestinal tract: expert review. Clin Res Hepatol Gastroenterol 2021; 19: 2252–2261

Bibliography

Endoscopy 2023; 55: E902–E903 DOI 10.1055/a-2119-0999 ISSN 0013-726X © 2023. The Author(s). This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

