


Combined laparoscopic–endoscopic resection of a bleeding giant duodenal Brunner’s gland hamartoma

OPEN
ACCESS




► **Fig. 1** Computed tomography image showing the large duodenal polyp (arrow).




► **Fig. 2** Macroscopic appearance of the specimen, which measured 50 × 35 × 15 mm.

Brunner’s gland hamartoma is a rare entity and constitutes 10.6% of all benign duodenal tumors [1]. In symptomatic patients, endoscopy represents the first-line treatment; however, there are many technical challenges that can limit endoscopic removal, including the size and location of the lesion. Therefore, surgery may be required for complex cases [2, 3]. We present the case of a healthy 41-year-old woman who presented with melena. Upper gastrointestinal endoscopy and computed tomography scanning revealed a large polyp with ulceration on the anterior wall of the duodenal bulb (► **Fig. 1**). Endoscopic ultrasonography confirmed a hypoechoic submucosal



► **Video 1** Laparoscopic-assisted polypectomy of the giant Brunner’s gland hamartoma.



► **Fig. 3** Microscopic section stained with hematoxylin and eosin (magnification ×10) showing a lining epithelium of atypical cells (probably dysplastic), stromal edema, and mild vascular congestion in relation to the Brunner’s glands.

pedunculated polyp with a 10-mm base and 50-mm head. No malignancy was revealed on biopsy.

Initially, a standard polypectomy was attempted but the head of the lesion was too large to pass through the pylorus. Therefore, a combined laparo-

scopic–endoscopic approach was planned (► **Video 1**). During the laparoscopic exploration of the abdominal cavity, the duodenal polyp with full endophytic growth was recognized. Even with laparoscopic assistance, passage of the lesion into the stomach was not possible.

Therefore, a 15-mL epinephrine solution (diluted 1:20 000) was injected into the head of the polyp to achieve volume reduction and reduce bleeding [4]. Next, the head of the lesion was pushed into the stomach by gently pressing the laparoscopic forceps along the duodenum towards the pylorus, and piecemeal resection of the head was carried out; the polypectomy was completed with the en bloc removal of the peduncle and all of the fragments were collected (► Fig. 2).

The total operative time was 80 minutes. The patient's postoperative course was uneventful. Pathology confirmed the lesion was a duodenal Brunner's gland hamartoma (► Fig. 3). No recurrence was detected at the 6-month follow-up endoscopy.

In the present case, the application of laparoscopy overcame the polyp size-related constraints, allowing endoscopic resection; the combined approach provided a safe and curative therapeutic strategy, avoiding a more invasive surgical treatment.

Endoscopy_UCTN_Code_CPL_1AH_2AZ

Competing interests

C. G. De Angelis is a consultant for Boston Scientific, Olympus, and Medi-Globe. A. Marano, M. Sacco, L. M. Rorato, S. Caronna, F. Di Giovanni, and M. Santarelli declare that they have no conflicts of interest.

The authors

Alessandra Marano^{1‡}, Marco Sacco^{2‡}, Lisa Marie Rorato¹, Stefania Caronna², Fabrizia Di Giovanni³, Mauro Santarelli¹, Claudio Giovanni De Angelis²

- 1 General and Specialist Surgery Department, Emergency General Surgery Unit, Azienda Ospedaliero Universitaria Città della Salute e della Scienza di Torino, Turin, Italy
- 2 Gastroenterology Department, Endoscopy Unit, Azienda Ospedaliero Universitaria Città della Salute e della Scienza di Torino, Turin, Italy
- 3 Pathology Unit, Azienda Ospedaliero Universitaria Città della Salute e della Scienza di Torino, Turin, Italy

Corresponding author

Marco Sacco, MD, PhD

Gastroenterology Unit, A.O.U. Città della Salute e della Scienza di Torino, Corso Bramante 88, 10126, Turin, Italy
marco.sacco10@gmail.com

References

- [1] Zhu M, Li H, Wu Y et al. Brunner's gland hamartoma of the duodenum: a literature review. *Adv Ther* 2021; 38: 2779–2794
- [2] Hu BW, Liu ZZ, Ding HF. Facts to be known for the better understanding of Brunner's gland adenoma and its management. *Asian J Surg* 2022; 45: 1295–1296
- [3] Pantaleón Sánchez M, Pioche M, Figueiredo Ferreira M et al. Giant Brunner's gland hamartoma of the duodenal bulb removed by endoscopic submucosal dissection. *Endoscopy* 2022; 54: E1018–E1019
- [4] Walker J, Howell DA, Gupta S et al. Combined volume reduction with the loop-and-leave technique permits safe endoscopic management of high-risk giant gastric polyps. *Gastrointest Endosc* 2016; 83: 655–656

Bibliography

Endoscopy 2023; 55: E983–E984

DOI 10.1055/a-2134-9639

ISSN 0013-726X

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited.

(<https://creativecommons.org/licenses/by/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



E-Videos is an open access online section of the journal *Endoscopy*, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high-quality video and are published with a Creative Commons CC-BY license. Endoscopy E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: <https://www.research4life.org/access/eligibility/>).

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>

[‡] Joint first authors