Time to Completion of Two-Step Screening for Gestational Diabetes and Adverse Outcomes

Sarah A. Nazeer, MD¹ Han-Yang Chen, PhD¹ Joycelyn A. Cornthwaite, CDCES¹ Sandra Sadek, MD¹ Tala Ghorayeb, MD¹ Nahla Daye, MD¹ Suneet P. Chauhan, MD, Hon DSc¹ Baha Sibai, MD¹ Michal F. Bartal, MD, MS^{1,2}

¹ Division of Maternal-Fetal Medicine, Department of Obstetrics, Gynecology, and Reproductive Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston, Houston, Texas ² Department of Obstetrics and Gynecology, Sheba Medical Center,

Tel Hashomer, Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel

Am J Perinatol 2024;41(suppl S1):e2679-e2685.

Abstract

Objective This study aimed to ascertain whether the length of time to complete the gestational diabetes mellitus (GDM) screening was associated with adverse neonatal outcomes.

Address for correspondence Sarah A. Nazeer, MD, Division of

Maternal-Fetal Medicine, Department of Obstetrics, Gynecology, and

Reproductive Sciences, McGovern Medical School, The University of

Texas Health Science Center at Houston 6431 Fannin Street, Suite

3.264, Houston, TX 77030 (e-mail: sarah.a.nazeer@uth.tmc.edu).

Study Design This was a retrospective cohort study of singleton, nonanomalous individuals who were screened for GDM at \geq 24 weeks' gestation at an academic hospital system. We compared outcomes among people who were diagnosed with GDM and completed the 3-hour glucose tolerance test (GTT) \leq 14 second versus >14 days from the 1-hour glucose challenge test (GCT). The primary outcome was a composite adverse neonatal outcome of the following: large for gestational age, shoulder dystocia, birth injury, respiratory distress, hypoglycemia, or fetal/neonatal death. The secondary outcomes included several individual neonatal and maternal morbidities. Multivariable Poisson's regression models were used to evaluate the association. Adjusted relative risk (aRR) and 95% confidence intervals (CI) were calculated.

Results Among the 313 individuals who completed the two-step screening for GDM and had an 1-hour GCT \geq 135 mg/dL; of them, 171 (54.6%) completed the 3-hour GTT \leq 14 days, 142 (45.4%) completed the 3-hour GTT > 14 days. Overall rate of the primary outcome was 44.1%. After multivariable adjustment, the risk of the primary outcome was similar between people who completed the two-step method in \leq 14 versus >14 days (aRR = 1.11, 95% CI = 0.81–1.52). There was no significant difference in all secondary adverse outcomes between the two groups. Subgroup analyses, limited to people diagnosed with GDM (N = 89, 23.4%), also found similar results as the full analyses.

Keywordsgestational diabetes

- screening
- glucose tolerance test
- large for gestational age

Conclusion Among individuals who completed the two-step screening for GDM, completion of the 3-hour GTT within \leq 14 versus \geq 14 days was not associated with an increase rate of the adverse outcomes.

received January 24, 2023 accepted after revision July 31, 2023 accepted manuscript online August 1, 2023 article published online September 15, 2023 © 2023. Thieme. All rights reserved. Thieme Medical Publishers, Inc., 333 Seventh Avenue, 18th Floor, New York, NY 10001, USA DOI https://doi.org/ 10.1055/a-2145-7899. ISSN 0735-1631.

Key Points

- Among pregnant people in an academic practice, 50% of people with abnormal 1-hour GTT completed GDM two-step screening in 14 days.
- Longer length of time to completion of diagnostic testing for GDM was not associated with an increased rate of adverse outcomes.
- Pregnant people that were diagnosed with GDM and completed the two-step method in >14 days did not have worse perinatal outcomes.

Gestational diabetes mellitus (GDM), defined as carbohydrate intolerance during pregnancy, affects up to 10% of all pregnancies in the United States.¹ GDM is associated with maternal and fetal complications such as hypertensive disorders, cesarean section, large for gestational age, shoulder dystocia, neonatal hypoglycemia, and neonatal intensive care unit (NICU admissions).^{2–6} Moreover, pregnant people diagnosed with GDM are at up to 60% risk of developing type 2 diabetes later in life.^{7–9} Timely diagnosis and treatment of GDM improves short-term maternal and fetal outcomes.^{2,3,10,11}

To diagnose GDM, American College of Obstetrics and Gynecologists (ACOG) recommends the two-step method, which entails completion of the screening 1-hour glucose challenge test (GCT); if elevated (\geq 135–140 mg/dL), the completion of the 3-hour glucose tolerance test (GTT) is required for diagnosis.^{12–14} Approximately, 90% of maternal–fetal medicine specialists recommend diagnosis with the two-step method.¹⁵ Unfortunately, the diagnostic 3-hour GTT requires a fasting state and a lengthy clinic visit, which may lead to a delay in testing, diagnosis, and ultimately intervention for GDM.¹⁶

There are no current guidelines on the optimal length of time to completion of the two-step method for GDM screening. In addition, there is a lack of data in examining the relationship between length of time to completion of the GDM, two-step screening method, and adverse outcomes. The aim of our study was to determine if the length of time to completion of the two-step method GDM screening was associated with adverse neonatal and maternal outcomes.

Materials and Methods

This was a retrospective cohort study taken place at the UT Health (University of Texas Health Science Center at Houston) McGovern Medical School, an academic, tertiary care system in Houston, TX from May 2021 to May 2022. An electronic medical record chart review was conducted to identify all singleton, nonanomalous pregnant people who were screened for GDM at \geq 24 weeks' gestation. Exclusion criteria included: people with pregestational diabetes, normal 50-g 1-hour GCT (<135 mg/dL) or individuals that did not complete 3-hour GTT. Pregnant people with 1-hour GCT \geq 135 mg/dL were considered positive screening based on an institutional threshold. After completion of the diagnostic 100-g 3-hour GTT, diagnosis of GDM was based on Carpenter–Coustan criteria of ≥ 2 abnormal values (fasting blood glucose >90 mg/dL, 1 hour > 180 mg/dL, 2 hour > 155 mg/dL, 3 hour > 140 mg/dL)¹. We compared outcomes among those who completed the 3hour GTT in ≤ 14 versus >14 days from the 1-hour GCT, irrespective of whether they had GDM.

The study was approved by the Institutional Review Board at UTHealth McGovern Medical School (IRB no.: HSC-MS-22-0292). Data were collected from the clinic and hospital medical record system. Maternal records were culled for prenatal visits, including laboratory values, diagnosis of GDM, need for hypoglycemic agents, and ultrasounds. Hospital records were abstracted for admissions during pregnancy, labor, delivery, and postpartum events. Neonatal records were also accessed for birth weight, Apgar scores, length of hospital stay, need for NICU admission, or any other neonatal complications after delivery.

The explanatory variable in this study was time to completion of the 3-hour GTT from the 1-hour GCT, categorized as \leq 14 versus > 14 days. As there is no recommended time to completion of two-step GDM screening, the authors arbitrarily designated 14 days to completion as the cutoff between the two groups. The hypothesis was that 14 days is a clinically practical amount of time to allow for completion of screening as well as initiation of GDM care, if required, and greater than 14 days may impact perinatal outcomes.

The primary outcome was a composite neonatal morbidity and mortality (CNM) consisting of any of the following: large for gestational age (defined as birth weight > 90% for gestational age per Duryea et al nomogram),¹⁷ shoulder dystocia or birth injury, respiratory distress, neonatal hypoglycemia, or fetal or neonatal death. Shoulder dystocia or birth injury was defined as additional maneuvers other than gentle downward traction for delivery. Birth injury included clavicular fracture or brachial plexus injury. Respiratory distress was defined as the need for at least 4 hours of respiratory support with supplemental oxygen, continuous positive airway pressure, or ventilation at the first 24 hours of life. Neonatal hypoglycemia was defined as blood glucose < 40 mg/dL in the first 24 hours of life or <50 mg/dL after or requiring medical therapy.

The secondary adverse neonatal outcomes included: rates of preterm delivery, Apgar score <7 at 5 minutes, need for continuous positive airway pressure ventilation neonatal jaundice requiring phototherapy, and length of hospital stay. Secondary adverse maternal outcomes that were collected included: hypertensive disorders of pregnancy, chorioamnionitis, cesarean delivery (CD), postpartum hemorrhage, endometritis, or postpartum readmission \leq 6 weeks. Preterm birth was defined as delivery at <37 weeks' gestation. Hypertensive disorder of pregnancy included gestational hypertension, preeclampsia, or superimposed preeclampsia based on the

current Task Force Criteria for Hypertension in pregnancy.¹⁸ Gestational hypertension defined as new onset hypertension (systolic \geq 140 mm Hg or diastolic \geq 90 mm Hg) without proteinuria (either \geq 300 mg per 24 hours or protein/Cr ratio \geq 0.3) after 20 weeks of gestation. Preeclampsia defined as hypertension (systolic \geq 140 mm Hg or diastolic \geq 90 mm Hg) with proteinuria or serum laboratory abnormalities (platelets \leq 100,000, serum aspartate aminotransferase \geq 80 IU/mL, creatinine \geq 1.1 mg/dL). Preeclampsia with severe features defined as systolic \geq 160 mm Hg, diastolic \geq 110 mm Hg, persistent headache, pulmonary edema, or any serum laboratory abnormalities as above.

Statistical Analysis

We examined the differences in baseline characteristics and outcomes between time groups (completion of 3-hour GTT in \leq 14 versus > 14 days from abnormal 1-hour GCT) using the chi-square test or Fischer's exact test for categorical variables and Student's t-tests for continuous variables. Multivariable Poisson regression models were used to evaluate the association between time groups (\leq 14 days [reference] vs. >14 days) and the primary outcome, while adjusting for maternal age ($<20, 20-34, \geq 35$ years), maternal race, and ethnicity (non-Hispanic White, non-Hispanic Black, Hispanic, non-Hispanic-Other, unknown), private insurance (private, government-issued insurance or no insurance), nulliparity, obesity (body mass index [BMI] \geq 30 kg/m²), hypertensive disorder, thyroid disease and gestational age at time of 1-hour GCT. The results were presented as adjusted relative risk (aRR) with 95% confidence interval (CI). Using the same approach, we also performed subgroup analyses among those diagnosed with GDM. Statistical significance was noted to be a p-value of < 0.05 or if the 95% CI did not include the integer 1. Statistical

analysis was completed using STATA software, version 17 (Stata-Corp., College Station, TX).

Results

Among the 401 pregnancies with 1-hour GCT \geq 135 mg/dL during the study period, 342 (85.3%) pregnant people had 3-hour GTT ordered. A total of 59 individuals did not have 3-hour GTT ordered due to the following reasons: GDM was diagnosed based on elevated 1-hour GCT (N=25) and 3-hour GTT not ordered by primary obstetrician (N=34). A total of 29 people did not complete the ordered 3-hour GTT.

The final study sample included 313 (78.1%) pregnant people who completed the 3-hour GTT. Among our study population, 171 (54.6%) completed the two-step method in \leq 14 days and 142 (45.4%) completed >14 days (**~Fig. 1**).

Baseline characteristics for the two groups were similar except for insurance status (\neg Table 1). Those who completed the 3-hour GTT >14 days were more likely to be insured with Medicaid (\leq 14 days 45% vs. >14 days 74%, p = 0.004).

Overall, the rate of the primary outcome was 28.8%; 25.7% in the group of \leq 14 days, whereas 32.4% in the group of >14 days (p = 0.195). The most common component of the primary outcome was neonatal hypoglycemia (15.7%; 14.0% in \leq 14 days and 17.6% >14 days, p = 0.387). After multivariable adjustment, the risk of the primary outcome was similar between people who completed the two-step method in \leq 14 days versus >14 days (aRR = 1.11; 95% CI = 0.81–1.52,). There was no difference in any component of the primary outcome between the two groups (**¬Table 2**).

Table 3 presents the secondary adverse neonatal and maternal outcomes. The results showed that there was no



Fig. 1 Flow chart of pregnant people: eligibility and sample size. GDM, gestational diabetes; 1hGCT, 1-hour glucose challenge test; 3hGCT, 3-hour glucose tolerance test.

Table 1 Baseline characteristics				
Characteristics	Two-step completed		p-Value	
	≤14 d	>14 d		
	n = 171 (%)	n = 142 (%)		
Maternal age (y)				
< 20	5 (2.9)	6 (4.2)	0.550	
20–34	134 (78)	104 (73)		
≥35	32 (18.7)	32 (22.5)		
Race/ethnicity				
Non-Hispanic White	47 (27)	28 (20)	0.100	
Non-Hispanic Black	20 (12)	29 (20)		
Hispanic	27 (16)	21 (15)		
Other	69 (40)	52 (37)		
Unknown	8 (4.7)	12 (8.5)		
Insurance status				
Private	94 (55.0)	62 (44)	0.004	
Medicaid	77 (45.0)	74 (52.1)		
Self-pay/no insurance	0 (0.0)	6 (4.2)		
Nulliparous	27 (15.8)	27 (19.0)	0.452	
Obese (BMI \geq 30 kg/m ²)	85 (49.7)	67 (47.2)	0.656	
Hypertension	13 (7.6)	12 (8.5)	0.783	
Thyroid disease	12 (7.0)	8 (5.6)	0.618	
Substance use (tobacco or illicit drug)				
Yes	5 (2.9)	5 (3.5)	0.496	
No	150 (87.7)	118 (83.1)		
Unknown	16 (9.4)	19 (13.4)		
GA at screening 1 h GCT	26.4 (2.0)	26.7 (2.1)	0.321	
1 h GCT	157.0 (16.7)	155.4 (15.8)	0.395	
1 abnormal value of 3 h GTT ^a	73.0 (42.7)	50.0 (35.2)	0.177	
Diagnosis of GDM	48.0 (28.1)	41.0 (28.9)	0.875	
Diabetes educator visit	42.0 (24.6)	36.0 (25.4)	0.872	
Need for hypoglycemic agent	23.0 (13.5)	11.0 (7.7)	0.106	

Abbreviations: BMI, body mass index; CGM, continuous glucose monitoring; GA, gestational age; GCT, glucose challenge test; GTT, glucose tolerance test.

Note: Data are presented as number (percentage) or mean (standard deviation).

^aGTT defined as abnormal based on Carpenter–Coustan criteria of \geq 2 abnormal values: fasting blood glucose > 90, 1 hour > 180, 2 hours > 155, 3 hours > 140 mg/dL.

difference in all the secondary adverse outcomes between the time groups. The most common maternal complication was CD (40.4% in \leq 14 days and 43.7% in >14 days, p = 0.554; **-Table 3**).

There were a total of 89 (28.4%) pregnant people that were diagnosed with GDM. Of them, 48 (53.9%) people completed the two-step method in \leq 14 days and 41 (46.1%) people completed in >14 days. Subgroup analysis showed there was no significant difference in any component of the CNM between the two groups, similar to the primary analysis (aRR = 1.01; 95% CI = 0.59–1.78). The most common CNM in both groups was neonatal hypoglycemia (29.2% in \leq 14 days and 29.3% >14 days). In addition, we did not find any

differences in all secondary neonatal and maternal outcomes in those diagnosed with GDM (**-Table 4**).

Discussion

In this retrospective cohort study of the length of time to completion of two-step method for diagnosis of GDM, there was no association with worse adverse neonatal or maternal outcomes in people with longer period of time to completion of testing. In our study, we observed that approximately 50% of people that received care at our institution completed the two tests within 14 days, while 67% of people completed the two-step method in \leq 21 days.

Table 2 The primary outcome: composite neonatal adverse outcome					
	Two-step method		p-Value	Adjusted RR	
	Completed in \leq 14 d	Completed in $>$ 14 d		(95% CI)	
	n = 171 (%)	n = 142 (%)			
Composite neonatal outcome	44 (25.7)	46 (32.4)	0.20	1.21 (0.85–1.73)	
Large for gestational age ^a	14 (8.2)	14 (9.9)	0.61		
Shoulder dystocia or birth injury ^b	5 (2.9)	5 (3.5)	0.77		
Respiratory distress ^c	8 (4.7)	14 (9.9)	0.07		
Hypoglycemia ^d	24 (14.0)	25 (17.6)	0.39		
Fetal or neonatal death	0 (0)	0 (0)	N/C		

Abbreviations: CI, confidence interval; CPAP, continuous positive airway pressure; N/C, not calculable; RR, relative risk.

Notes: Data are presented as number (percentage) or median (standard deviation). Adjusted for insurance status.

^aBirth weight above 90th percentile using the nomogram by Duryea et al.

^bNeed for any extra maneuvers for delivery, clavicular fracture, or brachial plexus injury.

^cNeed of at least 4 hours of respiratory support with supplemental oxygen, continuous positive airway pressure, or ventilation at the first 24 hours of life.

^dBlood glucose <40 mg/dL in the first 24 hours of life or <50 mg/dL after or requiring medical therapy.

Table 3 The secondary neonatal and maternal adverse outcome					
Outcomes	Two-step method	p-Value			
	Completed in \leq 14 d	Completed in $>$ 14 d			
	n = 171 (%)	n = 142 (%)			
Neonatal adverse outcomes					
Preterm delivery (<37 wk)	23 (13.5)	26 (18.3)	0.239		
Birth weight (g)	3,214.4 (±467.0)	3,149.1 (±566.6)	0.265		
Apgar score <7 at 5 min	1 (0.6)	0 (0.0)	1.000		
СРАР	11 (6.4)	11 (7.7)	0.651		
Neonatal jaundice requiring phototherapy	25 (14.6)	16 (11.3)	0.381		
Length of hospital stay (d)	2.6 (±2.8)	3.4 (±6.2)	0.118		
Maternal adverse outcomes					
Hypertensive disorders of pregnancy ^a	39 (24.9)	37 (28.2)	0.474		
Preeclampsia with severe features	11 (6.9)	6 (4.6)	0.444		
Chorioamnionitis	6 (3.5)	10 (7.0)	0.158		
Cesarean delivery	69 (40.4)	62 (43.7)	0.554		
Postpartum hemorrhage	9 (5.3)	7 (4.9)	0.894		
Endometritis	2 (1.2)	1(0.7)	1.000		
Postpartum readmission \leq 6 wk	7 (4.3)	4 (2.9)	0.557		

Abbreviation: CPAP, continuous positive airway pressure.

Note: Data are presented as number (percentage) or mean (standard deviation).

^aHypertensive disorder of pregnancy including gestational hypertension, preeclampsia, or superimposed preeclampsia.

In addition, subgroup analyses of those that were diagnosed with GDM did not show a significantly higher rate of adverse outcomes among those that took >14 days to diagnosis. Majority of people were able to be diagnosed with GDM in less than a month's time allowing for appropriate intervention.

There have been no prior studies examining length of time to completion of GDM screening and neonatal and maternal morbidity. However, many studies have explored the barriers to completion of the 1-hour GCT. Several social and institutional factors have been identified: inability to tolerate test, compliance with multiple prenatal appointments, younger maternal age, and mental/social stressors.^{19,20} These prior reports are consistent with our findings of those who completed the GDM screening in \leq 14 days had a significantly higher rate of private insurance as compared with >14 days to completion (p = 0.004). Although we did not find a significant difference in the primary outcome based on the length of time to completion of the two-step method of GDM screening, we detected a higher-than-average rate of neonatal hypoglycemia. Prior research reported that up to 10% of low risk, term newborns experience

Table 4 Subgroup analyses: the primary and secondary outcomes in those diagnosed with gestational diabetes mellitus				
	Two-step method $Completed in > 14 d$		p-Value	Adjusted RR (95% CI)
	n = 48 (%)	n = 41 (%)		
Composite neonatal outcome	18 (37.5)	17 (41.5)	0.703	1.01 (0.59–1.78)
Large for gestational age ^a	4 (8.3)	4 (9.8)	1.000	
Shoulder dystocia or birth injury ^b	1 (2.1)	2 (4.9)	0.593	
Respiratory distress ^c	1 (2.1)	4 (9.8)	0.176	
Hypoglycemia ^d	14 (29.2)	12 (29.3)	0.992	
Fetal or neonatal death	0 (0.0)	0 (0.0)	N/C	
Secondary neonatal outcomes	5 (10.4)	10 (24.4)	0.17	
Preterm delivery (<37 wk)	5 (10.4)	10 (24.4)	0.079	
Birth weight (g)	3,221.7 (±452.3)	3,114.3 (±612.7)	0.345	
Apgar score <7 at 5 min	1 (2.1)	0 (0.0)	1.000	
СРАР	3 (6.3)	4 (9.8)	0.699	
Neonatal jaundice requiring phototherapy	2 (4.2)	5 (12.2)	0.241	
Length of hospital stay (d)	2.3 (±1.5)	3.9 (±5.9)	0.079	
Secondary maternal outcomes				
Hypertensive disorders of pregnancy ^e	7 (14.9)	12 (29.3)	0.102	
Chorioamnionitis	1 (2.1)	5 (12.2)	0.091	
Cesarean delivery	22 (45.8)	25 (61.0)	0.154	
Postpartum hemorrhage	3 (6.3)	3 (7.3)	1.000	
Endometritis	0 (0.0)	0 (0.0)	N/C	
Postpartum readmission \leq 6 wk	4 (8.7)	2 (5.3)	0.685	

Abbreviations: CI, CPAP, continuous positive airway pressure; N/C, not calculable; RR, relative risk.

Notes: Data are presented as number (percentage) or median (standard deviation). Adjusted for insurance status.

^aBirth weight above 90th percentile using the nomogram by Duryea et al.

^bNeed for any extra maneuvers for delivery, clavicular fracture or brachial plexus injury.

^cNeed of at least 4 hours of respiratory support with supplemental oxygen, continuous positive airway pressure, or ventilation at the first 24 hours of life.

 d Blood glucose <40 mg/dL in the first 24 hours of life or <50 mg/dL after or requiring medical therapy.

^eHypertensive disorder of pregnancy including gestational hypertension, preeclampsia, or superimposed preeclampsia.

hypoglycemia defined as <40 to 45 mg/dL.^{21,22} In our study, however, the rate of the neonatal hypoglycemia was 15.7%. The increased rate may be due to high-risk pregnancies (those with GDM diagnosis or 48.5% of patients with BMI \geq 30) being included in our study population. This also reflects that hyperglycemia, below the threshold of diagnosis, is associated with worse perinatal outcomes consistent with the Landon et al findings study.² Despite completing and passing the two-step screening method, poor neonatal outcomes that are associated with GDM were observed. Future studies should focus on exploring other risk factors for neonatal hypoglycemia and whether the current method for GDM diagnosis in pregnancy is sufficient for diagnosis and treatment of people with elevated blood sugars during pregnancy.

Limitations and Strengths

We acknowledge the limitations of this study. It was a single-center, retrospective analyses and was specific to a university practice. Causes of delay might be due to several and all factors of social determinants of health, reflecting both a clinic system's issue as well as disparities in access to health care.^{19,20} We performed a multivariable adjustment analysis on the primary outcome of the composite neonatal adverse outcome. However, this analysis was not completed on the secondary outcomes due to small case numbers, and the bivariate analyses showed that there were no significant results in all the secondary outcomes. Another limitation was the risk of Type 2 error given small cohort sample; a significant difference between the two groups may have been observed with a larger sample size, given the high rate of neonatal hypoglycemia.

Our study has several strengths. To the best of our knowledge, this is the first study that has explored neonatal and maternal outcomes associated with length of time to completion of GDM screening. Although the study population was from a single center, our sample included people from a greater metropolitan area. In addition, our overall rate of GDM detection was approximately 6%, consistent with the overall national average.^{5,13,23–25}

Conclusion

In conclusion, we did not find a difference in adverse neonatal and maternal outcomes associated with the length of time to completion of the two-step GDM diagnosing method. This study provides a clinically applicable quantification of the delay that happens almost 50% of the time in the journey to diagnosis of GDM. This highlights that completion of the two-step method is fraught with challenges and a more practical form of GDM screening is needed for the ease of pregnant people.

Funding

None.

Conflict of Interest

None declared.

References

- 1 Gregory E, Ely DM. Trends and Characteristics in Gestational Diabetes: United States, 2016–2020, in National Vital Statistics Reports, Vol. 71. 2022
- 2 Landon MB, Spong CY, Thom E, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. A multicenter, randomized trial of treatment for mild gestational diabetes. N Engl J Med 2009;361(14):1339–1348
- ³ Crowther CA, Hiller JE, Moss JR, McPhee AJ, Jeffries WS, Robinson JSAustralian Carbohydrate Intolerance Study in Pregnant Women (ACHOIS) Trial Group. Effect of treatment of gestational diabetes mellitus on pregnancy outcomes. N Engl J Med 2005;352(24): 2477–2486
- 4 Hartling L, Dryden DM, Guthrie A, Muise M, Vandermeer B, Donovan L. Benefits and harms of treating gestational diabetes mellitus: a systematic review and meta-analysis for the U.S. Preventive Services Task Force and the National Institutes of Health Office of Medical Applications of Research. Ann Intern Med 2013;159(02):123–129
- 5 Shah NS, Wang MC, Freaney PM, et al. Trends in gestational diabetes at first live birth by race and ethnicity in the US, 2011-2019. JAMA 2021;326(07):660–669
- 6 Metzger BE, Lowe LP, Dyer AR, et al; HAPO Study Cooperative Research Group. Hyperglycemia and adverse pregnancy outcomes. N Engl J Med 2008;358(19):1991–2002
- 7 Bellamy L, Casas JP, Hingorani AD, Williams D. Type 2 diabetes mellitus after gestational diabetes: a systematic review and metaanalysis. Lancet 2009;373(9677):1773–1779
- 8 Kim C, Newton KM, Knopp RH. Gestational diabetes and the incidence of type 2 diabetes: a systematic review. Diabetes Care 2002;25(10):1862–1868

- 9 Bochkur Dratver MA, et al. Longitudinal changes in glucose during pregnancy in women with gestational diabetes risk factors. Diabetologia 2021
- 10 Association ADAmerican Diabetes Association. 14. Management of diabetes in pregnancy: standards of medical care in diabetes-2020.. Diabetes Care 2020;43(Suppl 1):S183–S192
- 11 Sohn J, Lim HJ, Kim S, et al. Delayed diagnosis of gestational diabetes mellitus and perinatal outcomes in women with large for gestational age fetuses during the third trimester. Obstet Gynecol Sci 2020;63(05):615–622
- 12 Practice Bulletin No. Practice bulletin no. 180: gestational diabetes mellitus. Obstet Gynecol 2017;130(01):e17–e37
- 13 Moyer VAU.S. Preventive Services Task Force. Screening for gestational diabetes mellitus: U.S. Preventive Services Task Force recommendation statement. Ann Intern Med 2014;160(06):414–420
- 14 Harper LM, Mele L, Landon MB, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) Maternal-Fetal Medicine Units (MFMU) Network. Carpenter-Coustan compared with national diabetes data group criteria for diagnosing gestational diabetes. Obstet Gynecol 2016;127(05):893–898
- 15 Bimson BE, Rosenn BM, Morris SA, Sasso EB, Schwartz RA, Brustman LE. Current trends in the diagnosis and management of gestational diabetes mellitus in the United States. J Matern Fetal Neonatal Med 2017;30(21):2607–2612
- 16 Agarwal MM. Gestational diabetes mellitus: screening with fasting plasma glucose. World J Diabetes 2016;7(14):279–289
- 17 Duryea EL, Hawkins JS, McIntire DD, Casey BM, Leveno KJ. A revised birth weight reference for the United States. Obstet Gynecol 2014;124(01):16–22
- 18 Gestational Hypertension and Preeclampsia. Gestational hypertension and preeclampsia: ACOG practice bulletin summary, number 222. Obstet Gynecol 2020;135(06):1492–1495
- 19 Nielsen KK, Kapur A, Damm P, de Courten M, Bygbjerg IC. From screening to postpartum follow-up - the determinants and barriers for gestational diabetes mellitus (GDM) services, a systematic review. BMC Pregnancy Childbirth 2014;14:41
- 20 Lachmann EH, Fox RA, Dennison RA, Usher-Smith JA, Meek CL, Aiken CE. Barriers to completing oral glucose tolerance testing in women at risk of gestational diabetes. Diabet Med 2020;37(09):1482–1489
- 21 Edwards T, Harding JE. Clinical aspects of neonatal hypoglycemia: a mini review. Front Pediatr 2021;8:562251
- 22 Braun D, Braun E, Chiu V, et al. Trends in neonatal intensive care unit utilization in a large integrated health care system. JAMA Netw Open 2020;3(06):e205239
- 23 ACOG Practice Bulletin No. ACOG practice bulletin no. 190: gestational diabetes mellitus. Obstet Gynecol 2018;131(02):e49–e64
- 24 Hamilton BE, Martin JA, Osterman MJK. Births: provisional data for 2021. Vital Statistics Rapid Release no 20. Hyattsville, MD: National Center for Health Statistics. ; May 2022
- 25 Vandorsten JP, Dodson WC, Espeland MA, et al. NIH consensus development conference: diagnosing gestational diabetes mellitus. NIH Consens State Sci Statements 2013;29(01):1–31