


A light in the dark: Peroral cholangioscopy provides a new strategy for difficult cannulation in pancreaticobiliary maljunction

OPEN
ACCESS




► **Fig. 1** Magnetic resonance cholangiopancreatography showed the confluence between dilated biliary and pancreatic ducts.



► **Fig. 2** Pancreatic duct opacification when conducting cholangiography.

A 56-year-old man suffered from epigastric pain for 5 days with elevated amylase (2600 IU/L), and computed tomography indicated acute pancreatitis. Magnetic resonance cholangiopancreatography showed the confluence between dilated biliary and pancreatic ducts (► **Fig. 1**), and endoscopic ultrasound indicated a 1.7-cm common channel, suggesting a pancreaticobiliary maljunction (PBM). Subsequent endoscopic retrograde cholangiopancreatography (ERCP) showed pancreatic duct opacification when conducting cholan-



► **Fig. 3** **a** The guidewire could not be inserted into the pancreatic duct owing to the long common channel. **b** The peroral choledochoscope was used to observe the opening of pancreatic duct and insert the guidewire under direct visualization.



► **Video 1** Peroral cholangioscopy-assisted pancreatic duct cannulation in a patient with a pancreaticobiliary maljunction.

giography (► **Fig. 2**). Considering the symptoms of acute pancreatitis and high amylase, we attempted stent implantation to conduct pancreatic juice drainage and relieve the abdominal pain. However, owing to the long common channel and the sharp angle, the guidewire could not be inserted into the pancreatic duct during prior attempts (► **Fig. 3 a**). Therefore, a novel peroral

choledochoscope (Eye-Max CDS11001, 9Fr; Micro-Tech, Nanjing, China) with a 1.8-mm working channel was used to observe the opening of pancreatic duct under direct visualization (► **Fig. 3 b**, ► **Video 1**). The fusion junction was presented clearly inside the field (► **Fig. 4**), and the guidewire was inserted into the pancreatic duct successfully with the assistance of the peroral choledocho-



► **Fig. 4** The fusion junction visible with the help of a peroral choledochoscope.

scope. After pancreatic duct stent implantation, the amylase gradually decreased and the patient's recovery was uneventful during 5-month follow-up. Peroral cholangioscopy has been widely applied in diagnosing pancreatobiliary diseases and shown its vital role in selective cannulation of complex biliary strictures [1]. PBM is a rare congenital malformation with the pancreatic and bile ducts united outside of the duodenal wall, resulting in dysfunction of the sphincter of Oddi and regurgitation of bile and pancreatic juice [2,3]. Endoscopic pancreatic duct stenting is an effective way to relieve the symptoms, although the diversity of anatomic variation sometimes makes cannulation difficult [4]. To our knowledge, this is the first report that applies peroral cholangioscopy in pancreatic duct cannulation in patients with PBM. Peroral cholangioscopy makes the fusion junction visible and provides a new strategy for difficult cannulation in PBM.

Endoscopy_UCTN_Code_TTT_1AR_2AI


Funding

Natural Science Foundation of Shandong Province, China
ZR2020LZL003

Competing interests

The authors declare that they have no conflict of interest.

The authors

Wen-Lin Zhang^{1,2,3}, Ning Zhong^{1,2,3},
Rui Ji^{1,2,3} 

- 1 Department of Gastroenterology, Qilu Hospital, Cheeloo College of Medicine, Shandong University, Jinan, Shandong Province, China
- 2 Laboratory of Translational Gastroenterology, Qilu Hospital, Cheeloo College of Medicine, Shandong University, Jinan, Shandong Province, China
- 3 Robot Engineering Laboratory for Precise Diagnosis and Therapy of GI Tumor, Qilu Hospital of Shandong University, Jinan, Shandong, China

Corresponding author

Rui Ji, MD

Qilu Hospital of Shandong University, 107 Wenhuxi Road, 250012 Jinan, China
qljrui@email.sdu.edu.cn

References

- [1] Kastelijin JB, Didden P, Bogte A et al. Digital single-operator cholangioscopy to guide selective cannulation of complex biliary strictures. *Surg Endosc* 2022; 36: 9476–9480. doi:10.1007/s00464-022-09665-1
- [2] Kamisawa T, Takuma K, Anjiki H et al. Pancreaticobiliary maljunction. *Clin Gastroenterol Hepatol* 2009; 7: S84–88. doi:10.1016/j.cgh.2009.08.024
- [3] Ono A, Arizono S, Isoda H et al. Imaging of pancreaticobiliary maljunction. *RadioGraphics* 2020; 40: 378–392. doi:10.1148/rg.2020190108
- [4] Jin Z, Bie LK, Tang YP et al. Endoscopic therapy for patients with pancreaticobiliary maljunction: a follow-up study. *Oncotarget* 2017; 8: 44860–44869. doi:10.18632/oncotarget.16228

Citation Format

Endoscopy Endoscopy 2023; 55: E792–E793. doi: 10.1055/a-2096-1950.

Bibliography

Endoscopy 2023; 55: 968–969

DOI 10.1055/a-2154-3461

ISSN 0013-726X

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited.

(<https://creativecommons.org/licenses/by/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



E-Videos is an open access online section of the journal *Endoscopy*, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high-quality video and are published with a Creative Commons CC-BY license. Endoscopy E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: <https://www.research4life.org/access/eligibility/>).

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>