



Contrast-enhanced endoscopic ultrasound and fine-needle biopsy of a rare mediastinal mass: a mediastinal schwannoma

OPEN
ACCESS



► **Fig. 1** Computed tomography 3D reconstruction with a nodular mass of approximately 7 cm in diameter in the left apical pleura with compression of the subclavian artery.



► **Video 1** Contrast-enhanced endoscopic ultrasonography examination and fine-needle biopsy of a mediastinal schwannoma.




Contrast-enhanced endoscopic ultrasonography-guided fine-needle biopsy (CE-EUS-FNB) is an important, minimally invasive tool for the diagnosis of mediastinal masses. The specimen obtained allows a wider range of analyses to be performed and a better description of morphology and immunophenotype compared with fine-needle aspiration (FNA). The main advantages of CE-EUS include real-time imaging of microvasculature and microperfusion, and impressively good detail resolution [1].

We present the case of a 20-year-old woman with Horner syndrome. On computed tomography scan, a nodular mass of approximately 7 cm in diameter in the left apical pleura was described, with compression of both the pulmonary parenchyma and subclavian artery (► **Fig. 1**). After multidisciplinary board discussion, it was decided that tissue sampling should be performed for histological definition (► **Video 1**).

We performed CE-EUS, which revealed a 7-cm hypoechoic mass in close apposition to the aortic arch, left subclavian artery, and left common carotid artery (► **Fig. 2**), with diffuse and inhomogeneous hypo-enhancement following



► **Fig. 2** Endoscopic ultrasound view of the lesion showing a 7-cm hypoechoic mass.




► **Fig. 3** Endoscopic ultrasound view of the lesion showing a diffuse and inhomogeneous hypo-enhancing pattern following contrast injection.

contrast injection (► **Fig. 3**) (Sonovue; Bracco, Milan, Italy). A transesophageal EUS-FNB was performed with a 22-G needle (SharkCore; Medtronic, Minneapolis, Minnesota, USA). Our pathologists defined the tumor as a benign peripheral nerve sheath tumor, most compatible with schwannoma, thanks to the features described by the wide range of immunohistochemical stains (positivity for S100, GFAP, SOX10, D2.40, and negativity for CAM5.2, CD117, MelanA, CD34) (► **Fig. 4**). Thus, the patient underwent a

thoracoscopic mass resection, with a regular postoperative course. The surgical specimen (► **Fig. 5**) confirmed the final diagnosis of schwannoma.

Mediastinal schwannoma is a rare mediastinal mass [2], and only a single case of EUS-FNA cytological diagnosis has been reported [3]. To the best of our knowledge, neither contrast-enhancement behavior nor endoscopic FNB has been described for this rare mediastinal lesion. Nevertheless, the histological specimen



► **Fig. 4** Tumor cells demonstrated strong and diffuse expression of S100 (original magnification $\times 3$).

5 Department of Surgical, Oncological and Oral Sciences, University of Palermo, Palermo, Italy

Corresponding author

Dario Quintini, MD

Department of Surgical Oncological and Gastroenterology Sciences, University of Padua, Via Nicolò Giustiniani, 2, 35128 Padova, PD Padova 35122, Italy
quintinidario@gmail.com

References

- [1] Dietrich CF, Sharma M, Hocke M. Contrast-enhanced endoscopic ultrasound. *Endosc ultrasound* 2012; 1: 130–136
- [2] Kapoor A, Singhal MK, Narayan S et al. Mediastinal schwannoma: a clinical, pathologic, and imaging review. *South Asian J Cancer* 2015; 4: 104–105
- [3] McGrath KM, Ballo MS, Jowell PS. Schwannoma of the mediastinum diagnosed by EUS-guided fine needle aspiration. *Gastrointest Endosc* 2001; 53: 362–365

Bibliography

Endoscopy 2023; 55: E1071–E1072

DOI 10.1055/a-2163-1544

ISSN 0013-726X

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited.

(<https://creativecommons.org/licenses/by/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



► **Fig. 5** Macroscopic appearance of the resected tumor in sections, with smooth surface and firm, light-tan tissue with focal patches of hemorrhage.

acquired by EUS-FNB allows for the use of a wider range of immunohistochemical stains, increasing the specificity of diagnosis. Furthermore, contrast-enhanced evaluation offers real-time guidance for EUS-FNB and is likely to increase sensitivity.

Endoscopy_UCTN_Code_CCL_1AF_2AC

Funding

Ministero della Salute
<http://dx.doi.org/10.13039/501100003196>
Ricerca corrente 2023

Competing interests

The authors declare that they have no conflict of interest.

The authors

Dario Ligresti¹, **Dario Quintini**^{1,2}, **Ilaria Tarantino**¹, **Alessandro Bertani**³, **Agita Jukna**⁴, **Giacomo Emanuele Maria Rizzo**^{1,5}, **Mario Traina**¹

1 Endoscopy Service, Department of Diagnostic and Therapeutic Services, IRCCS – ISMETT, Palermo, Italy

2 Department of Surgical Oncological and Gastroenterology Sciences, University of Padua, Padua, Italy

3 Unit of Thoracic Surgery and Lung Transplantation, Department for the Treatment and Study of Cardiothoracic Diseases and Cardiothoracic Transplantation, IRCCS – ISMETT, Palermo, Italy

4 Pathology Unit, Department of Diagnostic and Therapeutic Services, IRCCS – ISMETT, Palermo, Italy