# Endoscopic ultrasound-guided caudate lobe liver abscess drainage





► Fig. 1 Computed tomography showing a 4.5×5.1-cm abscess in the caudate lobe of the liver.

Endoscopic ultrasound (EUS)-guided liver abscess drainage is an alternative method in cases where percutaneous drainage (PCD) is difficult to perform [1,2]. However, reports on EUS-guided caudate lobe liver abscess drainage remain scarce, with only 12 reported cases [3]. Of these cases, 33% (4/12) were treated with a transesophageal approach, which can cause mediastinitis, mediastinal emphysema, and pneumothorax [4,5]. Herein, we present a case of safe and successful drainage via this method using marking clips and adjusting the scope position.

A 50-year-old man with type 2 diabetes mellitus was hospitalized for fever and epigastralgia evolving for 4 days. Computed tomography (CT) revealed a 4.5× 5.1-cm abscess in the liver caudate lobe (**Fig.1**). PCD was not performed because of poor visualization on abdominal ultrasonography. After 2 weeks of antibiotic treatment with little improvement, he was referred to our hospital, where EUS-guided caudate lobe liver abscess drainage was performed (**> Video 1**).

First, a forward-viewing endoscope was used to mark the esophagogastric junction with a clip so that its location could be determined under fluoroscopy. Subsequently, an echoendoscope was introduced into the stomach. By advancing the scope further from the clip and applying an upward angle, the abscess was successfully visualized from within the stom-



**Video 1** Endoscopic ultrasound-guided caudate lobe liver abscess drainage.



▶ Fig. 2 A 7-Fr double-pigtail plastic stent and a 6-Fr naso-abscess tube are visible in the abscess cavity.

ach. The abscess was punctured using a 19-gauge needle with fluoroscopic confirmation of the marking clip, and a 0.025-inch guidewire was inserted into the abscess cavity. Another 0.035-inch guidewire was placed in the abscess cavity using a double-lumen catheter. After dilation using an electrocautery dilator, a 7-Fr double-pigtail plastic stent and a 6-Fr naso-abscess tube were placed (**> Fig. 2**). *Klebsiella pneumoniae* was cultured from the abscess contents (**> Fig. 3**). The nasal tube was removed, and the patient was



**Fig. 3** Red-white pus is aspirated through the naso-abscess tube.

discharged on day 10 without any complications. The plastic stent was removed 3 months later without liver abscess recurrence.

Endoscopy\_UCTN\_Code\_TTT\_1AS\_2AD

The authors declare that they have no conflict of interest.

# The authors

Junichi Kaneko<sup>®</sup>, Hiroki Tamakoshi, Moeka Watahiki, Daisuke Kusama, Tomoyuki Niwa, Masaki Takinami<sup>®</sup>, Takanori Yamada<sup>®</sup> Department of Gastroenterology, Iwata City Hospital, Shizuoka, Japan

### Corresponding author

#### Junichi Kaneko, MD

Department of Gastroenterology, Iwata City Hospital, 512-3 Ookubo, Iwata-shi, Shizuoka 438-0002, Japan Fax: +81-538-38-5050 meganerock10@gmail.com

#### References

 [1] Noh SE, Park DH, Kim YR et al. EUS-guided drainage of hepatic abscesses not accessible to percutaneous drainage (with videos). Gastrointest Endosc 2010; 71: 1314–1319

- [2] Kawakami H, Kawakubo K, Kuwatani M et al. Endoscopic ultrasonography-guided liver abscess drainage using a dedicated, wide, fully covered self-expandable metallic stent with flared-ends. Endoscopy 2014; 46 (Suppl. 01): E982–E983
- [3] Zanwar S, Agrawal A. Endosonographyguided caudate lobe liver abscess drainage: a case and review of methods. J Digest Endosc 2022; 13: 119–124
- [4] Rana SS, Ahmed S, Sharma R et al. Safety and efficacy of EUS-guided drainage of liver abscess: a single-center experience. Endosc Ultrasound 2020; 9: 350–351
- [5] Okuno N, Hara K, Mizuno N et al. Risks of transesophageal endoscopic ultrasonography-guided biliary drainage. Gastrointest Interv 2017; 6: 82–84

#### **Bibliography**

Endoscopy 2023; 55: E1099–E1100 DOI 10.1055/a-2174-5604 ISSN 0013-726X © 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



## ENDOSCOPY E-VIDEOS https://eref.thieme.de/e-videos



*E-Videos* is an open access online section of the journal *Endoscopy*, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high-quality video and are published with a Creative Commons CC-BY license. Endoscopy E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: https:// www.research4life.org/access/eligibility/).

This section has its own submission website at

https://mc.manuscriptcentral.com/e-videos