


## The “furrow sign” in confirming proper gastric extent and direction of the myotomy at the end of peroral endoscopic myotomy

OPEN ACCESS




**▶ Video 1** Video clinical cases that demonstrate the practical application and utility of the “furrow sign.”


Peroral endoscopic myotomy (POEM), first performed by Inoue in 2008, stands as a primary treatment for achalasia [1]. Ensuring the appropriate length of the gastric myotomy is crucial, as a myotomy that is too short can lead to an inadequate response. Conversely, excessive length can result in a higher incidence of moderate reflux esophagitis without enhancing clinical effectiveness [2, 3]. While various landmarks and methods guide the proper extent of the tunnel [4, 5], a straightforward approach to confirm the end of the myotomy is lacking.

Herein, we introduce the concept of the “furrow sign,” characterized by a mucosal depression over a muscular defect in the wall post-myotomy. This sign becomes noticeable during gastric retroflexion under full insufflation, serving to confirm the extent and direction of the myotomy before closing the mucosal incision (▶ **Fig. 1**). To evaluate this sign, full carbon dioxide insufflation is applied in the gastric lumen for 60 seconds after myotomy (▶ **Fig. 2**).

Between May and September 2022, we assessed this sign in seven patients who underwent posterior POEM with comprehensive follow-up. Clinical success was



**▶ Fig. 1** Endoscopic images of the “furrow sign,” characterized by a mucosal depression over a muscular defect in the wall post-myotomy after full carbon dioxide insufflation in the stomach.




**▶ Fig. 2** Evaluation of the “furrow sign”. **a** After myotomy. **b** “Furrow sign” is evaluated by applying full carbon dioxide insufflation in the gastric lumen for 60 seconds.

evident in all cases (▶ **Table 1**), and no adverse events linked to full carbon dioxide insufflation during furrow sign evaluation were recorded. On each occasion, the furrow sign was confirmed by at least two endoscopists.

We verified via double-scope transillumination that the depression area seen in the furrow sign aligns with the myotomy area (▶ **Fig. 3**), with its persistence observed even 12 months after POEM (▶ **Video 1**).

In conclusion, the furrow sign corresponds to the myotomy area and can play a crucial role in confirming or fine tuning the accurate length (and also confirming the direction) of the gastric myotomy at the conclusion of the POEM procedure, prior to closing the mucosal incision. To establish its prevalence, sensitivity, specificity, and interobserver agreement, further studies are imperative.

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AN



► **Fig. 3** The depression area seen in the furrow sign aligns with the myotomy area, as confirmed by double-scope transillumination.

► **Table 1** Characteristics and follow-up of patients in whom the “furrow sign” was evaluated during posterior peroral endoscopic myotomy (May–September 2022).

Baseline and follow-up characteristics	N = 7
Baseline	
▪ Age, median (IQR), years	63 (54–72)
▪ Female sex, n (%)	5 (71.4)
▪ Type of achalasia, n (%)	
▪ Type II	5 (71.4)
▪ Type III	2 (28.6)
▪ Pre-POEM IRP, median (IQR)	27.3 (20.2–30.4)
▪ Pre-POEM Eckardt score, median (IQR)	7 (6–7)
Follow-up, median (IQR)	
▪ Gastric myotomy length, cm	2 (2–2.5)
▪ Post-POEM Eckardt score	0 (0–0)
▪ Supine IRP post-POEM, mmHg	13.3 (11.4–15)
▪ Upright IRP post-POEM, mmHg	8.1 (5.5–9.6)
▪ Time until HRM, months	4 (3–5)

IQR, interquartile range; POEM, peroral endoscopic myotomy; IRP, integrated relaxation pressure; HRM, high resolution manometry.

## Conflict of Interest

H. Uchima is a consultant for Lumendi, collaborates with ERBE Spain, Olympus Iberia, Izasa, and has received congresses registrations from Casen-Recordati. R. Muñoz-González, A. Calm, N. Caballero, J. Espinós, V. Moreno de Vega, and I. Marín declare that they have no conflict of interest.

## The authors

**Hugo Uchima**<sup>1,2</sup>, **Raquel Muñoz-González**<sup>1,2</sup>, **Anna Calm**<sup>1</sup>, **Noemí Caballero**<sup>1</sup>, **Jorge Espinos**<sup>2</sup>, **Vicente Moreno de Vega**<sup>1</sup>, **Ingrid Marín**<sup>1</sup>

- 1 Endoscopy Unit, Gastroenterology Department, Hospital Universitari Germans Trias i Pujol, Badalona, Spain
- 2 Endoscopy Unit, Teknon Medical Center, Barcelona, Spain

## Corresponding author

**Hugo Uchima, MD**

Endoscopy Unit, Gastroenterology Department, Hospital Universitari Germans Trias i Pujol, Carretera Canyet s/n, Badalona 08916, Spain  
huchima.germanstrias@gencat.cat

## References

- [1] Werner YB, Hakanson B, Martinek J et al. Endoscopic or surgical myotomy in patients with idiopathic achalasia. *N Engl J Med* 2019; 381: 1209–1210. doi:10.1056/NEJMoa1905380
- [2] Grimes KL, Bechara R, Shimamura Y et al. Gastric myotomy length affects severity but not rate of post-procedure reflux: 3-year follow-up of a prospective randomized controlled trial of double-scope per-oral endoscopic myotomy (POEM) for esophageal achalasia. *Surg Endosc* 2020; 34: 2963–2968. doi:10.1007/s00464-019-07079-0
- [3] Teitelbaum EN, Sternbach JM, El Khoury R et al. The effect of incremental distal gastric myotomy lengths on EGJ distensibility during POEM for achalasia. *Surg Endosc* 2016; 30: 745–750
- [4] Tanaka S, Kawara F, Toyonaga T et al. Two penetrating vessels as a novel indicator of the appropriate distal end of per-oral endoscopic myotomy. *Dig Endosc* 2017; 30: 206–211
- [5] Uchima H, Colán-Hernández J, Aguilar A et al. A simple method to determine the proper length of the gastric myotomy during per-oral endoscopic myotomy for achalasia. *Endoscopy* 2022; 54: E85–E87

## Bibliography

*Endoscopy* 2023; 55: E1209–E1210

DOI 10.1055/a-2199-6956

ISSN 0013-726X

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited.

(<https://creativecommons.org/licenses/by/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

