Endoscopic esophageal cicatricotomy assisted by submucosal injection for the treatment of esophageal stenosis after endoscopic submucosal dissection



Esophageal stenosis is a common complication after endoscopic submucosal dissection (ESD) for early esophageal cancer [1,2]. Post-ESD esophageal stenosis is considered to be related to scar tissue proliferation and fibrosis during the healing process [3]. Symptom relief can be achieved through dilation and radial incision in 83% [4]. However, these invasive strategies do not allow visual control of the operation, and it is difficult to control complications such as bleeding or perforation caused by these treatments.

A 77-year-old patient with esophageal stricture after ESD for early esophageal cancer underwent three endoscopic esophageal stricture expansion procedures over 6 months. Endoscopic examination showed that the patient had irregular stenosis in the middle esophagus, which the gastroscope could not pass. Under white-light and narrowband imaging (**> Fig. 1**), the approximate starting position of the scar could be determined. Submucosal injection and submucosal incision were then performed on the oral side of the scar (> Fig. 2, > Video 1). Under the blue background of the submucosal injectate, which contained indigo carmine, the white scar was clearly distinguishable. By using this contrast in visual appearance, it was easy to accurately identify and release all scar tissue using submucosal incision under direct visualization, until the gastroscope could pass smoothly through the esophageal stricture (**Fig. 3**). The incision length was 27–31 cm from the incisors, and the operation time was about 30 minutes. There were no bleeding or perforation complications during the process.

The esophagus that has undergone multiple expansions will have multiple complex fibrotic areas and severe submucosal adhesions. Submucosal injection-assisted cicatricotomy can aid visualization



Fig.1 Imaging of the stricture. **a** White-light imaging. **b** Narrow-band imaging.



► Fig.2 Submucosal injection and incision were performed on the oral side of the stricture.

of the scar and submucosa during the operation, allowing the endoscopists to accurately control the incision site and depth, minimize the damage to normal mucosa, and successfully reduce the risk of perforation and bleeding.

Endoscopy_UCTN_Code_TTT_1AO_2AH

Conflict of Interest

The authors declare that they have no conflict of interest.



Video 1 Endoscopic cicatricotomy of the esophagus, assisted by submucosal injection.

Funding

Chongqing Science and Technology Plan Project: Specific Project for Chongqing Research Institutions cstc2017jxjl130028



▶ Fig. 3 Treatment of the esophageal stricture. a Submucosal incision and release of all scar tissue under direct visualization. b After treatment, the gastroscope could pass through the esophagus.

The authors

Wen Ji^{1,‡}, Senrong Luo^{1,‡}, Shijun Wang¹, Huan He¹, Weiqing Chen¹

1 Department of Gastroenterology, Chongqing University Cancer Hospital, Chongqing, China

Corresponding author

Weiqing Chen, MD

Department of Gastroenterology, Chongqing University Cancer Hospital, Hanyu Road 181, 400030 Chongqing, China chenwq620712@163.com

References

- Pérez-Cuadrado Robles E, Moreels TG, Piessevaux H et al. Risk factors of refractory post-endoscopic submucosal dissection esophageal strictures. Rev Esp Enferm Dig 2021; 113: 813–819
- [2] Zhang M, Zhu C, Wang Y et al. Differential diagnosis for esophageal protruded lesions using a deep convolution neural network in endoscopic images. Gastrointest Endosc 2021; 93: 1261–1272 doi:10.1016/j. gie.2020.10.00533065026
- [3] Besendörfer M, Müller H, Weiss C et al. Association of clinical factors with postoperative complications of esophageal atresia. Pediatr Neonatol 2021; 62: 55–63
- [4] Hsu WH, Shih HY, Shen CS et al. Prevention and management of esophageal stricture after esophageal ESD: 10 years of experience in a single medical center. J Formos Med Assoc 2023; 122: 486–492 doi:10.1016/j.jfma.2022.12.00636572585

Bibliography

Endoscopy 2023; 55: E1254–E1255 DOI 10.1055/a-2210-0428 ISSN 0013-726X © 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS https://eref.thieme.de/e-videos



E-Videos is an open access online section of the journal *Endoscopy*, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high-quality video and are published with a Creative Commons CC-BY license. Endoscopy E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: https:// www.research4life.org/access/eligibility/).

This section has its own submission website at

https://mc.manuscriptcentral.com/e-videos

[‡] Contributed equally to this article