



# Endoscopic submucosal dissection for superficial esophageal cancer with ulcer scarring using a combination of pocket creation, gel immersion, and red dichromatic imaging




**▶ Video 1** Successful pharyngeal endoscopic submucosal dissection using a novel clip-traction band device.

In recent years, the usefulness of endoscopic resection (ER) methods, such as underwater endoscopic submucosal dissection, has been widely reported [1]. Moreover, recently, from the viewpoint of securing the visual field during endoscopy, gel immersion endoscopy has been reported to be useful [2,3]. Gel immersion endoscopy and underwater ER are used in challenging endoscopic procedures owing to their buoyancy effects. The pocket creation method is widely used to overcome endoscopic submucosal dissection (ESD) difficulties [4]. Red dichromatic imaging, which improves the visibility of deep blood vessels and bleeding points using longer wavelengths of light, is recently being extensively used, and its effectiveness has been reported [5]. We used red dichromatic imaging to improve visibility of the submucosal and muscular layers during ESD.

Here, we describe a case of successful ESD of a superficial esophageal carcinoma on a scar after ER using the pocket creation method, underwater conditions, gel immersion endoscopy, and other techniques.




**▶ Fig. 1** The target lesion. The lesion was recognized as a brownish area, 30 mm in size, on the posterior wall, 35 cm from the incisor row. The area covered by the yellow enclosure indicates the scar from previous treatment, and the red arrow indicates the target lesion. **a** White light imaging. **b** After spraying with Lugol's solution, the lesion was observable in the Lugol's voiding area.




**▶ Fig. 2** Schema image of the treatment process. **a** Fibrosis due to previous treatment was considered to have formed in the shallow submucosal layer, and the visibility of the fibrosis was ensured through filling the pocket with gel. **b** Using the pocket creation method, submucosal dissection was performed to a greater extent beyond the lesion border to break through the fibrosis.

A 68-year-old man, who had previously undergone curative resection of esophageal cancer in the middle thoracic region using ESD 30 years prior, was referred for follow-up. An upper gastrointestinal endoscopy revealed a new superficial carcinoma of the esophagus (30 mm in size) located on the post-treatment scar (▶ Fig. 1). ESD was planned. During this treatment, in addition to the pocket

creation method to break through the fibrosis, underwater endoscopic submucosal dissection and gel immersion endoscopy devices were used inside the pocket to add to the buoyancy effect (▶ Fig. 2, ▶ Fig. 3). Red dichromatic imaging was also used to visualize the fibrosed area and to ensure a good visual field (▶ Fig. 4). These measures made it possible to complete submucosal dissec-



► **Fig. 3** Pocket creation method. **a** Mucosal incision was initiated from the oral side of the lesion, and a pocket was created. **b** Inside the pocket, the gel provided a good visual field.



► **Fig. 4** Red dichromatic imaging. **a** White light observation may cause gels and water to become cloudy. **b** Red dichromatic imaging, using a long wavelength band that is not easily scattered or absorbed proximally, improves target visibility.

## Corresponding author

**Tomoaki Tashima, MD, PhD**

Department of Gastroenterology, Saitama Medical University International Medical Center, 1397-1 Yamane, Hidaka City, Saitama 350-1298, Japan  
tomoaki\_tashima@yahoo.co.jp

## References

- [1] Nagata M. Usefulness of underwater endoscopic submucosal dissection in saline solution with a monopolar knife for colorectal tumors (with videos). *Gastrointest Endosc* 2018; 87: 1345–1353. doi:10.1016/j.gie.2017.11.032
- [2] Miura Y, Yano T, Takezawa T et al. Gel immersion endoscopy simplifies hemostasis during endoscopic submucosal dissection using the pocket-creation method. *Endoscopy* 2018; 50: E294–E295
- [3] Nakano Y, Tashima T, Jinushi R et al. Conversion from conventional esophageal endoscopic submucosal dissection to the gel immersion method. *Endosc Int Open* 2021; 9: E1756–E1757
- [4] Yoshida N, Naito Y, Yasuda R et al. The efficacy of the pocket-creation method for cases with severe fibrosis in colorectal endoscopic submucosal dissection. *Endosc Int Open* 2018; 6: E975–E983
- [5] Yahagi N, Fujimoto A, Horii J et al. Dual red imaging: a novel endoscopic imaging technology visualizing thick blood vessels in the gastrointestinal wall. *Endosc Int Open* 2019; 7: E1632–E1625. doi:10.1055/a-0749-0075

tion with clear visibility of the fibrosed area (► **Video 1**).

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AC

## Funding Information

JSPS KAKENHI  
23K15052

Tomoaki Tashima was supported by JSPS KAKENHI.

## Acknowledgement

We would like to thank Editage ([www.editage.jp](http://www.editage.jp)) for English language editing.

## Conflict of Interest

The authors declare that they have no conflict of interest.

## The authors

**Tsubasa Ishikawa<sup>1</sup>**, **Tomoaki Tashima<sup>1</sup>**, **Takahiro Muramatsu<sup>1</sup>**, **Yumi Mashimo<sup>1</sup>**, **Shomei Ryozaawa<sup>1</sup>**

<sup>1</sup> Department of Gastroenterology, Saitama Medical University International Medical Center, Hidaka, Japan

## Bibliography

*Endoscopy* 2024; 56: E87–E88  
DOI 10.1055/a-2234-8435  
ISSN 0013-726X  
© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited.  
(<https://creativecommons.org/licenses/by/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14,  
70469 Stuttgart, Germany

