Novel method for retrieving a migrated plastic stent using an 11.5-Fr pusher sheath: The stent encapsulation method



Plastic stent placement using endoscopic retrograde cholangiopancreatography (ERCP) is a widely used biliary drainage technique [1]. Migration of the plastic stent is a major adverse effect. Retrieval grasping techniques using baskets, snares, forceps, and balloon catheters have been reported [2]; however, these techniques are not always successful because of biliary stenosis and tight anchoring of plastic stent flaps [3]. Herein, we introduce a novel method for retrieving a migrated plastic stent using an 11.5-Fr pusher sheath.

A 58-year-old woman with a history of multiple endoscopic treatments, including placement of a metallic stent and a plastic stent inside the metallic stent, for distal biliary obstruction caused by malignant lymphoma, was admitted to our institution because of cholangitis (**Fig.1**). Computed tomography revealed a dilated intrahepatic bile duct, which was suspected to be related to stent occlusion. We performed endoscopic retrograde cholangiopancreatography (ERCP) and attempted to retrieve the plastic stent using forceps, a snare, and a balloon catheter; however, these attempts failed because of the tight anchoring of a flap of the plastic stent to the metallic stent and the migration of the plastic stent into the bile duct.

Therefore, we decided to try a stent encapsulation method (> Video 1). We passed a 0.025-inch guidewire through the plastic stent and advanced an 11.5-Fr pusher sheath (Oasis; Cook Medical, Bloomington, Indiana, USA) over the quidewire. Forceps (E634045; Olympus Medical Systems, Tokyo, Japan) were then inserted through the pusher sheath (> Fig. 2). The end of the plastic stent was then grasped with the forceps (**Fig.3**). The 11.5-Fr pusher sheath device was advanced into the bile duct over the forceps encasing the entire length of the plastic stent (> Fig. 4), including the flap. The plastic stent was



▶ Fig. 1 X-ray of 58-year-old woman showing possibly occluded plastic stent coaxial within a metallic stent, placed for distal biliary obstruction caused by malignant lymphoma.



• **Fig. 2** Forceps are passed through an 11.5-Fr pusher sheath on the guidewire.

then removed successfully through the pusher sheath. Finally, another metallic stent was deployed inside the previously placed metallic stent (> Fig. 5).



► Fig. 3 The forceps are used to grasp the end of the migrated plastic stent.



Video 1 Successful removal of a migrated and tightly stuck biliary plastic stent using a novel stent encapsulation method.

This novel method can be useful for removing a migrated plastic stent after advancement of a sheath device beyond a stricture.

Fig.4 The 11.5-Fr pusher sheath is advanced over the forceps to encase the entire length of the migrated stent.

Endoscopy_UCTN_Code_TTT_1AR_2AZ

Conflict of Interest

The authors declare that they have no conflict of interest.

The authors

Fumitaka Niiya^{1©}, Masataka Yamawaki¹, Jun Noda¹, Tetsushi Azami¹, Yuichi Takano^{1©}, Fumiya Nishimoto¹, Masatsugu Nagahama¹

1 Internal Medicine, Division of Gastroenterology, Showa University Fujigaoka Hospital, Yokohama, Japan

Corresponding author

Fumitaka Niiya, MD, PhD

Division of Gastroenterology, Department of Internal Medicine, Showa University Fujigaoka Hospital, 1-30 Fujigaoka, Aoba-ku, Yokohama, Kanagawa, Japan f.niiya@med.showa-u.ac.jp

References

- Dumonceau JM, Tringali A, Blero D et al. Biliary stenting: indications, choice of stents and results: European Society of Gastrointestinal Endoscopy (ESGE) Clinical Guideline. Endoscopy 2012; 44: 277–292. doi:10.1055/a-0659-9864
- [2] Tarnasky PR, Cotton PB, Baillie J et al. Proximal migration of stents: attempted endoscopic retrieval in forty-one patients. Gastrointest Endosc 1995; 42: 513–520. doi:10.1016/s0016-5107(95)70003-x
- [3] Calcara C, Broglia L, Comi G et al. Plastic biliary stent migration during multiple stents placement and successful endoscopic removal using intra-stent balloon inflation technique: a case report and literature review. Am J Case Rep 2016; 17: 65–69

Bibliography

Endoscopy 2024; 56: E134–E135 DOI 10.1055/a-2239-3065 ISSN 0013-726X © 2024. The Author(s).

► Fig.5 Another metallic stent is deployed inside the previously placed metallic stent.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS https://eref.thieme.de/e-videos

E-Videos is an open access online section of the journal *Endoscopy*, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high-quality video and are published with a Creative Commons CC-BY license. Endoscopy E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: https:// www.research4Life.org/access/eligibility/).

This section has its own submission website at https://mc.manuscriptcentral.com/e-videos