Endoscopic sutured purse-string resection: a novel technique for resection of large gastric subepithelial lesions





Video 1 Endoscopic sutured pursestring resection for the removal of subepithelial lesions in the gastrointestinal tract

Subepithelial lesions (SELs) in the gastrointestinal (GI) tract are common and often necessitate removal, particularly when > 20 mm [1, 2]. The choice of endoscopic resection depends on various factors, including lesion characteristics, location, and evidence of deeper tissue involvement [2,3]. Challenges in achieving full-thickness resection have driven the development of innovative over-thescope devices [4]. However, these devices are typically restricted to lesions < 30 mm, and their size and rigidity often hinder passage beyond the pharynx. We present here a novel technique for accomplishing full-thickness resection of SELs, known as endoscopic sutured purse-string resection (ESPR). ESPR employs the Overstitch device (Apollo Endosurgery, Austin, Texas, USA), a wellestablished tool for placing full-thickness endoscopic sutures. It involves creating a purse-string configuration around the lesion (> Fig. 1) before resection, enabling the safe and complete removal of even larger lesions.

Patient selection followed European Society of Gastrointestinal Endoscopy guidelines, with a preference for lesions located on the greater curve of the stom-



▶ Fig. 1 Schematic showing lesion marking and suture placement for endoscopic sutured purse-string resection. a Lesion (L) of diameter (d), which allows placement of mucosal diathermy marks (m) and purse-string sutures (ps). b Lesion after tightening of the purse string, with corresponding sites of mucosal marks and suture sites. Snare resection is conducted at or underneath the diathermy marks. SM, submucosal layer; MP, muscularis propria.

ach, primarily due to the easier access provided by the Overstitch device. Lesions underwent thorough characterization using endoscopic direct visualization, endoscopic ultrasound, and crosssectional imaging (computed tomography), following a standardized protocol.

ESPR is conducted as follows (**> Video 1**): marking the boundaries of the lesion, applying an endoscopic purse string, tenting the lesion with forceps, and creating a pseudopolyp by tightening the purse string. Resection was carried out using a large snare, and the site underwent meticulous inspection for completeness before being oversewn with a Z-shaped suture (**Fig.2**). In the two cases where this technique was employed, no perioperative complications were encountered. Patients were discharged on the same day, and histological examination confirmed complete resection, including the muscularis propria and serosa layers, along with omental fat in one case.

This technique presents a safe and viable solution for resecting gastric SELs, effectively addressing the limitations of existing techniques. Its potential applicability extends beyond the stomach, offering a promising avenue for further exploration and adoption in the field of GI endoscopy.



▶ Fig. 2 Image showing the two lesions (a and b) resected using the endoscopic sutured purse-string resection technique. a1 30-mm gastric antral subepithelial lesion. b1 15-mm greater curvature lesion. a2, b2 Resected site showing complete resection. a3, b3 Closure of the resected site with Z-shaped suture (4-suture point).

Endoscopy_UCTN_Code_CCL_1AB_2AD_3AB

Acknowledgement

BH would like to thank Jacob and Isaac Hayee for their assistance in preparing schematic figure 1.

Conflict of Interest

The authors declare that they have no conflict of interest.

The authors

Olaolu Olabintan¹, Theodoros Voulgaris¹, Homira Ayubi¹, Sri Thrumurthy^{1,2}, Amyn Haji^{1,3}, Bu'Hussain Hayee^{1,2,3}

- 1 Gastroenterology & Advance Therapeutic Endoscopy Department, King's College Hospital NHS Foundation Trust, London, United Kingdom
- 2 Upper GI Surgery & Advance Therapeutic Endoscopy Department, King's College Hospital NHS Foundation Trust, London, United Kingdom
- 3 Colorectal Surgery & Advance Therapeutic Endoscopy Department, King's College Hospital NHS Foundation Trust, London, United Kingdom

Corresponding author

Olaolu Olabintan, MD

Gastroenterology & Advance Therapeutic Endoscopy Department, King's College Hospital NHS Foundation Trust, Denmark Hill, London SE5 9RS, United Kingdom olaolu.olabintan@nhs.net

References

- [1] Sharzehi K, Sethi A, Savides T. AGA clinical practice update on management of subepithelial lesions encountered during routine endoscopy: expert review. Clin Gastroenterol Hepatol 2022; 20: 2435–2443. doi:10.1016/j.cgh.2022.05.054
- [2] Deprez P, Moons L, O'Toole D et al. Endoscopic management of subepithelial lesions including neuroendocrine neoplasms: European Society of Gastrointestinal Endoscopy (ESGE) Guideline. Endoscopy 2022; 54: 412– 429
- [3] Moons L, Bastiaansen B, Richir M et al. Endoscopic intermuscular dissection for deep submucosal invasive cancer in the rectum: a new endoscopic approach. Endoscopy 2022; 54: 993–998. doi:10.1055/a-1748-8573
- [4] Hajifathalian K, Ichkhanian Y, Dawod Q et al. Full-thickness resection device (FTRD) for treatment of upper gastrointestinal tract lesions: the first international experience. Endosc Int Open 2020; 8: 1291–1301. doi:10.1055/a-1216-1439

Bibliography

Endoscopy 2024; 56: E352–E353 DOI 10.1055/a-2299-1899 ISSN 0013-726X © 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS https://eref.thieme.de/e-videos



E-Videos is an open access online section of the journal *Endoscopy*, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high-quality video and are published with a Creative Commons CC-BY license. Endoscopy E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: https:// www.research4Life.org/access/eligibility/).

This section has its own submission website at https://mc.manuscriptcentral.com/e-videos