Peroral cholangioscopy-guided lithotripsy using a novel thin cholangioscope under balloon enteroscopy for Roux-en-Y anastomosis





► Fig.1 Computed tomography revealed large stones (red arrows) in the intrahepatic bile duct.

Stone extraction using endoscopic retrograde cholangiopancreatography (ERCP) is less invasive than surgical procedures. However, stone extraction in patients with surgically altered anatomy, such as those who have undergone a Roux-en-Y procedure, is challenging. Although balloon enteroscopy is useful for such cases, there is still room for improvement [1,2]. Peroral cholangioscopy (POCS)-guided lithotripsy can aid in the extraction of stones that are difficult to remove [3,4]. However, performing POCS-guided lithotripsy under balloon enteroscopy is difficult because cholangioscopes have an approximate diameter of 10Fr and cannot pass through the forceps channel of the balloon enteroscope. This report describes a patient with a Roux-en-Y anastomosis who was successfully treated with POCS-guided lithotripsy using a novel thin cholangioscope under balloon enteroscopy.

A 51-year-old woman who had previously undergone a diversion operation and hepaticojejunostomy with Roux-en-Y for congenital biliary dilatation 7 years earlier was referred to our center. Computed tomography revealed large stones in the intrahepatic bile duct (► Fig. 1). Consequently, we performed ERCP using a short-type single-balloon enteroscope (SIF-H290; Olympus, Tokyo, Japan) with



Fig. 2 Thin cholangioscope (eyeMAX; Micro-Tech, Nanjing, China) with a length of 219 cm and diameter of 9 Fr.



► Fig. 3 Cholangiography revealed large stones (red arrows) in the intrahepatic bile duct.

a working length of 152 cm and a working channel with a diameter of 3.2 mm [1,2]. Additionally, we performed POCS-guided lithotripsy using a thin cholangioscope (eyeMAX; Micro-Tech, Nanjing, China) with a length of 219 cm and diameter of 9 Fr [5] (**> Fig. 2, > Video 1**) as complete stone extraction was difficult without POCS. Cholangiography revealed large stones in the intrahepatic bile duct (**> Fig. 3**). Subsequently, POCS was per-



Video 1 Successful peroral cholangioscopy-guided lithotripsy using a novel thin cholangioscope under balloon enteroscopy in a patient who had previously undergone a Roux-en-Y procedure.

formed using a thin cholangioscope, revealing multiple large stones in the intrahepatic bile duct (**> Fig. 4 a**). POCS-guided lithotripsy was performed while main-



Fig.4 Cholangioscopy findings. **a** Cholangioscopy revealed multiple large stones in the intrahepatic bile duct. **b**, **c** Peroral cholangioscopy-guided lithotripsy is performed while maintaining a clear field of view. **d** Stone fragmentation has been successfully achieved.

taining a clear field of view (▶ Fig.4b, c). Successful stone fragmentation was achieved (▶ Fig.4d), followed by complete stone extraction (▶ Fig.5). The thin cholangioscope was effective for POCS-guided lithotripsy even though a balloon enteroscope was used. This novel thin cholangioscope can improve the success rate of stone extraction in patients with a surgically altered anatomy.

Endoscopy_UCTN_Code_TTT_1AR_2AH

Acknowledgement

We thank Editage (www.editage.com) for English language editing.

Conflict of Interest

The authors declare that they have no conflict of interest.

The authors

Yuki Tanisaka^{, 1}^o, Masafumi Mizuide¹, Akashi Fujita¹^o, Takahiro Shin¹, Kei Sugimoto¹, Ryuhei Jinushi¹, Shomei Ryozawa¹

1 Gastroenterology, Saitama Medical University International Medical Center, Hidaka, Japan

Corresponding author

Yuki Tanisaka, MD, PhD

Department of Gastroenterology, Saitama Medical University International Medical Center, 1397-1 Yamane, Hidaka, Saitama 350-1298, Japan tanisaka1205@gmail.com

References

[1] Tanisaka Y, Ryozawa S, Itoi T et al. Efficacy and factors affecting procedure results of short-type single-balloon enteroscopyassisted ERCP for altered anatomy: a multicenter cohort in Japan. Gastrointest Endosc 2022; 95: 310–318.e1



► **Fig.5** Endoscopy revealed successful stone extraction.

- [2] Tanisaka Y, Mizuide M, Fujita A et al. Factors affecting complete stone extraction in the initial procedure in short type single-balloon enteroscopy-assisted endoscopic retrograde cholangiography for patients with Roux-en-Y gastrectomy. Scand J Gastroenterol 2022; 57: 1390–1396
- [3] Maydeo AP, Rerknimitr R, Lau JY et al. Cholangioscopy-guided lithotripsy for difficult bile duct stone clearance in a single session of ERCP: results from a large multinational registry demonstrate high success rates. Endoscopy 2019; 51: 922–929
- [4] Angsuwatcharakon P, Kulpatcharapong S, Ridtitid W et al. Digital cholangioscopyguided laser versus mechanical lithotripsy for large bile duct stone removal after failed papillary large-balloon dilation: a randomized study. Endoscopy 2019; 51: 1066– 1073. doi:10.1055/a-0848-8373
- [5] Liu WH, Huang XY, Hu X et al. Initial experience of visualized biliary cannulation during ERCP. Endoscopy 2023; 55: 1037–1042

Bibliography

Endoscopy 2024; 56: E360–E361 DOI 10.1055/a-2299-2477 ISSN 0013-726X © 2024. The Author(s). This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

