Innovative push percutaneous endoscopic gastrostomy by means of gastropexy and Foley catheter in a patient with advanced head and neck and esophageal cancer





**Fig.1** Computed tomography (CT) of the neck in a 54-year-old woman demonstrated an irregular circumferential enhancing mass involving the cervical esophagus, left epiglottis, left aryepiglottic fold, and the left pyriform sinus, with multiple cervical and paraesophageal lymph nodes (arrow): **a** coronal view; **b** sagittal view; **c**, **d** horizontal views.





Push percutaneous endoscopic gastrostomy (PEG) is a noninvasive endoscopically assisted procedure that avoids the disadvantages of surgical gastrostomy while minimizing the risks of cancer seeding to the stroma, especially in patients with head and neck or esophageal cancers [1–3]. However, it still has some drawbacks in terms of tube-related complications and the cost of the PEG kit [2, 4]. There are many different techniques for gastropexy with dedicated kits available [3]. In addition, the Foley catheter may be used safely as a gastrostomy tube and is available at most institutions and inexpensive compared with commercial devices [5]. With these considerations we developed a safe and simple push PEG technique for oncology patients. A 54-year-old woman who had been diagnosed with malignant supraglottic and cervical esophageal cancer needed a push-PEG procedure for nutritional support (**> Fig. 1, > Video 1**). A pediatric gastroscope was used for endoscopic

visualization, and the PEG site was selec-

ted using gastric transillumination and finger palpation.

- The gastropexy steps were as follows:
- Extracorporeally, a 2.0 nylon thread was passed through an 18-G spinal needle and the distal end of the thread was turned and passed back along the outside of the needle. Needle and thread were inserted into the stomach and manipulated to form a loop.
- Another 18-G spinal needle was inserted into the stomach about 2 cm



Video 1 Push percutaneous endoscopic gastrostomy using gastropexy and a Foley catheter in a patient with advanced head and neck and esophageal cancer.

from the first. The tip was manipulated to pass through the loop created by the first needle and thread. The end of a nylon thread was then passed through and out of the second spinal needle and therefore through the loop.

- This second needle was then retracted, leaving the second thread passing through the loop (> Fig. 2 a).
- The loop was then retracted, pulling with it the second thread.
- The ends of the second nylon thread were then knotted extracorporeally to suture together the gastric and abdominal walls.
- To complete the gastropexy, the same process was repeated at a distance of 3–4 cm from the previous location.

An 18-G over-the-needle catheter was inserted into the stomach in the area between the two gastropexy sutures, and a guidewire was threaded through this puncture site. An Alken metal dilator was used to gradually increase the size of the track from 6 Fr to 24 Fr (▶ Fig. 2b). A Foley catheter was prepared by cutting the tip to enable passage over the guidewire, and was used to replace the dilator (▶ Fig. 2c). To properly test the feeding tube, the balloon of the Foley catheter was inflated (> Fig. 2 d). Then, the position of the catheter was checked to ensure it was correctly placed. The feeding functionality was tested using water. Finally, the Foley catheter was sutured and tied to prevent the skin at the ostomy site from moving.

The patient was stable and was discharged from the hospital just 2 days after the procedure. At a follow-up visit 4 weeks later, the patient was being treated with chemoradiotherapy and having a full calorific intake from enteral nutrition, without any complications.

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AK

## **Conflict of Interest**

The authors declare that they have no conflict of interest.

# The authors

#### Prasit Mahawongkajit<sup>1©</sup>, Kanokwan Bunprachoen<sup>1</sup>

1 Surgery, Thammasat University Faculty of Medicine, Pathum Thani, Thailand

## Corresponding author

## Prasit Mahawongkajit, MD, PhD

Department of Surgery, Faculty of Medicine, Thammasat University (Rangsit Campus), 95/209 Moo 18, Paholyothin Road, Amphur Klongluang, Pathum Thani, Thailand, 12120 prasit\_md@yahoo.com

#### References

- Siu J, Fuller K, Nadler A et al. Metastasis to gastrostomy sites from upper aerodigestive tract malignancies: a systematic review and meta-analysis. Gastrointest Endosc 2020; 91: 1005–1014
- [2] Mahawongkajit P, Techagumpuch A, Limpavitayaporn P et al. Comparison of introducer percutaneous endoscopic gastrostomy with open gastrostomy in advanced esophageal cancer patients. Dysphagia 2020; 35: 117– 120
- [3] Nunes G, Fonseca J, Barata AT et al. Nutritional support of cancer patients without oral feeding: how to select the most effective technique? GE Port J Gastroenterol 2020; 27: 172–184

- [4] Kucha P, Zorniak M, Szmit M et al. To push or to pull? A clinical audit on the efficacy and safety of the pull and push percutaneous endoscopic gastrostomy techniques in oncological patients United European Gastroenterol J 2023; 11: 951–959. doi:10.1002/ ueq2.12467
- [5] Kadakia SC, Cassaday M, Shaffer RT. Comparison of Foley catheter as a replacement gastrostomy tube with commercial replacement gastrostomy tube: a prospective randomized trial. Gastrointest Endosc 1994; 40: 188–193. doi:10.1016/s0016-5107(94) 70165-2

## Bibliography

Endoscopy 2024; 56: E428–E429 DOI 10.1055/a-2316-1067 ISSN 0013-726X © 2024. The Author(s). This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



## ENDOSCOPY E-VIDEOS https://eref.thieme.de/e-videos



*E-Videos* is an open access online section of the journal *Endoscopy*, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high-quality video and are published with a Creative Commons CC-BY license. Endoscopy E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: https:// www.research4life.org/access/eligibility/).

This section has its own submission website at

https://mc.manuscriptcentral.com/e-videos