

Significance of the sFlt-1/PlGF Ratio in Certain Cohorts – What Needs to be Considered?

Wertigkeit des sFlt-1/PlGF-Quotienten in bestimmten Kollektiven – was gibt es zu beachten?



Authors

Oliver Graupner¹, Stefan Verloren², Tanja Groten³, Dietmar Schlembach⁴, Holger Stepan⁵, Bettina Kuschel¹, Anne Karge¹, Ulrich Pecks⁶

Affiliations

- 1 Klinik und Poliklinik für Frauenheilkunde, Universitätsklinikum rechts der Isar, Technische Universität München, München, Germany
- 2 Klinik für Geburtsgynäkologie, Charité – Universitätsmedizin Berlin, Berlin, Germany
- 3 Klinik für Geburtsgynäkologie, Universitätsklinikum Jena, Jena, Germany
- 4 Klinik für Geburtsgynäkologie, Klinikum Neukölln, Vivantes Netzwerk für Gesundheit GmbH, Berlin, Germany
- 5 Klinik für Geburtsgynäkologie, Universitätsklinikum Leipzig, Leipzig, Germany
- 6 Klinik für Gynäkologie und Geburtshilfe, Universitätsklinikum Würzburg, Würzburg, Germany

Keywords

sFlt-1/PlGF ratio, angiogenic factors, preeclampsia

Schlüsselwörter

sFlt-1/PlGF-Quotient, angiogene Marker, Präeklampsie

received 7.3.2024

accepted after revision 5.5.2024

Bibliography

Geburtsh Frauenheilk 2024; 84: 629–634

DOI 10.1055/a-2320-5843

ISSN 0016-5751

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

Correspondence

PD Dr. Oliver Graupner
Klinik und Poliklinik für Frauenheilkunde
Universitätsklinikum rechts der Isar,
Technische Universität München
Ismaninger Str. 22
81675 München, Germany
oliver.graupner@googlemail.com
oliver.graupner@mri.tum.de

Deutsche Version unter:
<https://doi.org/10.1055/a-2320-5843>.

ABSTRACT

The sFlt-1/PlGF ratio is an established tool in clinical practice, where it is part of a diagnostic algorithm and informs the prognosis of preeclampsia (PE). Maternal and gestational comorbidities can affect the performance of the sFlt-1/PlGF ratio and its constituent elements, and a good understanding of the potential pitfalls is required. The objective of this paper was to provide a current narrative review of the literature on the diagnostic and predictive performance of the sFlt-1/PlGF ratio in specific patient cohorts. Potential factors which can negatively affect the clinical interpretability and applicability of the sFlt-1/PlGF ratio include chronic kidney disease, twin pregnancy, and maternal obesity. Pathophysiological mechanisms related to these factors and disorders can result in different concentrations of sFlt-1 and/or PlGF in maternal blood, meaning that the use of standard cut-off values in specific cohorts can lead to errors. To what extent the cut-off values should be adapted in certain patient cohorts can only be clarified in large prospective cohort studies. This applies to the use of the ratio both for diagnosis and prognosis.

ZUSAMMENFASSUNG

Der sFlt-1/PlGF-Quotient hat sich bezogen auf den diagnostischen Algorithmus und die Prognoseabschätzung der Präeklampsie (PE) im klinischen Alltag etabliert. Mütterliche und gestationsbedingte Komorbiditäten können die Performance des sFlt-1/PlGF-Quotienten und seiner Bestandteile beeinflus-

sen und erfordern ein entsprechendes Wissen über mögliche Fallstricke. Ziel dieser Arbeit ist es, eine aktuelle, narrative Literaturübersicht im Kontext der diagnostischen und prädiktiven Performance des sFlt-1/PIGF-Quotienten in speziellen Patientinnenkollektiven zu geben. Zu potenziellen Störfaktoren der klinischen Interpretier- und Anwendbarkeit des sFlt-1/PIGF-Quotienten zählen hier u.a. eine chronische Nierenerkrankung, Zwillingsschwangerschaften und eine mütterliche Adi-

positas. Dabei kommt es auf Basis von pathophysiologischen Mechanismen zu unterschiedlichen Konzentrationen von sFlt-1 und/oder PIGF im Blut der Mutter, wodurch eine Übertragung der gängigen Cut-off-Werte fehleranfällig sein kann. Inwieweit bei diesen Patientinnengruppen eine Anpassung der Cut-off-Werte erfolgen sollte, kann nur in großen prospektiven Kohortenstudien geklärt werden. Dies gilt sowohl für die Frage der Diagnose als auch Prognose.

Introduction

The sFlt-1 (soluble fms-like tyrosine kinase 1)/PIGF (placental growth factor) ratio has been found to be useful in routine clinical practice as it provides information that feeds into the diagnostic algorithm for hypertensive disorders of pregnancy (HDPs) [1].

The pathophysiological consequences of an angiogenic imbalance (in favor of sFlt-1 and at the expense of PIGF – expressed by an increased sFlt-1/PIGF ratio) is endothelial dysfunction. The resulting vascular leakage and consequent intravascular lack of volume and the reduced perfusion of organs such as the kidneys, liver, or placenta leads to the well-known symptoms of preeclampsia (PE) [2]. It is important to emphasize at this point that in patients with endothelial disease, sFlt-1 and/or PIGF may be increased or decreased even outside of pregnancy [3, 4, 5, 6, 7]. The cut-off values for the sFlt-1/PIGF ratio to estimate the risk of PE and the degree of placental dysfunction but also to predict an unfavorable neonatal or maternal outcome and the associated short time to delivery have been published with a high level of evidence [8, 9, 10, 11, 12] (► **Table 1**).

Overview

The use of the sFlt-1/PIGF ratio and its constituent elements in special patient cohorts is described in more detail below. The focus is on diagnosing PE and on the predictive value of angiogenic factors with regards to adverse perinatal outcomes (APOs).

Chronic Kidney Disease

The prevalence of PE in women with chronic kidney disease (CKD) can be up to 40% [18, 19], in women with advanced CKD it may be as high as 70% [19, 20, 21]. In a CKD cohort, the overlapping clinical presentations of the two pathologies, both of which usually include pre-existing hypertension and proteinuria, can make it more difficult to identify the underlying cause [22]. Moreover, renal dysfunction disorders such as systemic lupus erythematosus (SLE), thrombocytopenic purpura (TTP) or atypical hemolytic uremic syndrome (aHUS) may first appear during pregnancy or be exacerbated by pregnancy, which makes differentiating placental complications as difficult as predicting the pregnancy outcome [23, 24]. Rolfo et al. were able to show that, despite overlapping characteristics (hypertension and proteinuria), it is possible to differentiate between PE and CKD using the sFlt-1/PIGF ratio. In their prospective cohort study, patients with CKD (n = 23) had signifi-

cantly lower sFlt-1/PIGF values (4.00 [interquartile range [IQR]: 0.51–136.59] vs. 435.79 [IQR: 160.90–1153.53]; $p < 0.001$) compared to patients without CKD (n = 34) but with PE (based on the diagnostic criteria hypertension and proteinuria before 20 + 0 weeks of gestation) [13]. Moreover, a higher APO rate and a shorter mean time to delivery in the CKD cohort (n = 171) was observed, which correlated with the level of anti-angiogenic imbalance indicated by the sFlt-1/PIGF ratio [25]. The usefulness of the sFlt-1/PIGF ratio to diagnose PE in a CKD cohort was emphasized in the S2 k-guideline “Chronic Kidney Disease and Pregnancy” [26]: “PIGF and sFlt-1 should be used as additional diagnostic parameters if preeclampsia is suspected in patients with CKD.” The special pathophysiological features of the circulating PIGF levels in CKD patients need to be considered as the figures can be misleading. Outside pregnancy, increased PIGF production and decreased sFlt-1 concentrations have been demonstrated in CKD cohorts [27, 28]. PIGF is filtered in the kidney and excreted in urine [29]. Progressive impairment of kidney function in pregnancy can therefore result in higher PIGF levels due to reduced renal clearance. In a “normal” population, a PIGF of less than 100 pg/ml is suspicious or associated with a higher risk of placental complications. To achieve the same sensitivity and specificity, the recommendation is that the threshold value should be increased for women with CKD [14]. Wiles et al. therefore recommended that the monitoring of CKD patients with a PIGF serum level of less than 150 pg/ml after 20 + 0 weeks of gestation should be increased, although it must be noted that the value of PIGF serum levels for predicting PE decreases as the impairment of kidney function increases [14].

Obesity

As with pre-existing diabetes mellitus, obesity is a risk factor for developing PE and, like PE, it is characterized by a pro-inflammatory micromilieu [30, 31]. The PROGNOSIS study, which evaluated and established gestational age-specific cut-off values for the sFlt-1/PIGF ratio, did not differentiate between obese (BMI $> 30 \text{ kg/m}^2$) and normal-weight (BMI $< 25 \text{ kg/m}^2$) women based on BMI [32] and it therefore also did not investigate whether different threshold values would be necessary for obese pregnant women [8]. Zera et al. observed an inverse correlation between sFlt-1 and maternal BMI, i.e., the higher the BMI, the lower the sFlt-1 level [15]. A recently published prospective observational study (n = 1450 PE vs. n = 1065 controls) showed that women with PE and obesity had significantly lower sFlt-1 concentrations in the

► **Table 1** Overview of established cut-off values [8, 10] for the sFlt-1/PlGF ratio and its constituent elements for the reference cohort, and the special pathophysiological features associated with specific maternal or gestational comorbidities.

	Reference cohort [8, 10]	Chronic kidney disease [13, 14]	Obesity [15, 16]	Twin pregnancy [17]
sFlt-1	Stable in 1 st and 2 nd trimester, increased in 3 rd trimester	Changes in sFlt-1 values the same as for the reference cohort	Significantly lower sFlt-1 values when BMI is > 30	Increase in sFlt-1 in the 3 rd trimester is more pronounced compared to singleton pregnancies
PlGF	Continuous increase, followed by a decrease from week 34 of gestation	In principle, higher PlGF values	Similar PlGF course	Similar PlGF course
sFlt-1/PlGF	Values decrease until the 3 rd trimester when the ratio increases again	Lower GFR correlates with lower informative value of the ratio	Ratio significantly lower in the 3 rd trimester if BMI is > 30	Ratio higher from 29 + 0 weeks of gestation
Possible cause of change	–	PlGF is eliminated renally and PlGF accumulates if GFR is reduced	Higher plasma volume TNF- α inhibits endogenous sFlt-1 production	Higher placental mass
sFlt-1/PlGF				
< 38	PE unlikely in the next 4 weeks	A ratio of less than 38 does not rule out PE	Data unclear	Established cut-off values after 29 + 0 weeks of gestation cannot be transferred
38–85 or 110	Development of PE over the course of pregnancy is possible, control within 7 days is recommended	–		
> 85 or 110	PE very probable	PE very probable		
PlGF < 100	Placental dysfunction is probable	A value of more than 100 does not rule out placental dysfunction		
MTTD	Ratio > 655: delivery within 48 h or 7 days probable	PlGF > 150 does not rule out high probability of delivery within the next 14 days	Data unclear	Ratio < 38: delivery within next 2 weeks unlikely
APO/AMO	Ratio < 38: unlikely Ratio > 655: probable	Data unclear	Ratio < 38 does not rule out an APO	Ratio < 38 does not rule out an APO

AMO = adverse maternal outcome; APO = adverse perinatal outcome; GFR = glomerular filtration rate; MTTD = medium time to delivery

2nd and 3rd trimester of pregnancy compared to overweight (BMI 25–30 kg/m²) and normal-weight patients with PE. The sFlt-1/PlGF ratio itself did not show any significant differences between groups [16]. Possible explanations for the decreased sFlt-1 levels in obese PE patients could be the generally higher plasma volumes as well as a higher extracellular matrix mass with heparan sulfate proteoglycans which can degrade sFlt-1 [33, 34]. TNF- α (TNF- α : tumor necrosis factor α) could also play a role, as it is a pro-inflammatory factor which is elevated in obese women while endogenous sFlt-1 expression is reduced in adipose tissue [35]. While it appears that BMI does affect sFlt-1, currently no studies have developed alternative threshold values with sufficient evidence. The sFlt-1/PlGF ratio should and can be used for the diagnostic workup of PE in obese pregnant women. But caution is advised when the risk of a potential APO is being estimated based on the ratio. A retrospective analysis was able to show that an APO

could not be ruled out despite an sFlt-1/PlGF ratio of less than 38 in the obese PE cohort [36].

Twin Pregnancies

The incidence of PE in twin pregnancies is double that reported for singleton pregnancies [37]. In twin pregnancies, the sFlt-1/PlGF ratio provides verifiable prognostic and diagnostic information [38, 39], although how chorionicity can affect test accuracy has not yet been conclusively determined [1]. Dröge et al. were able to show that the concentration levels of sFlt-1 and PlGF differ between singleton and twin pregnancies, with higher levels of sFlt-1 and PlGF observed in the twin cohort [40]. A gestational age-related reference value for normal twin pregnancies (n = 269) was recently published [17]: the data shows that sFlt-1/PlGF ratios in women with twin and singleton pregnancies are similar up until week 29 + 0 of gestation. From week 29 + 0 of gestation, however,

significantly higher sFlt-1/PIGF levels (caused by a significant increase in sFlt-1) were observed in twin pregnancies, which is why the ratio after 29 + 0 weeks of gestation appears to be less discerning when predicting PE in twin pregnancies; results should therefore be interpreted with caution [1]. One suggested explanation for the higher sFlt-1/PIGF concentrations in twin pregnancies currently being discussed is the larger placental mass [41]. Although retrospective studies with small case numbers have shown a clear correlation between the ratio and shorter times to delivery [39, 42], they did not show that the sFlt-1/PIGF ratio had an additional diagnostic value with regards to predicting an APO [39, 42, 43].

Chronic and Gestational Hypertension

Compared to patients with chronic or gestational hypertension, patients with PE have a significantly higher sFlt-1/PIGF ratio [44]. The term generally used when the sFlt-1/PIGF ratio is higher than 85 before 34 + 0 weeks of gestation or higher than 110 in or after 34 + 0 weeks of gestation is “angiogenic PE” [45]. Chronic hypertension (and gestational hypertension) is not associated per se with higher sFlt-1/PIGF levels and this allows them to be safely differentiated from superimposed PE [46]. Binder et al. showed that in pregnant women with chronic hypertension and suspected superimposed PE, adding the sFlt-1/PIGF ratio to the standard diagnostic criteria proposed by the International Society for the Study of Hypertension in Pregnancy (ISSHP) [47] significantly improved the detection rate of adverse perinatal and maternal outcomes (AMO) [48]. Accordingly, angiogenic markers can be routinely used in patients with chronic hypertension of pregnancy and gestational hypertension.

Pregnancy-related Liver Complications

Acute fatty liver of pregnancy (AFLP) and HELLP syndrome are two rare hepatic complications of pregnancy associated with high rates of maternal and fetal morbidity and mortality. In addition to the clinical characteristics, the constellation of laboratory findings is often similar and can include thrombocytopenia, hemolysis, and elevated concentrations of hepatic enzymes [49]. Differentiating HELLP syndrome from AFLP can sometimes be difficult, particularly if hypertension or PE are not additionally present. In patients with AFLP, the serum level of sFlt-1 is dramatically increased (compared to HELLP syndrome) [50, 51, 52, 53]. In addition to the Swansea criteria, an sFlt-1 value > 31 100 pg/ml appears to be an additional parameter which points to AFLP. In contrast, PIGF serum levels are only decreased in patients with HELLP syndrome. This would fit with the characteristic placental dysfunction which occurs in HELLP syndrome (and the normal placental function in AFLP) and underscores the different entities of the two pathologies [53]. Elevated sFlt-1 serum levels have also been reported outside of pregnancy in patients with chronic liver disease [54]. Studies into the role played by sFlt-1 in liver function impairment using a mouse model [55, 56] suggest that high sFlt-1 serum levels in patients with AFLP are not simply an epiphenomenon but a major cause of the disorder [53]. With regards to HELLP syndrome, in the rare case of isolated HELLP syndrome (without PE), the sFlt-1/PIGF ratio is significantly decreased [57].

Rheumatic Diseases

Patients with rheumatoid disorders such as systemic lupus erythematosus (SLE), Sjögren’s syndrome, systemic sclerosis, dermatomyositis, and rheumatoid arthritis have a higher risk of PE [58]. Making a differential diagnosis of PE and HELLP syndrome as opposed to lupus nephritis flares, vasculitis, or a renal crisis in the context of systemic sclerosis can often be difficult. In the differential diagnosis of PE, the sFlt1/PIGF ratio can be used as a sure sign of placental involvement [59]. A prospective multicenter observational study showed that in patients with SLE and/or antiphospholipid syndrome, the sFlt1/PIGF ratio in the 2nd trimester of pregnancy (in addition to other factors such as the necessity for anti-hypertensive medication) was one of the strongest predictors for an APO [60]. In its position paper “Management of Rheumatic Diseases During Pregnancy and Breastfeeding,” the section Maternal Diseases in Pregnancy of the Working Group for Obstetrics and Prenatal Medicine (AGG) noted: “Determination of [...] the sFlt1/PIGF ratio in the second and third trimester of pregnancy can be used as part of the differential diagnosis to determine prognosis when placental involvement is present” [59].

Differential Diagnosis of Thrombotic Microangiopathy/Thrombotic Thrombocytopenic Purpura

Diseases from the family of thrombotic microangiopathies (TMAs) such as aHUS and TTP can manifest in pregnancy and puerperium and present a clinical picture which overlaps with that of the obstetric disorders “severe PE” and (postpartum) “HELLP syndrome.” As the potential therapeutic options differ considerably, it would be useful to have biomarkers which are able to differentiate between TMAs and obstetric disorders and able to show whether the primary pathogenetic problem is placental dysfunction. A study published in 2023 investigated whether the sFlt1/PIGF ratio is different in patients with TTP. The study was able to show that the sFlt1/PIGF ratio was elevated in both the congenital and the acquired form of TTP in a relevant percentage of patients but not in all of them (27.3% – 52.2%). As regards the outcomes of these pregnancies (maternal and fetal survival), in patients with TTP there was no difference between cases with normal and cases with pathological angiogenesis. As TMA and placental dysfunction may develop in parallel and can mutually influence each other, angiogenic markers are not able to unequivocally identify the primary cause and have no prognostic value with regards to outcomes in cases with TTP [61].

Conclusion

Maternal and gestational comorbidities have the potential to frustrate the clinical interpretability and applicability of the sFlt-1/PIGF ratio and its constituent elements. Recent studies indicate that pregnant women with comorbidities may develop PE with a milder elevation of the sFlt-1/PIGF ratio compared to pregnant women without comorbidities. The presence of pre-existing endothelial dysfunction could be one reason for this as, under certain circum-

stances, this can lower the anti-angiogenic imbalance threshold [62]. Prospective studies with large case numbers are needed to clarify whether adapting established cut-off values would be useful for diagnosing PE and predicting an APO in specific patient cohorts.

Conflict of Interest

HS: lectures and consultancy work for Sanofi, Alexion, Roche Diagnostics and Norgine. SV: Speaker fees: Thermo Fisher Scientific, Roche Diagnostics, Comanche Biopharma, Alexion Advisory Board: Siemens, Beckman Coulter, Comanche Biopharma.

References/Literatur

- [1] Verlohren S, Brennecke SP, Galindo A et al. Clinical interpretation and implementation of the sFlt-1/PlGF ratio in the prediction, diagnosis and management of preeclampsia. *Pregnancy Hypertens* 2022; 27: 42–50. DOI: 10.1016/j.preghy.2021.12.003
- [2] Rana S, Burke SD, Karumanchi SA. Imbalances in circulating angiogenic factors in the pathophysiology of preeclampsia and related disorders. *Am J Obstet Gynecol* 2022; 226 (Suppl 2): S1019–S1034. DOI: 10.1016/j.ajog.2020.10.022
- [3] Greco M, Palumbo C, Sicuro F et al. Soluble Fms-Like Tyrosine Kinase-1 Is A Marker of Endothelial Dysfunction During Sepsis. *J Clin Med Res* 2018; 10: 700–706. DOI: 10.14740/jocmr3505w
- [4] Dupont V, Kanagaratnam L, Goury A et al. Excess Soluble fms-like Tyrosine Kinase 1 Correlates With Endothelial Dysfunction and Organ Failure in Critically Ill Coronavirus Disease 2019 Patients. *Clin Infect Dis* 2021; 72: 1834–1837. DOI: 10.1093/cid/ciaa1007
- [5] Greco M, Suppressa S, Lazzari RA et al. sFlt-1 and CA 15.3 are indicators of endothelial damage and pulmonary fibrosis in SARS-CoV-2 infection. *Sci Rep* 2021; 11: 19979. DOI: 10.1038/s41598-021-99470-y
- [6] Dewerchin M, Carmeliet P. PlGF: a multitasking cytokine with disease-restricted activity. *Cold Spring Harb Perspect Med* 2012; 2: a011056. DOI: 10.1101/cshperspect.a011056
- [7] Theilade S, Lajer M, Jorsal A et al. Evaluation of placental growth factor and soluble Fms-like tyrosine kinase 1 as predictors of all-cause and cardiovascular mortality in patients with Type 1 diabetes with and without diabetic nephropathy. *Diabet Med* 2012; 29: 337–344. DOI: 10.1111/j.1464-5491.2011.03482.x
- [8] Zeisler H, Lllurba E, Chantraine F et al. Predictive Value of the sFlt-1:PlGF Ratio in Women with Suspected Preeclampsia. *N Engl J Med* 2016; 374: 13–22. DOI: 10.1056/NEJMoa1414838
- [9] Cerdeira AS, O'Sullivan J, Ohuma EO et al. Performance of soluble fms-like tyrosine kinase-1-to-placental growth factor ratio of ≥ 85 for ruling in preeclampsia within 4 weeks. *Am J Obstet Gynecol* 2021; 224: 322–323. DOI: 10.1016/j.ajog.2020.11.007
- [10] Verlohren S, Herraiz I, Lapaire O et al. New gestational phase-specific cut-off values for the use of the soluble fms-like tyrosine kinase-1/placental growth factor ratio as a diagnostic test for preeclampsia. *Hypertension* 2014; 63: 346–352. DOI: 10.1161/HYPERTENSIONAHA.113.01787
- [11] Villalain C, Herraiz I, Valle L et al. Maternal and Perinatal Outcomes Associated With Extremely High Values for the sFlt-1 (Soluble fms-Like Tyrosine Kinase 1)/PlGF (Placental Growth Factor) Ratio. *J Am Heart Assoc* 2020; 9: e015548. DOI: 10.1161/JAHA.119.015548
- [12] Rana S, Powe CE, Salahuddin S et al. Angiogenic factors and the risk of adverse outcomes in women with suspected preeclampsia. *Circulation* 2012; 125: 911–919. DOI: 10.1161/CIRCULATIONAHA.111.054361
- [13] Rolfo A, Attini R, Nuzzo AM et al. Chronic kidney disease may be differentially diagnosed from preeclampsia by serum biomarkers. *Kidney Int* 2013; 83: 177–181. DOI: 10.1038/ki.2012.348
- [14] Wiles K, Bramham K, Seed PT et al. Placental and endothelial biomarkers for the prediction of superimposed pre-eclampsia in chronic kidney disease. *Pregnancy Hypertens* 2021; 24: 58–64. DOI: 10.1016/j.preghy.2021.02.010
- [15] Zera CA, Seely EW, Wilkins-Haug LE et al. The association of body mass index with serum angiogenic markers in normal and abnormal pregnancies. *Am J Obstet Gynecol* 2014; 211: 247.e1–247.e7. DOI: 10.1016/j.ajog.2014.03.020
- [16] Jääskeläinen T, Heinonen S, Hämäläinen E et al. FINNPEC. Impact of obesity on angiogenic and inflammatory markers in the Finnish Genetics of Pre-eclampsia Consortium (FINNPEC) cohort. *Int J Obes (Lond)* 2019; 43: 1070–1081. DOI: 10.1038/s41366-018-0217-8
- [17] De La Calle M, Delgado JL, Verlohren S et al. Gestational Age-Specific Reference Ranges for the sFlt-1/PlGF Immunoassay Ratio in Twin Pregnancies. *Fetal Diagn Ther* 2021; 48: 288–296. DOI: 10.1159/000514378
- [18] Zhang JJ, Ma XX, Hao L et al. A Systematic Review and Meta-Analysis of Outcomes of Pregnancy in CKD and CKD Outcomes in Pregnancy. *Clin J Am Soc Nephrol* 2015; 10: 1964–1978. DOI: 10.2215/CJN.09250914
- [19] Leañós-Miranda A, Campos-Galicia I, Ramírez-Valenzuela KL et al. Urinary IgM excretion: a reliable marker for adverse pregnancy outcomes in women with chronic kidney disease. *J Nephrol* 2019; 32: 241–251. DOI: 10.1007/s40620-018-0536-9
- [20] Piccoli GB, Attini R, Vasario E et al. Pregnancy and chronic kidney disease: a challenge in all CKD stages. *Clin J Am Soc Nephrol* 2010; 5: 844–855. DOI: 10.2215/CJN.07911109
- [21] Williams D, Davison J. Chronic kidney disease in pregnancy. *BMJ* 2008; 336: 211–215. DOI: 10.1136/bmj.39406.652986.BE
- [22] Wiles K, Chappell LC, Lightstone L et al. Updates in Diagnosis and Management of Preeclampsia in Women with CKD. *Clin J Am Soc Nephrol* 2020; 15: 1371–1380. DOI: 10.2215/CJN.15121219
- [23] Hirashima C, Ogoyama M, Abe M et al. Clinical usefulness of serum levels of soluble fms-like tyrosine kinase 1/placental growth factor ratio to rule out preeclampsia in women with new-onset lupus nephritis during pregnancy. *CEN Case Rep* 2019; 8: 95–100. DOI: 10.1007/s13730-018-0373-7
- [24] Verlohren S, Dröge LA. The diagnostic value of angiogenic and antiangiogenic factors in differential diagnosis of preeclampsia. *Am J Obstet Gynecol* 2022; 226 (Suppl 2): S1048–S1058. DOI: 10.1016/j.ajog.2020.09.046
- [25] Molina-Pérez CJ, Nolasco-Leaños AG, Carrillo-Juárez RI et al. Clinical usefulness of angiogenic factors in women with chronic kidney disease and suspected superimposed preeclampsia. *J Nephrol* 2022; 35: 1699–1708. DOI: 10.1007/s40620-022-01299-9
- [26] Deutsche Gesellschaft für Gynäkologie und Geburtshilfe e.V. (DGGG), Deutsche Gesellschaft für Nephrologie e.V. (DGfN). Pregnancy and renal disease. Guideline of the DGGG and OEGGG (S2k-Level, AWMF Registry No. 015/090, September 2021). 2021. Accessed June 09, 2024 at: <https://www.awmf.org/leitlinien/detail/ll/015-090.html>
- [27] Levine RJ, Thadhani R, Qian C et al. Urinary placental growth factor and risk of preeclampsia. *JAMA* 2005; 293: 77–85. DOI: 10.1001/jama.293.1.77
- [28] Matsui M, Takeda Y, Uemura S et al. Suppressed soluble Fms-like tyrosine kinase-1 production aggravates atherosclerosis in chronic kidney disease. *Kidney Int* 2014; 85: 393–403. DOI: 10.1038/ki.2013.339
- [29] Onoue K, Uemura S, Takeda Y et al. Reduction of circulating soluble fms-like tyrosine kinase-1 plays a significant role in renal dysfunction-associated aggravation of atherosclerosis. *Circulation* 2009; 120: 2470–2477. DOI: 10.1161/CIRCULATIONAHA.109.867929

- [30] Wei YM, Yang HX, Zhu WW et al. Risk of adverse pregnancy outcomes stratified for pre-pregnancy body mass index. *J Matern Fetal Neonatal Med* 2016; 29: 2205–2209. DOI: 10.3109/14767058.2015.1081167
- [31] Kwaifa IK, Bahari H, Yong YK et al. Endothelial Dysfunction in Obesity-Induced Inflammation: Molecular Mechanisms and Clinical Implications. *Biomolecules* 2020; 10: 291. DOI: 10.3390/biom10020291
- [32] WHO. Obesity: Preventing and Managing the Global Epidemic; Report of a WHO Consultation. WHO Technical Report Series Geneva, Switzerland: WHO; 2000. ; 894: 1–253
- [33] Spradley FT, Palei AC, Granger JP. Obese melanocortin-4 receptor-deficient rats exhibit augmented angiogenic balance and vasorelaxation during pregnancy. *Physiol Rep* 2013; 1: e00081. DOI: 10.1002/phy2.81
- [34] Mariman EC, Wang P. Adipocyte extracellular matrix composition, dynamics and role in obesity. *Cell Mol Life Sci* 2010; 67: 1277–1292. DOI: 10.1007/s00018-010-0263-4
- [35] Herse F, Fain JN, Janke J et al. Adipose tissue-derived soluble fms-like tyrosine kinase 1 is an obesity-relevant endogenous paracrine adipokine. *Hypertension* 2011; 58: 37–42. DOI: 10.1161/HYPERTENSIONAHA.111.171322
- [36] Karge A, Desing L, Haller B et al. Performance of sFlt-1/PlGF Ratio for the Prediction of Perinatal Outcome in Obese Pre-Eclamptic Women. *J Clin Med* 2022; 11: 3023. DOI: 10.3390/jcm11113023
- [37] von Versen-Höyneck F, Schaub AM, Chi YY et al. Increased Preeclampsia Risk and Reduced Aortic Compliance With In Vitro Fertilization Cycles in the Absence of a Corpus Luteum. *Hypertension* 2019; 73: 640–649. DOI: 10.1161/HYPERTENSIONAHA.118.12043
- [38] Rana S, Hacker MR, Modest AM et al. Circulating angiogenic factors and risk of adverse maternal and perinatal outcomes in twin pregnancies with suspected preeclampsia. *Hypertension* 2012; 60: 451–458. DOI: 10.1161/HYPERTENSIONAHA.112.195065
- [39] Binder J, Palmrich P, Pateisky P et al. The Prognostic Value of Angiogenic Markers in Twin Pregnancies to Predict Delivery Due to Maternal Complications of Preeclampsia. *Hypertension* 2020; 76: 176–183. DOI: 10.1161/HYPERTENSIONAHA.120.14957
- [40] Dröge L, Herraiz I, Zeisler H et al. Maternal serum sFlt-1/PlGF ratio in twin pregnancies with and without pre-eclampsia in comparison with singleton pregnancies. *Ultrasound Obstet Gynecol* 2015; 45: 286–293. DOI: 10.1002/uog.14760
- [41] Bdoah Y, Lam C, Rajakumar A et al. Twin pregnancy and the risk of preeclampsia: bigger placenta or relative ischemia? *Am J Obstet Gynecol* 2008; 198: 428.e1–428.e6. DOI: 10.1016/j.ajog.2007.10.783
- [42] Karge A, Seiler A, Flechsenhar S et al. Prediction of adverse perinatal outcome and the mean time until delivery in twin pregnancies with suspected pre-eclampsia using sFlt-1/PlGF ratio. *Pregnancy Hypertens* 2021; 24: 37–43. DOI: 10.1016/j.preghy.2021.02.003
- [43] Saleh L, Tahitu SIM, Danser AHJ et al. The predictive value of the sFlt-1/PlGF ratio on short-term absence of preeclampsia and maternal and fetal or neonatal complications in twin pregnancies. *Pregnancy Hypertens* 2018; 14: 222–227. DOI: 10.1016/j.preghy.2018.03.014
- [44] Verlohren S, Herraiz I, Lapaire O et al. The sFlt-1/PlGF ratio in different types of hypertensive pregnancy disorders and its prognostic potential in preeclamptic patients. *Am J Obstet Gynecol* 2012; 206: 58.e1–58.e8. DOI: 10.1016/j.ajog.2011.07.037
- [45] Deutsche Gesellschaft für Gynäkologie und Geburtshilfe e.V. (DGGG). Hypertensive Pregnancy Disorders: Diagnosis and Therapy. Guideline of the German Society of Gynecology and Obstetrics (S2k-Level, AWMF-Registry No. 015/018, March 2019). 2019. Accessed June 09, 2024 at: <https://register.awmf.org/de/leitlinien/detail/015-018#anmeldung>
- [46] Perni U, Sison C, Sharma V et al. Angiogenic factors in superimposed preeclampsia: a longitudinal study of women with chronic hypertension during pregnancy. *Hypertension* 2012; 59: 740–746. DOI: 10.1161/HYPERTENSIONAHA.111.181735
- [47] Brown MA, Magee LA, Kenny LC et al. International Society for the Study of Hypertension in Pregnancy (ISSHP). Hypertensive Disorders of Pregnancy: ISSHP Classification, Diagnosis, and Management Recommendations for International Practice. *Hypertension* 2018; 72: 24–43. DOI: 10.1161/HYPERTENSIONAHA.117.10803
- [48] Binder J, Kalafat E, Palmrich P et al. Should angiogenic markers be included in diagnostic criteria of superimposed pre-eclampsia in women with chronic hypertension? *Ultrasound Obstet Gynecol* 2022; 59: 192–201. DOI: 10.1002/uog.23711
- [49] Minakami H, Morikawa M, Yamada T et al. Differentiation of acute fatty liver of pregnancy from syndrome of hemolysis, elevated liver enzymes and low platelet counts. *J Obstet Gynaecol Res* 2014; 40: 641–649. DOI: 10.1111/jog.12282
- [50] Suzuki H, Nagayama S, Hirashima C et al. Markedly higher sFlt-1/PlGF ratio in a woman with acute fatty liver of pregnancy compared with HELLP syndrome. *J Obstet Gynaecol Res* 2019; 45: 96–103. DOI: 10.1111/jog.13786
- [51] Neuman RI, Hesselink ERM, Saleh L et al. Angiogenic markers are elevated in women with acute fatty liver of pregnancy. *Ultrasound Obstet Gynecol* 2020; 56: 465–466. DOI: 10.1002/uog.21912
- [52] Brügge M, Pecks U, Iannaccone A et al. Akute Schwangerschaftsfettleber – Fallserie einer seltenen Schwangerschaftskomplikation [Acute Fatty Liver of Pregnancy – Case Series]. *Z Geburtshilfe Neonatol* 2023; 227: 466–473. DOI: 10.1055/a-2096-6230
- [53] Trottmann F, Raio L, Amylidi-Mohr S et al. Soluble fms-like tyrosine kinase 1 (sFlt-1): A novel biochemical marker for acute fatty liver of pregnancy. *Acta Obstet Gynecol Scand* 2021; 100: 1876–1884. DOI: 10.1111/aogs.14218
- [54] Coulon S, Heindryckx F, Geerts A et al. Angiogenesis in chronic liver disease and its complications. *Liver Int* 2011; 31: 146–162. DOI: 10.1111/j.1478-3231.2010.02369.x
- [55] Oe Y, Ko M, Fushima T et al. Hepatic dysfunction and thrombocytopenia induced by excess sFlt1 in mice lacking endothelial nitric oxide synthase. *Sci Rep* 2018; 8: 102. DOI: 10.1038/s41598-017-18260-7
- [56] Uda Y, Hirano T, Son G et al. Angiogenesis is crucial for liver regeneration after partial hepatectomy. *Surgery* 2013; 153: 70–77. DOI: 10.1016/j.surg.2012.06.021
- [57] Trottmann F, Baumann M, Amylidi-Mohr S et al. Angiogenic profiling in HELLP syndrome cases with or without hypertension and proteinuria. *Eur J Obstet Gynecol Reprod Biol* 2019; 243: 93–96. DOI: 10.1016/j.ejogrb.2019.10.021
- [58] Sammaritano LR, Bermas BL, Chakravarty EE et al. 2020 American College of Rheumatology Guideline for the Management of Reproductive Health in Rheumatic and Musculoskeletal Diseases. *Arthritis Rheumatol* 2020; 72: 529–556. DOI: 10.1002/art.41191
- [59] Kuschel B, Schäfer-Graf UM, Schmidt M et al. AGG – Section Maternal Diseases in Pregnancy. Management of Rheumatic Diseases During Pregnancy and Breastfeeding: Position Paper of the Working Group for Obstetrics and Prenatal Medicine in the German Society for Gynecology and Obstetrics e.V. (AGG - Section Maternal Diseases in Pregnancy). *Geburtshilfe Frauenheilkd* 2024; 84: 130–143. DOI: 10.1055/a-2201-2680
- [60] Kim MY, Buyon JP, Guerra MM et al. Angiogenic factor imbalance early in pregnancy predicts adverse outcomes in patients with lupus and anti-phospholipid antibodies: results of the PROMISSE study. *Am J Obstet Gynecol* 2016; 214: 108.e1–108.e14. DOI: 10.1016/j.ajog.2015.09.066
- [61] Béranger N, Tsatsaris V, Coppo P et al. High sFlt-1 (Soluble fms-Like Tyrosine Kinase 1)/PlGF (Placental Growth Factor) Ratio in Pregnancy-Onset Thrombotic Thrombocytopenic Purpura. *Hypertension* 2023; 80: e140–e142. DOI: 10.1161/HYPERTENSIONAHA.123.20987
- [62] Tanner MS, de Guingand D, Reddy M et al. The effect of comorbidities on the sFlt-1:PlGF ratio in preeclampsia. *Pregnancy Hypertens* 2022; 29: 98–100. DOI: 10.1016/j.preghy.2022.06.008