



Cholangioscope laser lithotripsy used as a remedial treatment for pancreatic duct stone basket incarceration




► **Fig. 1** Computed tomography image showing chronic pancreatitis and pancreatic duct stones.



► **Fig. 2** Radiographic image showing the appearance after three sessions of extracorporeal shock wave lithotripsy.



► **Video 1** A basket became entrapped while ensnaring a pancreatic duct stone and attempts to drag the basket out after balloon expansion were unsuccessful; a cholangioscope laser is used to fragment the stone into smaller pieces and the basket is then pulled out and the stone fragments removed.



► **Fig. 3** Fluoroscopic image during endoscopic retrograde cholangiopancreatography showing: **a** balloon dilation of the narrowed pancreatic duct up to 7 mm; **b** a 6-mm dilation balloon being used to dilate the pancreatic duct.

A 60-year-old woman with chronic pancreatitis and pancreatic duct stones (► **Fig. 1**) was admitted to hospital because of abdominal pain. The patient underwent three sessions of extracorporeal shock wave lithotripsy (ESWL) and was scheduled for further endoscopic stone removal (► **Fig. 2**).

The endoscopic retrograde cholangiopancreatography (ERCP) images showed a narrowing in the head of pancreatic duct and multiple filling defects, with

the shadow of a chunky stone visible in the tail of pancreas. A papillotomy was first performed, followed by balloon dilation of the narrowed pancreatic duct up to 7 mm (► **Fig. 3 a**). A spiral stone retrieval basket (COOK) was then used to attempt to retrieve the stone; however, the basket became entrapped with the pancreatic duct stone, which was measured as having a maximum diameter of 14 mm. A dilation balloon (Boston Hurricane, 6 mm) was used to dilate the

pancreatic duct, but an attempt to drag the basket out was still unsuccessful (► **Fig. 3 b**). Subsequently, the handle of basket was cut off. A cholangioscope (SpyGlass) was then introduced via the biopsy channel of the endoscope, and a laser was introduced through the accessory channel. The U100 plus electrohydraulic lithotripsy fiber was inserted through the working tube and placed onto the surface of the stone, with intracavitary laser lithotripsy performed with a frequency of 120 Hz (► **Fig. 4**). After the stone had been fragmented into smaller pieces, the basket was pulled out under radiographic guidance. The remaining stone fragments were then successfully removed by the basket (► **Video 1**).

Stone extraction via ERCP is an effective method to treat pancreatic duct stones [1,2]. Basket incarceration is however a tricky situation that usually requires surgical treatment. In this report, we



► **Fig. 4** Image from a choledochoscope that had been introduced via the biopsy channel of the endoscope, with a laser introduced through the accessory channel to fragment the stone.

used a choledochoscope laser to fragment the stones within the incarcerated basket in the pancreatic duct, ultimately resolving the basket incarceration.

Endoscopy_UCTN_Code_TTT_1AR_2AH

Conflict of Interest

The authors declare that they have no conflict of interest.

The authors

Ping-Ping Zhang^{‡1}, Yan-Wei Lv^{‡1}, Ting Yang¹, Liang-Hao Hu¹ 

¹ Gastroenterology, Changhai Hospital, The Second Military Medical University, Shanghai, China

[‡] These authors contributed equally.

Corresponding author

Liang-Hao Hu, MD

Department of Gastroenterology, Changhai Hospital, Naval Military Medical University, Shanghai 200433, China
lianghao-hu@smmu.edu.cn

References

- [1] Nguyen-Tang T, Dumonceau JM. Endoscopic treatment in chronic pancreatitis, timing, duration and type of intervention. *Best Pract Res Clin Gastroenterol* 2010; 24: 281–298. doi:10.1016/j.bpg.2010.03.002
- [2] Moole H, Jaeger A, Bechtold ML et al. Success of extracorporeal shock wave lithotripsy in chronic calcific pancreatitis management: a meta-analysis and systematic review. *Pancreas* 2016; 45: 651–658. doi:10.1097/MPA.0000000000000512

Bibliography

Endoscopy 2024; 56: E764–E765

DOI 10.1055/a-2387-4112

ISSN 0013-726X

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited.

(<https://creativecommons.org/licenses/by/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



E-Videos is an open access online section of the journal *Endoscopy*, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high-quality video and are published with a Creative Commons CC-BY license. Endoscopy E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: <https://www.research4life.org/access/eligibility/>).

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>