

Inflammatory myofibroblastic tumor of the terminal ileum




Fig. 1 Upper gastrointestinal images of a 37-year-old male presenting with weight loss and melena. **a** Abdominal computed tomography revealed one soft-tissue mass with wall thickening of the ileum (arrow). **b** Single-balloon enteroscopy showed one well-defined round tumor within the terminal ileum (arrow). **c** The base of the tumor showed hyperemic swollen mucosa with ulceration (arrow). **d** Representative hematoxylin and eosin-stained histological sections of the tumor (original magnification $\times 100$ and $\times 400$).

A 37-year-old male was admitted to our hospital due to weight loss and passage of tarry stools. Upper gastrointestinal panendoscopy showed only gastritis. Abdominal computed tomography revealed one mass-like lesion with wall thickening of the ileum (Fig. 1a). Single-balloon enteroscopy revealed one well-defined round tumor (Fig. 1b; Video 1), measuring about $4 \times 4 \times 5$ cm in the terminal ileum, up to 30 cm from the ileocecal valve. The base of the tumor showed swollen mucosa with ulceration (Fig. 1c).

The patient underwent laparotomy with segmental resection of the ileum on the

day after enteroscopy. Histologic features of the tumor showed myofibroblastic proliferation in an inflammatory background (Fig. 1d); immunohistochemical staining for S100 and c-Kit revealed negative results. The pathologic result demonstrated inflammatory myofibroblastic tumor (IMT).

Gastrointestinal IMT can occur in the alimentary tract or mesentery/omentum [1]. The most common sites of alimentary tract IMT have been noted in ileal locations [1]. The majority of cases have been reported in childhood and early adulthood [1–3]; it is very rare in adults. This is the first reported case of an adult with

ileal IMT, which was viewed clearly by single-balloon enteroscopy. Due to a high recurrence rate, the mainstay of therapy is surgical resection [4].

Endoscopy_UCTN_Code_CCL_1AC_2AC

Competing interests: None

Video 1

Anal approach single-balloon enteroscopy showed one round tumor (about $4 \times 4 \times 5$ cm in size) within the terminal ileum.

T. Y. Huang¹, P. J. Chen¹, Y. C. Liu²,
J. S. Jin³

¹ Division of Gastroenterology, Department of Internal Medicine, Tri-Service General Hospital, Taiwan, Republic of China

² Department of Surgery, Tri-Service General Hospital, Taiwan, Republic of China

³ Department of Pathology, Tri-Service General Hospital, Taiwan, Republic of China

References

- 1 Demirkan NC, Akalin T, Yilmaz F et al. Inflammatory myofibroblastic tumor of small bowel wall in childhood: report of a case and a review of the literature. *Pathol Int* 2001; 51: 47–49
- 2 Coffin CM, Watterson J, Priest JR, Dehner LP. Extrapulmonary inflammatory myofibroblastic tumor (inflammatory pseudotumor). A clinicopathologic and immunohistochemical study of 84 cases. *Am J Surg Pathol* 1995; 19: 859–872
- 3 Biselli R, Ferlini C, Fattorossi A et al. Inflammatory myofibroblastic tumor (inflammatory pseudotumor): DNA flow cytometric analysis of nine pediatric cases. *Cancer* 1996; 77: 778–784
- 4 Difiore JW, Goldblum JR. Inflammatory myofibroblastic tumor of the small intestine. *J Am Coll Surg* 2002; 194: 502–506

Bibliography

DOI 10.1055/s-0030-1255821

Endoscopy 2011; 43: E14–E15

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

T. Y. Huang, MD, PhD

Division of Gastroenterology
Department of Internal Medicine
Tri-Service General Hospital
No. 325 Sec. 2, Cheng-Kung Road
Taipei 114
Taiwan
Republic of China
Fax: 886-2-87927139
teinyu.chun@msa.hinet.net