

## Diminutive submucosally invasive cancers of the colon and rectum

**Tab. 1** Baseline patient data, endoscopic and pathological findings, and outcomes of seven patients with diminutive submucosally invasive colorectal cancers.

Patient number sex; age	Tumor location; type; size, mm	Endoscopic diagnosis	Treatment	Pathological diagnosis	Resection margin	Growth type	Invasion depth, $\mu\text{m}$	ly	v	pN	Meta-static risk	Outcome
1 M; 67	Ascending colon; Ip; 5	Adenoma	Polypectomy then surgery	Adenocarcinoma (tub1)	Negative	Non-polypoid	1750	0	1	0	High	Lost to follow-up
2 M; 85	Transverse colon; Is; 5	Adenoma	EMR then surgery	Adenocarcinoma (tub1)	Negative	Non-polypoid	1750	0	0	0	High	Alive, disease free
3 M; 76	Transverse colon; Is; 5	Adenoma	EMR	Adenocarcinoma (tub1)	Negative	Non-polypoid	300	0	0	N/A	Low	Alive, disease free
4 F; 86	Rectosigmoid; Is; 4	Adenoma	EMR	Adenocarcinoma (tub1)	Negative	Non-polypoid	1370	0	0	N/A	High	Alive, disease free
5 M; 57	Sigmoid colon; IIc; 5	Intramucosal cancer	EMR	Adenocarcinoma (tub1)	Negative	Non-polypoid	250	0	0	N/A	Low	Alive, disease free
6 M; 53	Sigmoid colon; IIc; 5	Intramucosal cancer	EMR	Adenocarcinoma (tub1)	Negative	Non-polypoid	200	0	0	N/A	Low	Alive, disease free
7 F; 61	Sigmoid colon; IIa + IIc; 5	Submucosal invasive cancer	Surgery	Adenocarcinoma (tub1)	N/A	Non-polypoid	3500	1	0	0	High	Alive, disease free

ly, lymphatic permeation; v, vascular permeation; pN, pathological lymph node metastasis; M, male; F, female; tub1, well-differentiated tubular adenocarcinoma; EMR, endoscopic mucosal resection; N/A, not applicable.




**Fig. 1** Images from patient #2. **a** Conventional colonoscopy showing a protruded lesion (5 mm in diameter) in the transverse colon. **b** Chromoendoscopy obtained using an indigo carmine spray showing an indistinct depressed area surrounding a protrusion. **c** Pathology of the specimen from endoscopic mucosal resection (EMR) showing submucosal invasive cancer (1750  $\mu\text{m}$ ) of non-polypoid growth type.

Only a few cases of submucosally invasive colorectal cancer (SM-CRC) in diminutive colorectal polyps of  $\leq 5\text{ mm}$  have been described [1], and as yet there is no detailed knowledge of these cancers.

Pathological SM-CRCs were selected from our colonoscopy database. We examined the frequency of these lesions relative to all colonoscopy examinations and to colorectal neoplasms  $\leq 5\text{ mm}$ . The growth type of the lesions was divided into two categories: polypoid growth and non-polypoid growth [2].

A total of 32 692 colonoscopies were performed between September 2002 and December 2012, from which 5690 colorectal neoplasms were detected and treated. Only seven cases of SM-CRC occurred in lesions  $\leq 5\text{ mm}$ , accounting for only 0.5% (7/1358) of colorectal neoplasms  $\leq 5\text{ mm}$ .

Baseline patient information, endoscopic and pathological findings, and outcomes are summarized in **Table 1**. Four cases were of protruded type and three cases were of depressed type. All the protruded-type lesions were initially diagnosed before treatment as being adenomas and were resected endoscopically. All the depressed-type lesions were diagnosed before treatment as being malignant. Endoscopic treatment was indicated for six of the patients (#1 – #6), and additional surgery was performed for two patients (#1 and #2) because of histological



**Fig. 2** Images from patient #3. **a** Conventional colonoscopy showing a protruded lesion (5 mm in diameter) in the transverse colon. **b** Chromoendoscopy obtained using an indigo carmine spray showing an indistinct depressed area surrounding a protrusion. **c** Pathology of the specimen from endoscopic mucosal resection (EMR) showing submucosally invasive cancer (300  $\mu$ m) of non-polypoid growth type.

findings that suggested a high metastatic risk. All cases were of pathologically well-differentiated adenocarcinomas of non-polypoid growth type. No lymph node metastasis was evident in the surgically resected cases. The appearances in patients #2 and #3 are shown in **Fig. 1** and **Fig. 2** respectively.

The most important finding of the present study was that all diminutive SM-CRCs were pathologically diagnosed as being of non-polypoid growth type. Shimoda et al. [2] reported that colorectal cancers showing non-polypoid growth tended to in-

vade the submucosa when of a smaller size than those showing polypoid growth. Chromoendoscopy, magnified endoscopy, and image-enhanced endoscopy have been shown to be effective for the precise diagnosis of invasion depth in colorectal cancers [3, 4]. Careful endoscopic observation is strongly recommended when adopting the policies of the DISCARD trial [5, 6].

Endoscopy\_UCTN\_Code\_CCL\_1AD\_2AB

**Competing interests:** None

**Kinichi Hotta, Kenichiro Imai, Yuichiro Yamaguchi, Noboru Kawata, Masaki Tanaka, Naomi Kakushima, Kohei Takizawa, Hiroyuki Matsubayashi, Hiroyuki Ono**

Division of Endoscopy, Shizuoka Cancer Center, Shizuoka, Japan

## References

- 1 Kudo SE, Takemura O, Ohtsuka K. Flat and depressed types of early colorectal cancers: from East to West. *Gastrointest Endosc Clin N Am* 2008; 18: 581–593
- 2 Shimoda T, Ikegami M, Fujisaki J et al. Early colorectal carcinoma with special reference to its development de novo. *Cancer* 1989; 64: 1138–1146
- 3 Matsuda T, Fujii T, Saito Y et al. Efficacy of the invasive/non-invasive pattern by magnifying chromoendoscopy to estimate the depth of invasion of early colorectal neoplasms. *Am J Gastroenterol* 2008; 103: 2700–2706
- 4 Ikematsu H, Matsuda T, Emura F et al. Efficacy of capillary pattern type IIIA/IIIB by magnifying narrow band imaging for estimating depth of invasion of early colorectal neoplasms. *BMC Gastroenterol* 2010; 10: 33
- 5 Ignjatovic A, East JE, Suzuki N et al. Optical diagnosis of small colorectal polyps at routine colonoscopy (Detect InSpect ChAracterise Resect and Discard; DISCARD trial): a prospective cohort study. *Lancet Oncol* 2009; 10: 1171–1178
- 6 Rex DK, Kahi C, O'Brien M et al. The American Society for Gastrointestinal Endoscopy PIVI (Preservation and Incorporation of Valuable Endoscopic Innovations) on real-time endoscopic assessment of the histology of diminutive colorectal polyps. *Gastrointest Endosc* 2011; 73: 419–422

## Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1377400>  
*Endoscopy* 2015; 47: E2–E3  
 © Georg Thieme Verlag KG  
 Stuttgart · New York  
 ISSN 0013-726X

## Corresponding author

**Kinichi Hotta, MD**  
 Division of Endoscopy  
 Shizuoka Cancer Center  
 1007 Shimonagakubo  
 Nagaizumi-cho, Sunto-gun  
 Shizuoka 411-8777  
 Japan  
 Fax: +81-55-9895551  
 k.hotta@scchr.jp