

# Oxidative Stress and Overview of Pediatric Disease Biomarkers

Ahmet Ayaz<sup>1</sup>

<sup>1</sup>Division of Biochemistry, Department of Bioengineering, Yıldız Technical University, Istanbul, Turkey

Address for correspondence Ahmet Ayaz, PhD, Division of Biochemistry, Department of Bioengineering, Yıldız Technical University, 34349 Beşiktaş/Istanbul, Turkey (e-mail: aayaz@yildiz.edu.tr).

J Pediatr Biochem 2015;5:8–11.

## Abstract

Free radicals are small molecules enabled to react with other biological molecules. The antioxidant system in metabolism is responsible for balancing antioxidant levels and the amount of free radicals. Excessive free radicals production will result in oxidative stress. A certain amount of reactive oxygen species are required to maintain normal physiological activity. However, elevated oxidative stress levels will damage molecules and produce enzymatic malfunction. Several pediatric diseases are associated with increased oxidative stress. A biomarker is a specific molecule that acts as an indicator for a specific condition. There are some oxidative stress biomarkers currently in uses and further analysis of their application is required. Protein molecules may serve as potential biomarker according to proteomics analysis.

## Keywords

- ▶ oxidative stress
- ▶ biomarker
- ▶ proteomics
- ▶ pediatric diseases

## Introduction

### Reactive Oxygen Species and Free Radicals

Free radicals can be defined as a molecule that includes one or more unpaired electron. Reactive oxygen species (ROS) are a kind of free radical associated with a single oxygen atom having a higher reactivity than  $O_2$ .<sup>1,2</sup> They are natural by-products of cellular function and ionizing radiation. ROS can be categorized into four groups: hydroxyl radical (OH), superoxide anion ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ), and singlet oxygen ( $^1O_2$ ). The reactivity of these molecules is relatively low, yet they can produce hydroxyl radicals in the presence of transition metal by Fenton or Haber-Weiss reaction. Some other free radicals have biological importance as well. Lipid peroxide (ROOH), lipid peroxy radical (ROO), and lipid alkoxy radical are associated with lipid molecules in the cell membrane. Some species are considered as reactive nitrogen species (RNS) such as nitric oxide (NO), nitrogen dioxide ( $NO_2$ ), and peroxyxynitrite ( $ONOO^-$ ) whereas thyl radical (RS) have unpaired electron.<sup>2–4</sup>

### Oxidative Stress and Nitric Oxide System

Nitric oxide (NO) formation and oxidative stress can be classified together. NO is an uncharged, diatomic, free radical that has the ability to diffuse and affect biological molecules.<sup>1</sup> This molecule is produced by nitric oxide synthases (NOSs)

from L-arginine and nicotinamide adenine dinucleotide phosphate hydrogen (NADPH) using heme, flavin adenine dinucleotide (FAD), flavin mononucleotide (FMN), and tetrahydrobiopterin.<sup>5</sup>

NO synthases is found in the three metabolically active isoforms in mammals: Neuronal ( $NOS_1$ ), inducible ( $NOS_2$ ), and endothelial ( $NOS_3$ ).  $NOS_1$  and  $NOS_2$  are soluble and  $NOS_3$  is membrane bound. Neuronal and endothelial NO is present in various types of cells and is activated in the cell temporarily when calcium influx increases. NO binds to heme iron of soluble guanylate cyclase to produce cyclic guanosine monophosphate (cGMP) that modulates mediators such as ion channels, phosphodiesterase, and protein kinases.<sup>6</sup> Inducible NO,  $NOS_2$ , is activated against inflammatory and immunologic response. This isoform produces NO in the intracellular calcium state. This massive production of NO is about a 1,000 times more when compared other NO isoforms.<sup>7</sup>

NO can be produced by nonenzymatically and can be synthesized from nitrite at acidic pH levels under reducing conditions. Nonenzymatic NO generation may result in similar biological and enzymatically produced NO. In addition, nitrite acts as hypoxic buffer and may contribute to hypoxic vasodilation and modulation in infarction and ischemia-reperfusion tissue injury.<sup>8</sup>

### Proxynitrite Formation

NO can react with  $O_2^-$  and produce  $ONOO^-$ . Similarly, superoxide dismutase (SOD) can react with  $O_2^-$  as an antioxidant. NO reacts with  $O_2^-$  at a three times higher rate than SOD activity. The calculated diffusion rate is  $6.7 \times 10^9 M^{-1}s^{-1}$  for the NO reaction whereas SOD activity is  $2 \times 10^9 M^{-1}s^{-1}$ . A competition exists between NO and SOD in capturing  $O_2^-$ . Under the normal physiologic conditions, the amounts of NO and  $O_2^-$  are in much more lower concentrations when compared with SOD amount. Therefore, a limited amount of  $ONOO^-$  is generated in normal conditions. In pathologic states NO and  $O_2^-$  increase in the setting of low SOD activity, resulting in toxic levels  $ONOO^-$ .<sup>9</sup>

### Antioxidants

ROS has an important function in cell signaling mechanisms, especially in autocrine and paracrine systems as a defense against microbial and tumor cells.<sup>1,2</sup> Excess amounts of ROS may contribute to disease states or inflammation. The amount of ROS is restricted by several systems such as electron transfer, enzymatic removal, and ROS scavenging. Under normal physiologic states, the balance between ROS generation and ROS elimination is maintained by antioxidant enzymes and nonenzymatic antioxidants.<sup>10</sup>

Antioxidants in biological systems can be classified into three groups: enzymes, proteins, and low-molecular-weight proteins. Antioxidant enzymes are SOD, catalase, glutathione peroxidase, glutathione reductase, glutathione-S-transferase (GST), thioredoxin reductase, and hemeoxygenase. Proteins involved in antioxidant mechanisms include albumin, ferritin, transferrin, lactoferrin, ceruloplasmin, and thioredoxin (TRX). Other low-molecular-weight antioxidant molecules include bilirubin, tocopherols, carotenoids, ubiquinol/ubiquinone, ascorbate, glutathione, cysteine, and urate. An imbalance between antioxidant system and free radicals metabolism results in disequilibrium and causes oxidative stress.<sup>1,2,11,12</sup>

It is well known that high ROS and oxidative stress are correlated with some pathological conditions such as male infertility, atherosclerosis, hypertension, renal failure, neurodegeneration, carcinogenesis, and other inflammatory and degenerative conditions.<sup>13</sup>

### Oxidative Stress Biomarkers

Biological molecules are prone to damage by reactive oxygen species. Oxidative stress may damage carbohydrates, lipids, proteins, and nucleic acids.<sup>14</sup> ROS are small molecules with a short half-life that requires special techniques for in vivo detection biomarkers because oxidative would not only be useful to detect oxidative damage but would also help determine the source of oxidative stress.<sup>3,15,16</sup> A specific biomarker would be important to determine byproduct of oxidation and to prevent further destructive effect of oxidative stress.

Oxidative stress biomarkers can be generally classified into two groups:

1. Formation of modified molecules by the effect of ROS
2. Measurement of enzymes or antioxidants

Biomarkers from body fluids such as blood, urine, semen, or cerebrospinal fluid<sup>17</sup> would allow the monitoring of oxidative stress in vivo that cannot be done with other invasive tests. A test that could visualize biomarkers by using a fluorescent probe would be useful, but it is not practical as a routine laboratory test.<sup>2,8</sup>

The first group of biomarkers includes ROS-generated byproducts that affect other biologically important molecules. This interaction can be mediated by scission, cross-linking, or covalent binding between free radicals and biological molecules. Some byproducts can be repaired or removed, but the others can stay longer if the molecule is the part of an intra- or extracellular compartment. ROS mainly attacks carbohydrates, lipids, proteins, and nucleic acids.<sup>18</sup> The effect of ROS on these molecules can be measured by analyzing stable byproducts produced by them. Malondialdehyde-lysine, 4-hydroxy-2-nonenal-lysine, acrolein-lysine, 8-hydroxy-2-deoxyguanosine (8-OHdG), carboxymethyl-lysine, pentosidine, and nitrite/nitrate are some clinically applicable biomarker for diagnosis of oxidative stress.<sup>19</sup>

Better biomarkers are necessary to improve diagnosis, targeted therapy, and therapeutic response to overcome the effect of oxidative stress. A proteomics approach based on mass spectrometry may serve as a potential protein biomarker for a clinical sample. New discoveries in proteomics methodology may allow for predicting clinically useful biomarkers with further advances in qualification, verification, assay optimization, validation, and commercialism.<sup>20</sup> Discovery of protein biomarker for oxidative stress may serve as a potential biomarker for clinical tests.

Studies including children are needed to investigate role of oxidative stress in pediatric disease. It could provide a better approach to understanding the effect of oxidative stress in pediatric diseases when compared with adolescents. However, a limited number of studies have reported on oxidative stress in similar children.<sup>21,22</sup> In most studies, oxidative stress parameters were identified in blood samples that allow glutathione peroxidase and glutathione reductase activity,<sup>23</sup> antioxidant vitamins,<sup>24</sup> ubiquinol/ubiquinone,<sup>25</sup> and SOD and catalase activities.<sup>26</sup> Schock and colleagues reported oxidative stress biomarkers including antioxidant vitamins in 83 healthy children.<sup>21</sup> Kaufmann and colleagues identified urinary levels of F2-isoprostane in 342 children with less than 7 years population.<sup>24</sup> In conclusion, further studies are needed to determine reference values of oxidative stress in pediatric urine samples.

### Oxidative Stress in Pediatric Diseases

Oxidative stress is associated with the progression and pathogenesis of many diseases.<sup>6,27-30</sup> Certain drugs such as analgesic, anticancer drugs may also contribute tissue damage by increasing oxidative stress.<sup>31-33</sup> Oxidative stress biomarkers have been measured in cerebrospinal fluid, joint fluid,<sup>34</sup> nasal lavage fluid,<sup>35</sup> several types of tissues,<sup>11,12,36-38</sup> or combination with blood or urine samples.<sup>39-41</sup>

Oxidative stress levels were conventionally measured by some analytical techniques including high-performance liquid chromatography (HPLC) and gas chromatography-mass

spectrometry (GC-MS).<sup>42-44</sup> Recent developments in monoclonal antibody technology allows for specific biomarkers via enzyme-linked immuno assay (ELISA) or Western blot analysis.<sup>3,42</sup> ELISA and Western blot provides a strong biomarker candidate as other analytical tests labor intensive and costly.<sup>28,45,46</sup> Recently, proteins were studied as potential biomarker to detect effect of oxidative stress on metabolism.<sup>47-49</sup> This technique needs to be improved to be useful in clinical medicine to provide rapid results. The majority of oxidative stress parameters in different pathologic states are performed by high-throughput ELISA techniques.<sup>45,46,50</sup>

A respiratory burst is an oxidative burst that results in the rapid production of reactive oxygen species, including superoxide and hydrogen peroxide. NADPH oxidase is an enzyme that has an ability to produce highly reactive free radicals. It is particularly involved in vascular disease and spontaneous recombination with other molecules leading to free radical production. A systematic cascade initiates the reaction of superoxide molecules with NO. This reaction results in the formation of peroxynitrite and reduces necessary bioactive NO. In addition, the superoxide anion peroxynitrite and other reactive oxygen species cause some pathology as a result of oxidation of protein and lipids. Free radicals affect redox signaling pathways and posttranslational modifications.<sup>51</sup> Many studies have reported a correlation between infection and the respiratory burst effect on biologically important molecules. Cemek and colleagues identified an oxidative burst in children with hepatitis A.<sup>52</sup> Caksen and colleagues identified lipid peroxidation and antioxidant status in children with tonsillitis.<sup>53</sup> Bayiroğlu monitored increased lipid peroxidation and antioxidant status in pediatric gastroenteritis patients.<sup>54</sup>

Proteins may serve as better potential biomarkers when compared with genome.<sup>55</sup> Proteomics is an evolving field in the evaluation of diseases. In addition to protein identification, quantitative proteomics provides information about the physiologic and pathologic function of identified proteins as well as their cellular localization and biological processes.<sup>56</sup> Differential expression of selected proteins may be used as a biomarker as a noninvasive diagnostic tool.<sup>47,57,58</sup> Understanding the protein expression level may be the key to understanding the cellular processes and/or pathways of a disease.<sup>59</sup>

Oxidative stress has destructive effect on several diseases by decomposing of biologically important molecules. Carbohydrates, lipids, nucleic acids, and proteins are prone to be denatured by the effect of free radicals and cause a loss of function. Specific biomarkers are needed to improve noninvasive diagnostic tools and develop better treatments. Recent developments have shown that proteins are potential molecules with the specificity detect qualitatively and quantitatively oxidation. Differential expression of candidate proteins can be determined by proteomic tools and quantitative proteomics may help find that target protein.

## Conclusion

The challenge for further analysis in pediatric disease is elucidating molecular mechanism behind disorders and

engenders the oxidative stress in mechanisms. Before accepting antioxidant therapy in clinical practice, comprehensive studies must be conducted to evaluate oxidative stress biomarkers with together clinical endpoints of patients.

This review provided updated information about noninvasive value of evaluating oxidative stress in pediatric medicine. Many of the mentioned biomarkers can be measured in urine samples where protein extraction will be needed for proteomics analysis. Measurement of these parameters has great potential for managing and treating oxidative stress-related pediatric diseases.

## References

- Halliwell B, Zhao K, Whiteman M. Nitric oxide and peroxynitrite. The ugly, the uglier and the not so good: a personal view of recent controversies. *Free Radic Res* 1999;31(6):651-669
- Toyokuni S. Reactive oxygen species-induced molecular damage and its application in pathology. *Pathol Int* 1999;49(2):91-102
- Noiri E, Tsukahara H. Parameters for measurement of oxidative stress in diabetes mellitus: applicability of enzyme-linked immunosorbent assay for clinical evaluation. *J Investig Med* 2005;53(4):167-175
- Kato M, Minakami H, Kuroiwa M, et al. Superoxide radical generation and Mn- and Cu-Zn superoxide dismutases activities in human leukemic cells. *Hematol Oncol* 2003;21(1):11-16
- Lavine JE, Schwimmer JB. Nonalcoholic fatty liver disease in the pediatric population. *Clin Liver Dis* 2004;8(3):549-558, viii-ix
- Renke J, Popadiuk S, Korzon M, Bugajczyk B, Wozniak M. Protein carbonyl groups' content as a useful clinical marker of antioxidant barrier impairment in plasma of children with juvenile chronic arthritis. *Free Radic Biol Med* 2000;29(2):101-104
- Hayashi M, Araki S, Arai N, et al. Oxidative stress and disturbed glutamate transport in spinal muscular atrophy. *Brain Dev* 2002;24(8):770-775
- Gladwin MT, Schechter AN, Kim-Shapiro DB, et al. The emerging biology of the nitrite anion. *Nat Chem Biol* 2005;1(6):308-314
- Giray B, Hincal F. Oxidative DNA base damage, antioxidant enzyme activities and selenium status in highly iodine-deficient goitrous children. *Free Radic Res* 2002;36(1):55-62
- Djordjević VB, Stanković T, Cosić V, et al. Immune system-mediated endothelial damage is associated with NO and antioxidant system disorders. *Clin Chem Lab Med* 2004;42(10):1117-1121
- Cemek M, Dede S, Bayiroğlu F, Caksen H, Cemek F, Yuca K. Oxidant and antioxidant levels in children with acute otitis media and tonsillitis: a comparative study. *Int J Pediatr Otorhinolaryngol* 2005;69(6):823-827
- Cemek M, Dede S, Bayiroğlu F, Caksen H, Cemek F, Mert N. Oxidant and non-enzymatic antioxidant status in measles. *J Trop Pediatr* 2007;53(2):83-86
- Bryan NS. Nitrite in nitric oxide biology: cause or consequence? A systems-based review. *Free Radic Biol Med* 2006;41(5):691-701
- Tsukahara H. Oxidative stress biomarkers in pediatric medicine. In: Laher I, ed. *Systems Biology of Free Radicals and Antioxidants*. Germany Springer Berlin Heidelberg Publishing; 2014:689-715
- Uchida K. Current status of acrolein as a lipid peroxidation product. *Trends Cardiovasc Med* 1999;9(5):109-113
- Jakus V, Rietbrock N. Advanced glycation end-products and the progress of diabetic vascular complications. *Physiol Res* 2004;53(2):131-142
- Kharitonov SA, Barnes PJ. Biomarkers of some pulmonary diseases in exhaled breath. *Biomarkers* 2002;7(1):1-32
- Reichenbach J, Schubert R, Schindler D, Müller K, Böhles H, Zielen S. Elevated oxidative stress in patients with ataxia telangiectasia. *Antioxid Redox Signal* 2002;4(3):465-469

- 19 Shimizu T, Satoh Y, Syoji H, et al. Effects of parenteral lipid infusion on DNA damage in very low birth weight infants. *Free Radic Res* 2002;36(10):1067–1070
- 20 Rifai N, Gillette MA, Carr SA. Protein biomarker discovery and validation: the long and uncertain path to clinical utility. *Nat Biotechnol* 2006;24(8):971–983
- 21 Schock BC, Young IS, Brown V, et al. Antioxidants and protein carbonyls in bronchoalveolar lavage fluid of children: normal data. *Pediatr Res* 2001;49(2):155–161
- 22 Stolarska M, Mlynarski W, Zalewska-Szewczyk B, Bodalski J. Cytoprotective effect of amifostine in the treatment of childhood neoplastic diseases—a clinical study including the pharmacoeconomic analysis. *Pharmacol Rep* 2006;58(1):30–34
- 23 Erden-Inal M, Sunal E, Kanbak G. Age-related changes in the glutathione redox system. *Cell Biochem Funct* 2002;20(1):61–66
- 24 Kauffman LD, Sokol RJ, Jones RH, Awad JA, Rewers MJ, Norris JM. Urinary F2-isoprostanes in young healthy children at risk for type 1 diabetes mellitus. *Free Radic Biol Med* 2003;35(6):551–557
- 25 Miles MV, Horn PS, Tang PH, et al. Age-related changes in plasma coenzyme Q10 concentrations and redox state in apparently healthy children and adults. *Clin Chim Acta* 2004;347(1–2):139–144
- 26 Andreazza AC, Bordin DL, Salvador M. Thiobarbituric acid reactive substances, seric superoxide dismutase and catalase activities in healthy subjects. *Clin Chim Acta* 2005;362(1–2):192–194
- 27 Manary MJ, Leeuwenburgh C, Heinecke JW. Increased oxidative stress in kwashiorkor. *J Pediatr* 2000;137(3):421–424
- 28 Tsukahara H, Haruta T, Todoroki Y, et al. Oxidant and antioxidant activities in childhood meningitis. *Life Sci* 2002;71(23):2797–2806
- 29 Németh I, Boda D. Xanthine oxidase activity and blood glutathione redox ratio in infants and children with septic shock syndrome. *Intensive Care Med* 2001;27(1):216–221
- 30 Cardin R, Saccoccio G, Masutti F, Bellentani S, Farinati F, Tiribelli C. DNA oxidative damage in leukocytes correlates with the severity of HCV-related liver disease: validation in an open population study. *J Hepatol* 2001;34(4):587–592
- 31 Parra Cid T, Conejo García JR, Carballo Alvarez F, de Arriba G. Antioxidant nutrients protect against cyclosporine A nephrotoxicity. *Toxicology* 2003;189(1–2):99–111
- 32 Conklin KA. Coenzyme q10 for prevention of anthracycline-induced cardiotoxicity. *Integr Cancer Ther* 2005;4(2):110–130
- 33 Kennedy DD, Ladas EJ, Rheingold SR, Blumberg J, Kelly KM. Antioxidant status decreases in children with acute lymphoblastic leukemia during the first six months of chemotherapy treatment. *Pediatr Blood Cancer* 2005;44(4):378–385
- 34 Lotito AP, Muscará MN, Kiss MH, et al. Nitric oxide-derived species in synovial fluid from patients with juvenile idiopathic arthritis. *J Rheumatol* 2004;31(5):992–997
- 35 Sienra-Monge JJ, Ramirez-Aguilar M, Moreno-Macias H, et al. Antioxidant supplementation and nasal inflammatory responses among young asthmatics exposed to high levels of ozone. *Clin Exp Immunol* 2004;138(2):317–322
- 36 Cemek M, Caksen H, Cemek F, et al. Investigation of antioxidant status in children with acute otitis media and tonsillitis. *Int J Pediatr Otorhinolaryngol* 2004;68(11):1381–1385
- 37 Ayvaz S, Inan M, Aksu B, et al. Desferrioxamine effectively attenuates testicular tissue at the end of 3 h of ischemia but not in an equal period of reperfusion. *J Pediatr Urol* 2014;10(3):550–558
- 38 Caksen H, Ozkan M, Cemek M, Cemek F. Oxidant and antioxidant status in children with subacute sclerosing panencephalitis. *J Child Neurol* 2014;29(11):1448–1452
- 39 Müller T, Langner C, Fuchsichler A, et al. Immunohistochemical analysis of Mallory bodies in Wilsonian and non-Wilsonian hepatic copper toxicosis. *Hepatology* 2004;39(4):963–969
- 40 Yilmaz T, Koçan EG, Besler HT. The role of oxidants and antioxidants in chronic tonsillitis and adenoid hypertrophy in children. *Int J Pediatr Otorhinolaryngol* 2004;68(8):1053–1058
- 41 Hargreaves IP, Sheena Y, Land JM, Heales SJ. Glutathione deficiency in patients with mitochondrial disease: implications for pathogenesis and treatment. *J Inher Metab Dis* 2005;28(1):81–88
- 42 Tsukahara H, Toyo-Oka M, Kanaya Y, et al. Quantitation of glutathione S transferase-pi in the urine of preterm neonates. *Pediatr Int* 2005;47(5):528–531
- 43 Wong RH, Kuo CY, Hsu ML, et al. Increased levels of 8-hydroxy-2-deoxyguanosine attributable to carcinogenic metal exposure among schoolchildren. *Environ Health Perspect* 2005;113(10):1386–1390
- 44 Stephensen CB, Marquis GS, Douglas SD, Wilson CM. Plasma cytokines and oxidative damage in HIV-positive and HIV-negative adolescents and young adults: a protective role for IL-10? *Free Radic Res* 2005;39(8):859–864
- 45 Omata N, Tsukahara H, Ito S, et al. Increased oxidative stress in childhood atopic dermatitis. *Life Sci* 2001;69(2):223–228
- 46 Sebeková K, Podracká L, Blazíček P, Syrová D, Heidland A, Schinzel R. Plasma levels of advanced glycation end products in children with renal disease. *Pediatr Nephrol* 2001;16(12):1105–1112
- 47 Sharma R, Agarwal A, Mohanty G, et al. Proteomic analysis of seminal fluid from men exhibiting oxidative stress. *Reprod Biol Endocrinol* 2013;11:85
- 48 Ming X, Stein TP, Brimacombe M, Johnson WG, Lambert GH, Wagner GC. Increased excretion of a lipid peroxidation biomarker in autism. *Prostaglandins Leukot Essent Fatty Acids* 2005;73(5):379–384
- 49 Hata I, Kaji M, Hirano S, Shigematsu Y, Tsukahara H, Mayumi M. Urinary oxidative stress markers in young patients with type 1 diabetes. *Pediatr Int* 2006;48(1):58–61
- 50 Matsubasa T, Uchino T, Karashima S, et al. Oxidative stress in very low birth weight infants as measured by urinary 8-OHdG. *Free Radic Res* 2002;36(2):189–193
- 51 Inoguchi T, Sonta T, Tsubouchi H, et al. Protein kinase C-dependent increase in reactive oxygen species (ROS) production in vascular tissues of diabetes: role of vascular NAD(P)H oxidase. *J Am Soc Nephrol* 2003;14(8, Suppl 3):S227–S232
- 52 Cemek M, Dede S, Bayiroglu F, Caksen H, Cemek F, Mert N. Relationship between antioxidant capacity and oxidative stress in children with acute hepatitis A. *World J Gastroenterol* 2006;12(38):6212–6215
- 53 Caksen H, Dede S, Cemek M, Dulger H, Cemek F. Evaluation of antioxidant status in children with acute bacterial meningitis and encephalitis. *Int J Neurosci* 2003;113(11):1497–1504
- 54 Bayiroğlu F, Cemek M, Caksen H, Cemek F, Dede S. Altered antioxidant status and increased lipid peroxidation in children with acute gastroenteritis admitted to a pediatric emergency service. *J Emerg Med* 2009;36(3):227–231
- 55 Agarwal A, Durairajanayagam D, Halabi J, Peng J, Vazquez-Levin M. Proteomics, oxidative stress and male infertility. *Reprod Biomed Online* 2014;29(1):32–58
- 56 Zhou T, Zhou ZM, Guo XJ. Bioinformatics for spermatogenesis: annotation of male reproduction based on proteomics. *Asian J Androl* 2013;15(5):594–602
- 57 Milardi D, Grande G, Vincenzoni F, Castagnola M, Marana R. Proteomics of human seminal plasma: identification of biomarker candidates for fertility and infertility and the evolution of technology. *Mol Reprod Dev* 2013;80(5):350–357
- 58 Sharma R, Agarwal A, Mohanty G, et al. Proteomic analysis of human spermatozoa proteins with oxidative stress. *Reprod Biol Endocrinol* 2013;11:48
- 59 Amaral A, Castillo J, Ramalho-Santos J, Oliva R. The combined human sperm proteome: cellular pathways and implications for basic and clinical science. *Hum Reprod Update* 2014;20(1):40–62