

Predictive Factors, 30-Day Clinical Outcomes, and Costs Associated with Cerebrospinal Fluid Leak in Pituitary Adenoma Resection

Adish Parikh¹ Arjun Adapa¹ Stephen E. Sullivan² Erin L. McKean^{1,2}

¹Department of Otolaryngology Head and Neck Surgery, University of Michigan, Ann Arbor, Michigan, United States

²Department of Neurosurgery, University of Michigan, Ann Arbor, Michigan, United States

Address for correspondence Erin McKean, MD, MBA, Department of Otolaryngology Head and Neck Surgery, University of Michigan, 1500 E Medical Center Dr, Ann Arbor, MI, 48109, United States (e-mail: elmck@med.umich.edu).

J Neurol Surg B 2020;81:43–55.

Abstract

Cerebrospinal fluid (CSF) leak is a complication of endoscopic endonasal pituitary adenoma resection. Previous studies examining complications of pituitary adenoma resection have not examined associations of an exhaustive list of clinical and financial variables with CSF leak. We designed a retrospective analysis of 334 consecutive patients that underwent endoscopic endonasal pituitary adenoma resection at a single institution over 5 years, analyzing associations between CSF leak and demographic data, operative data, comorbidities, clinical complications and outcomes, costs, charges, and payments. Of the 20 preoperative variables studied, none were positively associated with CSF leak in between-groups comparison, although multivariate analysis revealed an association with a history of radiation to the skull base (odds ratio [OR], 8.67; 95% confidence interval [CI], 0.94–57.03; $p < 0.05$). CSF leak was associated with a significantly higher rate of postoperative diabetes insipidus ($\Delta = 33.4\%$, $p = 0.040$) and increased length of stay after operation in between-groups comparison. Multivariate analysis on postoperative variables revealed significant associations between CSF leak and intracerebral hemorrhage (OR, 17.44; 95% CI, 0.65–275.3; $p < 0.05$) and postoperative intracranial infection (OR, 28.73; 95% CI, 2.04–438.7; $p < 0.05$). Also, CSF leak was associated with significantly higher costs ($\Delta = \$15,643$, $p < 0.05$) and hospital charges ($\Delta = \$46,026$, $p < 0.05$). Operating room time, room and board, and supplies and implants were the strongest cost drivers. This study highlights the difficulty of utilizing preoperative variables to predict CSF leak, the clinical complications and outcomes of leak, and the financial subcategories that drive the costs, charges, and payments associated with this complication.

Keywords

- ▶ endoscopic endonasal approach
- ▶ pituitary adenoma
- ▶ cerebrospinal fluid leak
- ▶ CSF leak
- ▶ costs
- ▶ complications
- ▶ pituitary surgery

Introduction

The endoscopic endonasal approach is a common method for pituitary adenoma resection. Many studies have reinforced that the operation is safe with overall favorable postoperative outcomes relative to the microscopic approach.^{1–3} However, there are notable risks for complications such as diabetes

insipidus (DI), cranial nerve injury, and cerebrospinal fluid (CSF) leak, among others, that can lead to morbidity, mortality, and substantial financial costs.

One of the most common complications is CSF leak due to the intracranial maneuvering needed for the resection of pituitary tumors.^{4–6} Piek et al in Germany found a 5.36% leak rate in their 168 patients, whereas Grotenhuis in the

received
September 22, 2018
accepted after revision
January 11, 2019
published online
February 18, 2019

© 2020 Georg Thieme Verlag KG
Stuttgart · New York

DOI <https://doi.org/10.1055/s-0039-1679896>
ISSN 2193-6331.

Netherlands found a 10.7% rate in 412 patients.^{7,8} As discussed by Lobatto et al identifying patients with risk factors for complications like CSF leak is imperative for appropriate consideration of preoperative planning and timing of the operation, patient counseling, consultation or referral to a center of excellence, alternative treatments, and postoperative management such as short versus long stay.⁹

Some literature has identified risk factors for postoperative CSF leak. Nishioka et al suggest that prior transsphenoidal surgery and prior radiotherapy to the skull are risk factors, while Dlouhy et al and Ivan et al found that increased body mass index (BMI) was correlated with CSF leak.^{10–12} Lobatto et al conducted a systematic review on general endoscopic pituitary surgical outcomes and identified 14 studies that examined risk factors for CSF leaks, specifically.⁹ Only interventricular extension was consistently associated with CSF leak, although extension was defined in different ways in different studies. Some studies have identified younger age,^{13,14} increased BMI,^{11,13,14} female gender,^{13,14} increased tumor size,¹⁵ and surgeons' learning¹⁶ curve as less consistent associations with CSF leaks. Again, differences in variable definitions make these comparisons difficult (e.g., age categories, tumor size categories, and learning curve).

From our literature review, there are no studies specifically examining clinical outcomes of CSF leak, such as mortality, 30-day emergency department (ED) visits, and associations with other complications, with the exception of one study noting that the length of stay (LOS) was twice as long in patients with CSF leakage.¹¹ There is some research on the costs of CSF leak following endonasal endoscopic resection of pituitary adenoma. Piek et al noted that without CSF leak, costs were €14,079.37 per case with a reimbursement of €14,856.05 per case (net +€780 per case). The costs were increased in CSF leaks—€25,286.60/case with a reimbursement of €25,499.39 (net +€210 per case).⁷ It is not surprising that a complication like CSF leak leads to higher costs. To help alleviate this increased cost, it is helpful to identify the drivers of this cost so providers and administrators can help mitigate the financial burden. Cost drivers of CSF leak are currently poorly understood. A retrospective study of 27 patients that underwent pituitary adenoma resection found that total operating room (OR) and total bed-assignment costs constituted the highest expense, representing more than 60% of the cost of hospitalization.¹⁷ Because CSF leak has been shown to be associated with longer LOS and has the potential for reoperation, these cost drivers may be the biggest sources of increased cost in CSF leak patients, although this has not been specifically studied.

Despite the prevalence of postoperative CSF leak in endoscopic endonasal pituitary surgery, many studies into the risk factors for CSF leak appear to assess singular factors, such as BMI or prior pituitary operation, rather than exploring an exhaustive list of demographic data and comorbidities within a large set of patients. In addition, there is limited literature on the cost drivers of postoperative CSF leak in this patient group, particularly in the United States, and 30-day postoperative outcomes in these patients.

The main objective of our study was to create a large database of patient-specific variables, as well as postoperative outcomes and costs to better understand the causes and results, both clinically and financially, of CSF leak. Identification of those at risk for CSF leak could potentially change perioperative management. In addition, because previous studies have had controversial conclusion on risk factors of CSF leak, we hope that our study can better clarify the impact of these variables (i.e., gender, BMI, and tumor size). Also, due to the breadth of the variables in our database, we hope to study potential risk factors and associations with CSF leak that have not been previously studied (i.e., medical comorbidities and 30-day postoperative outcomes) and to create a more robust multivariate linear regression analysis. Lastly, understanding cost and charge data may allow quantification of the financial impact of CSF leak via specific cost drivers, allowing us to pinpoint specific variables that hospitals may focus their efforts on to minimize costs of CSF leak and optimize value of the operation.

Methods

We designed a retrospective analysis on 334 consecutive patients that underwent endoscopic endonasal pituitary adenoma resection at the University of Michigan by the single team of senior authors from March 2011 to March 2016. Patients with craniopharyngiomas or Rathke's cleft cyst and patients that underwent concomitant or early staged craniotomy (i.e., for intracranial hemorrhage at presentation unrelated to the pituitary adenoma, sellar masses biopsied endoscopically but resected via craniotomy or giant adenomas requiring planned multistaged approaches) were not included in this study. Using our secure internal health system self-service data search tool, demographic data including age, sex, and race was collected. Next, a search on this cohort using a text-recognition tool, Electronic Medical Record Search Engine (EMERSE), allowed for collection of operative data, comorbidities, and clinical complications/outcomes.¹⁸ EMERSE allowed us to search for "word bundles" in a patient's chart to find matching words and phrases. A word bundle included several synonyms for the variable of interest. After identifying matches, the corresponding clinical note was reviewed for confirmation of the variable (i.e., patient has a history of myocardial infarction in their preoperative note). Preoperative variables are shown in ► **Table 1** and postoperative variables, including complications and outcomes, are shown in ► **Table 2**. The vast majority of variables were binary or categorical (i.e., patient did or did not have a stroke in the postoperative period). Only postoperative data within 30 days of the operation date was collected.

Financial data on our cohort, including total hospital costs (what it costs internally to provide the care, allocated directly and indirectly), charges (the price tag to the payor for hospital and professional services), and payments (what we are actually paid for our care) for the hospital billing encounters associated with the pituitary operations, was collected via the Michigan Medicine Corporate Finance

Table 1 List of preoperative variables measured, definitions, and variable conditions. A total of 20 variables were measured and 14 of these are comorbidities

Variable name	Definition	Conditions of the variable
Age	Date of operation minus date of birth	Linear: 0–infinity
Gender	Gender	Male, female, NA
Race	White, Black, Hispanic, or other race	White, Black, Hispanic, or other race
Tumor type	Primary type of pituitary tumor based on pathology	Nonsecreting, prolactinoma, acromegaly, Cushing’s disease, TSH-oma
Macroadenoma	Tumor size > 1cm	Yes, No, NA
H/o MI	History of past myocardial infarction prior to operation	Yes, No, NA
H/o CHF	History of past congestive heart failure prior to operation	Yes, No, NA
H/o stroke	History of past stroke prior to operation	Yes, No, NA
Preop visual field cut	Field cut (no light) within 1 month prior to surgery	Yes, No, NA
Preop decreased visual acuity	Decreased visual acuity prior to surgery (uncorrectable via eye glasses or contacts)	Yes, No, NA
H/o pulmonary disease	Chronic lung disease requiring maintenance medications	Yes, No, NA
Diabetes mellitus	Diabetes mellitus diagnosis prior to operation	Yes, No, NA
Renal disease	CKD stage 3 or higher	Yes, No, NA
Liver disease	Liver disease diagnosis prior to operation	Yes, No, NA
Blood thinners	Use of aspirin, warfarin, dabigatran, apixaban, rivaroxaban, clopidogrel, or heparin noted in pre-op history and physical	Yes, No, NA
H/o radiation to skull base	Radiation therapy to the skull prior to operation	Yes, No, NA
H/o prior pituitary surgery	Pituitary surgery prior to operation	Yes, No, NA
Immune suppression	Chronic immune suppression including current (within 30 days preop) chemotherapy; autoimmune disease or transplant requiring immunosuppressants, or a diagnosis of primary immunodeficiency	Yes, No, NA
BMI	Body mass index	Linear 0–infinity
Obesity	BMI >30 kg/m ²	Yes, No, NA

Abbreviations: CKD, chronic kidney disease; CHF, chronic heart failure; MI, myocardial infarction; TSH, thyroid-stimulating hormone.

Office. These data, further defined in **Table 3**, encompass expenses from the admission date through the discharge date, up to and including 30 days postoperatively. In addition, these financial metrics were subdivided for each patient in our cohort by subcategory (e.g., anesthesia, OR time, and supplies/implants).

Of the 334 patients, 13 patients were found to have had a postoperative CSF leak within 30 days of operation. We stratified our database into two different groups: “CSF leak group” (*n* = 13) and “non-CSF leak group” (*n* = 321). This allowed us to compare the variables shown in **Tables 1** and **2** between these two groups to identify significant differences. Due to unequal sizes of the two groups, a two-tailed *t*-test assuming unequal variances was utilized to compare variables. To our knowledge, this is the most extensive set of clinical variables collected to analyze relationships with CSF leak in patients that underwent pituitary

adenoma resection. There are 20 preoperative variables, 15 postoperative complications (not including requirement of desmopressin for DI and type of CSF leak intervention), and 6 surgical outcome variables. To evaluate the relative contributions and associations of certain preoperative, operative, demographic, postoperative clinical and surgical variables to CSF leak, univariate logistic regression was performed for each variable against CSF leak. All variables with *p* < 0.2 were included in a multivariate logistic regression. We considered a two-sided *p* < 0.05 to be statistically significant. Data were analyzed using R Version 3.5.2 and R Studio 1.1.456.

Lastly, total costs and drivers were compared between the two groups. Average total hospital costs, charges, and payments were calculated for the CSF leak group and non-CSF leak group. To calculate the average net profit for a CSF leak patient, average hospital cost was subtracted from average

Table 2 List of postoperative variables: Variables collected in the 30-day postoperative period, definitions, and variable conditions. The list is categorized into complications and clinical outcomes. Fifteen complications (and two interventions) and six clinical outcome variables were measured

Variable name	Definition	Conditions of the variable
Complication within 30 days postoperative		
Diabetes insipidus	A urine-specific gravity of 1.005 or less, a urine osmolality less than 200 mOsm/kg, or physician diagnosis in note	Yes, No, NA
Desmopressin required	Desmopressin required postoperative to treat diabetes insipidus	Yes, No, NA
CSF leak	Clinical diagnosis of a CSF Leak	Yes, No, NA
CSF leak Intervention	What was used to treat CSF leak	NA, surgery, lumbar drain, acetazolamide, conservative management, MD
Intracerebral hemorrhage	Hemorrhage or hematoma requiring additional scans or other interventions	Yes, No, NA
Tension pneumocephalus	Pneumocephalus confirmed on imaging and causing compressive symptoms	Yes, No, NA
Intracranial infection	Clinical diagnosis of meningitis, cerebritis, or abscess requiring intervention (including delayed discharge for observation)	Yes, No, NA
Decreased visual field	Visual field cut not present preoperatively	Yes, No, NA
Decreased visual acuity	Decreased visual acuity as a result of surgery	Yes, No, NA
Cranial nerve Injury	New cranial neuropathy not present preoperatively	Yes, No, NA
MI	Clinical diagnosis of myocardial infarction based on EKG and/or troponins	Yes, No, NA
Stroke	Clinical diagnosis of stroke	Yes, No, NA
DVT or PE	Deep vein thrombosis or pulmonary embolism diagnosis	Yes, No, NA
Severe arrhythmia	Arrhythmia in the postoperative period requiring any intervention (new meds, delayed discharge, cardioversion)	Yes, No, NA
Hyponatremia	Serum sodium level < 136 mEq/L or diagnosis in note	Yes, No, NA
Hypernatremia	Serum sodium level >145 mEq/L or diagnosis in note	Yes, No, NA
Respiratory failure	Respiratory compromise requiring noninvasive or invasive positive pressure ventilation	Yes, No, NA
Clinical Outcomes within 30 days postoperative		
Length of stay	How many nights spent in the hospital after surgery prior to initial discharge	Linear 0–infinity
30-day re-admission	Readmission into Michigan Medicine or other hospital prior to 30 days postoperative, as noted within Michigan Medicine EMR	Yes, No, NA
Total length of stay	Total number of nights spent in hospital after surgery, including all readmissions prior to 30 days postoperatively	Linear 0–infinity
30-day ED visits	Emergency department visit at Michigan Medicine or any other ED, as noted within Michigan Medicine EMR	Yes, No, NA
30-day reoperation	Reoperation to treat CSF leak, remove residual tumor, or to treat bleeding	Yes, No, NA
30-day mortality	Death after surgery	Yes, No, NA

Abbreviations: CSF, cerebrospinal fluid; EKG, electrocardiogram; EMR, electronic medical record; MI, myocardial infarction.

Table 3 Definitions of financial metrics used in analysis

Financial metric	Definition
Hospital charges	Sum of all charges sent to the patient/payor for the services the hospital provides to the patient
Professional charges	Component of hospital charges; clinical caregiver charges (e.g., physician, resident) for a hospital billing encounter sent to the patient/payor
Hospital costs	Total money that the hospital spends on a patient's encounter; sum of direct and variable costs for a hospital billing encounter
Hospital direct variable costs	Component of hospital costs. What the hospital spends on worker supplies, patient care supplies, diagnostic and therapeutic supplies, and medications
Hospital payments	Insurance and patient payments received by the hospital

hospital payment. Next, hospital charges, costs, direct variable costs (a component of total costs), payments, and professional charges (a component of total hospital charges) were then subdivided to a more granular level, which we call a “financial subcategory” or simply “subcategory,” in the CSF leak group to examine the contribution of the subcategory to each variable. Of note, for one CSF leak patient treated in 2011, encounter details by financial subcategory could not be obtained due to differences in data entry prior to 2012. As such, this patient was excluded from the financial analysis separated by subcategory ($n = 12$). Finally, to examine the cost of CSF leak interventions used in our CSF leak group, average hospital charges, costs, professional charges, and LOS were compared between patients treated with surgery alone, lumbar drain, or a combination of both. These data included all 13 patients in the CSF leak group.

Results

Our total cohort included 334 patients. Two-hundred patients (60%) had nonsecreting pituitary adenomas. Of the 134 patients (40%) with secreting tumors, acromegaly was the most common disease ($n = 77$, 23%). Two-hundred eighty-two patients (84%) were diagnosed with macroadenoma. The average age at operation was 52.96 years old, 55% of patients were male, and 83% of patients were white. Summary of tumor characteristics and demographics is shown in ► **Table 4**.

Significant Differences in Preoperative Variables

After stratifying our total cohort for CSF leak ($n = 13$, 3.9%), we found several significant differences between the leak groups versus the nonleak group (► **Table 5**). There were four comorbidities that were significantly *less* common in the CSF leak group—history of congestive heart failure ($p < 0.001$), history stroke ($p = 0.001$), history of liver disease ($p < 0.001$), and immunosuppression ($p < 0.001$). Of the 13 patients with CSF leak, none had a history of any of these four conditions. Although the differences between the two groups were highly

Table 4 Tumor characteristics and basic demographics on 334 patients

	Amount	Percentage of total
Nonsecreting	200	59.9%
Secreting	134	40.1%
Acromegaly	77	23.1%
Cushing's disease	41	12.3%
Prolactinoma	14	4.2%
TSH-oma	2	0.6%
Total	334	
Macroadenomas	282	84.4%
White people	276	82.6%
Male	183	54.8%
Average age (years)	52.96	

Abbreviation: TSH, thyroid-stimulating hormone.

significant, these conditions were still fairly uncommon in the non-CSF leak group—4% had a history of CHF, 3% had stroke, 4% had liver disease, and 4% had immunosuppression. Of the 14 comorbidities measured, none were significantly more common in the CSF leak group.

Significant Differences in Clinical Complications

There were also three postoperative clinical complications that occurred significantly less often in the CSF leak group—venous thromboembolic event (VTE) ($p = 0.014$), decreased postoperative visual field ($p = 0.008$), and respiratory failure ($p = 0.025$)—as shown in ► **Table 5**. In the CSF leak group, none of the 13 patients had any of these complications. These complications were fairly uncommon in the nonleak group as well—2% had VTE, 2% had decreased postoperative visual field, and 2% had respiratory failure. One postoperative clinical complications occurred significantly more often in the CSF leak group—DI ($\Delta = 33.4\%$, $p = 0.040$).

Significant Differences in Surgical Outcomes

Surgical outcomes had several notable differences (► **Table 5**). LOS after operation in the leak group was 5.08 days compared with 2.23 days in the nonleak group ($\Delta = 2.84$ days, $p < 0.001$). Total LOS, which includes any additional days spent in the hospital on readmission, was 5.08 days in the leak group versus 2.79 days in the nonleak group ($\Delta = 2.29$ days, $p < 0.001$). Reoperation rate was significantly higher in the CSF leak group (54 vs. 2%, $p = 0.004$). Thirty-day readmission rate was significantly less in the leak group ($\Delta = 13\%$, $p < 0.001$). Readmission LOS was less in the leak group ($\Delta = 0.55$ days, $p < 0.001$). Postoperative ED visits were also fewer in the leak group ($\Delta = 18\%$, $p < 0.001$).

Notable Nonsignificant Differences

Contrary to some literature on CSF leak, there were no significant differences when comparing tumor characteristics (i.e., % macroadenoma, nonsecreting type, and Cushing's disease). Acromegaly was the only condition approaching significance

Table 5 Significant differences found between the CSF leak and nonleak groups: Categorized by preoperative variables, complications, and outcomes

Preop variables	H/o CHF	0.00%	4.36%	-4.36%	0.000
	H/o stroke	0.00%	2.80%	-2.80%	0.003
	Liver disease	0.00%	4.36%	-4.36%	0.000
	Immune suppression	0.00%	3.74%	-3.74%	0.000
Postoperative clinical complications	Diabetes insipidus	46.15%	12.77%	33.38%	0.040
	DVT or PE	0.00%	1.87%	-1.87%	0.014
	Postoperative visual field change	0.00%	2.18%	-2.18%	0.008
	Respiratory failure	0.00%	1.56%	-1.56%	0.025
Postoperative surgical outcomes	LOS (days)	5.08	2.23	2.84	0.000
	30-day readmission	0.00	0.13	-13.40%	0.000
	Readmission LOS (days)	0.00	0.55	-0.55	0.000
	Total LOS (days)	5.08	2.79	2.29	0.001
	30-Day ED visits	0.00%	17.76%	-17.76%	0.000
	30-Day Reoperation	53.85%	2.18%	51.67%	0.004

Abbreviations: CHF, chronic heart failure; ED, emergency department; DVT, deep vein thrombosis; LOS, length of stay; PE, pulmonary embolism.

(7.7% in leak group vs. 23.7% in nonleak group, $p = 0.07$). In addition, unlike some of the studies mentioned in the introduction, demographic data was not significantly different, including age at operation, % of male, % of white race, and BMI (►Table 6). Notably, in our patient population, the average BMI is in the “obese” range for both leak and nonleak patients.

Preoperative Variables that Predict CSF Leak Complication in Multivariate Analysis

Multivariate binary logistic regression demonstrated that of the various types of tumors studied, none were significantly

Table 6 Notable nonsignificant differences in tumor characteristics and patient demographics

	CSF leak	Non-CSF leak	Difference	p Value
Nonsecreting	53.8%	60.1%	-6.3%	0.68
Secreting	46.2%	39.9%	6.3%	0.68
Acromegaly	7.7%	23.7%	-16.0%	0.07
Cushing’s disease	15.4%	11.8%	3.5%	0.74
Prolactinoma	15.4%	3.7%	11.6%	0.29
TSH-oma	0.00%	0.62%	-0.6%	0.16
Macroadenomas	92.3%	84.1%	8.2%	0.32
White race	92.3%	82.2%	10.1%	0.23
Male	61.5%	54.5%	7.0%	0.63
Average age (years)	56.5	52.8	3.7	0.44
BMI	34.66	32.50	2.16	0.44

Abbreviations: BMI, body mass index; CSF, cerebrospinal fluid; TSH, thyroid-stimulating hormone.

associated with CSF leak, although prolactinomas approached significance (odds ratio [OR], 6.35; 95% confidence interval [CI], 0.71–40.45; $p = 0.062$). Of the comorbidities studied, only patients with a history of radiation to the skull base were found to have an association with CSF leak (OR, 8.67; 95% CI, 0.94–57.03; $p = 0.031$). Importantly, both of these variables had CIs that included the null hypothesis.

Postoperative Outcomes Variables Associated with CSF Leak in Multivariate Analysis

Multivariate binary logistic regression showed that patients with postoperative intracranial infection had 28.73 times greater odds of also having postoperative CSF leak (OR, 28.73; 95% CI, 2.04–438.7; $p = 0.012$). Interestingly, the presence of CSF leak did not affect LOS (OR, 1.00; 95% CI, 1.09–1.67; $p = 0.005$). Notably, both postoperative intracranial infection and LOS had CIs that did not include the null hypothesis. In addition, patients with postoperative intracerebral hemorrhage had 17.44 greater odds of also having CSF leak (OR, 17.44; 0.65–275.3; $p = 0.044$), although the CI of this relationship included the null hypothesis. While patients with postoperative DI had 11.02 increased odds of also having postoperative CSF leak, this finding only approached significance (OR, 11.02; 95% CI, 0.45–102.7; $p = 0.064$). ORs and CIs for variables with $p < 0.9$ in multivariate analysis are displayed (►Table 7).

Financial Consequences of CSF Leak

Average hospital costs for patients with CSF leak compared with those without CSF leak were \$33,684 and 18,041, respectively ($p < 0.05$). Average hospital charges for patients with CSF leak compared with those without CSF leak were \$99,281 and 53,255, respectively ($p < 0.05$). Average hospital payments for patients with CSF leak compared with those without CSF leak were \$30,467 and 22,868, respectively, but this difference was

Table 7 Univariate and multivariate logistic regression models of variables associated with CSF leak

		Univariate analysis			Multivariate analysis			
		p-Value	Odds ratio	95% Confidence interval	p-Value	Odds ratio	95% Confidence interval	
Operative data	Tumor type	Macroadenoma	0.443					
		Acromegaly	0.214					
		Cushing's disease	0.691					
		TSH-oma	0.994					
		Prolactinoma	0.061			0.062	6.35	0.71–40.45
	Nonsecreting	0.632						
	LOS (days)	0.0003	1.36	1.16–1.60	0.005	1.00	1.09–1.67	
Demographics	Age at surgery	0.415						
	Male gender	0.609						
	White race	0.360						
Preoperative variables	History of MI	0.703						
	History of CHF	0.993						
	History of stroke	0.991						
	Preop visual field cut	0.538						
	Preop visual acuity	0.510						
	History of pulmonary disease	0.659						
	Diabetes mellitus	0.813						
	Renal disease	0.143				0.462	3.06	0.08–39.25
	Liver disease	0.993						
	Blood thinners	0.911						
	History of radiation to skull base	0.048	5.12	0.74–22.20	0.031	8.67	0.94–57.03	
	History of prior pituitary surgery	0.526						
	Immunosuppression	0.900						
BMI	0.335							
Postoperative clinical complications	Diabetes insipidus	0.002	5.87	1.81–18.47	0.064	11.02	0.45–102.7	
	Desmopressin required	0.004	5.64	1.62–17.99	0.702	0.61	0.06–14.59	
	Intracerebral hemorrhage	0.040	13.29	0.59–148.4	0.044	17.44	0.65–275.3	
	Symptomatic pneumocephalus	0.990						
	Intracranial infection	0.001	29.08	3.24–259.8	0.012	28.73	2.04–438.7	
	Cranial nerve injury	0.993						
	DVT or PE	0.993						
	Severe arrhythmia	0.426						
	Hyponatremia	0.053				0.088	3.54	0.79–15.01
	Hypertatremia	0.151				0.496	0.52	0.07–3.17
	Postoperative visual field change	0.992						
	Postoperative visual acuity change	0.569						
	Respiratory failure	0.990						
Postoperative surgical outcomes	30-Day readmission	0.992						
	Readmission LOS	0.992						
	Total LOS	0.017	1.15	0.997–1.24	0.149	1.13	0.94–1.32	
	30-Day ED visits	0.991						
	30-Day reoperation	0.992						
	30-Day mortality	0.992						

Abbreviations: BMI, body mass index; CHF, chronic heart failure; CSF, cerebrospinal fluid; DVT, deep vein thrombosis; ED, emergency department; LOS, length of stay; MI, myocardial infarction; PE, pulmonary embolism; TSH, thyroid-stimulating hormone.

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.

Table 8 Hospital charges, costs, and payments for cases with CSF leak

Financial metrics (\$)	CSF leak group (n = 13)	Non-CSF leak group (n = 322)	p-Value
Average total hospital charges	99,281	53,255	<0.05
Average total hospital costs	33,684	18,041	<0.05
Average total hospital payments	30,467	22,868	0.14
Average net hospital profit (payment-cost)	-3,217 (loss)	4,385 (gain)	0.06

Abbreviation: CSF, cerebrospinal fluid.

not significant ($p = 0.14$). The average net profit (payment minus cost) from patient cases with CSF leak compared with those without CSF leak was $-\$3,217$ (loss) and $4,385$ (gain), respectively, which approached significance ($p = 0.06$). Average hospital costs, charges, and payments for CSF leak and nonleak patients are shown in **Table 8**.

When identifying cost drivers and variability in cost by category, the average percentage of contribution of a particular financial subcategory (e.g., pharmacy, magnetic resonance imaging/magnetic resonance angiography [MRI/MRA], and supplies/implants) to a cost metric was calculated and plotted (**Fig. 1**). Those subcategories for which less than 2% of either hospital charges, costs, or payments was attributed were not plotted but are listed in **Table 9**.

For hospital charges, OR time ($23.12 \pm 5.35\%$), supplies/implants (e.g., absorbable packing or tissue glue) ($18.78 \pm 4.43\%$), and room and board ($15.78 \pm 5.62\%$) were the top contributors in order (**Fig. 1**). In contrast, room and board constituted the highest percentage of hospital total costs ($27.60 \pm 10.02\%$), followed by time-based OR ($21.99 \pm 6.10\%$) and supplies/implants ($19.42 \pm 7.13\%$). For hospital payments, time-based OR, supplies/implants, and room and board were the primary contributors in order, with

the same percentages as for hospital charges (**Fig. 1**). Notably, room and board are the largest hospital cost driver yet only the third highest factor in both hospital charge and hospital payment.

Hospital direct variable costs, a component of total costs, were primarily driven by supplies/implants ($26.78 \pm 10.63\%$), room and board ($23.37 \pm 8.86\%$), and time-based OR ($22.32 \pm 7.96\%$). Professional charges, a component of total hospital charges, were also examined by a different set of subcategories (less than those listed in **Fig. 1**). These data are plotted in **Fig. 2**. The primary driver for professional charges was surgical procedures ($77.69 \pm 5.10\%$). Among the other subcategories, anesthesia was the only one that represented greater than 10% of average professional charges ($11.00 \pm 3.77\%$).

We next compared the various modalities of CSF leak intervention—lumbar drain, surgery alone, or a combination of both—for average total hospital costs, average total hospital charges, average professional charges, average total hospital payments, average profit, and average LOS. As shown in **Table 10**, average total hospital costs for lumbar drain ($n = 6$), surgery alone ($n = 4$), and combination ($n = 3$) were $\$26,353$, $31,285$, and $51,547$, respectively. Average hospital




Fig. 1 Average hospital charges, payments, and costs for cases with a cerebrospinal fluid leak complication, broken down by financial subcategory. Abbreviations: CT, computed tomography; CTA, computed tomography angiography; MRI, magnetic resonance imaging; MRA, magnetic resonance angiography.

Table 9 Financial subcategories with <2% financial contribution to hospital charges, payments, and costs

Financial subcategories
Drug admin
Hospital outpatient department
Blood/transfusion
Cardiac services
EEG
Emergency services
EKG
Peripheral vascular laboratories
Physical therapy/occupational therapy/speech
Reference laboratories
Market priced services
Neurology
Nuclear medicine/PET
Ophthalmologic evaluation services
Ultrasound
X-ray

Abbreviations: EEG, electroencephalogram; EKG, electrocardiogram; PT/OT, physical therapy/occupational therapy; PET, positron emission tomography.

charges for lumbar drain, surgery alone, and combination were \$84,610, 84,648, and 148,144, respectively. Average professional charges for lumbar drain, surgery alone, and combination were \$35,556, 37,729, and 42,333, respectively. Average total hospital payments for lumbar drain, surgery alone, and combination were \$19,056, 40,897, and 39,385,

respectively. Average profit for lumbar drain, surgery alone, and combination was -\$7,927 (loss), 9,612 (gain), and -12,162 (loss), respectively. Finally, average LOS (in days) for patients treated with lumbar drain, surgery alone, and combination were 6.3 ± 1.4 , 3.8 ± 1.0 , and 4.3 ± 2.1 , respectively. A comparison of differences among lumbar drain, surgery alone, and combination treatment revealed a significant difference only between lumbar drain and surgically treated patients in regard to average total hospital payments, average profit, and average LOS.

Discussion

CSF leak is one of the most common complications following pituitary adenoma resection.⁴⁻⁶ Although its prevalence is well understood, research into its risk factors, clinical implications, and costs is limited. Surgeons operating on patients with pituitary adenomas can potentially change their clinical management to decrease the frequency of CSF leak and reduce costs. Our study, to date, examines the largest set of clinical correlates and financial drivers of CSF leak in patients that have undergone pituitary adenoma resection.

It is worth discussing our standard procedures in this group. This analysis is for patients operated on by a single team of surgeons over a 5-year time course which was during rapid expansion of the program and was fairly early in our team's learning curve. Now, we perform well over 100 pituitary operations annually as a single team, and standardization of processes has increased over time. During the time period of study, the operations and postoperative care always included intraoperative image-guidance, frequently included fat graft reconstruction with or without pedicled flap, routinely included early inpatient postoperative MRI to evaluate completeness of resection, and routinely included postoperative




Fig. 2 Professional charges for cases with a cerebrospinal fluid leak complication, broken down by financial subcategory. Surgical procedures are highest contributor to professional charges. Abbreviations: CT, computed tomography; CTA, computed tomography angiography; MRI, magnetic resonance imaging; MRA, magnetic resonance angiography.

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.

Table 10 Hospital charges, total costs, professional charges and LOS among CSF leak interventions

Financial metrics (\$)	Lumbar drain (n = 6)	Surgery alone (n = 4)	Surgery and lumbar drain (n = 3)	p-Values		
				Drain versus surgery alone	Drain versus both	Surgery alone versus both
Average total hospital charges	84,610	84,648	148,133	1.0	0.5	0.5
Average professional charges	35,556	37,729	42,333	0.6	0.4	0.7
Average total hospital costs	26,353	31,285	51,547	0.3	0.4	0.5
Average total hospital payments	19,056	40,897	39,385	<0.05	0.4	0.9
Average total hospital profit (payment-cost)	-7,297	9,612	-12,162	<0.05	0.3	0.9
Average LOS (days)	6.3 ± 1.4	3.8 ± 1.0	4.3 ± 2.1	<0.01	0.2	0.7

Abbreviations: CSF, cerebrospinal fluid; LOS, length of stay.

staged endoscopic sinonasal debridement. Patients were sent to a neurosurgical care floor directly, with no intensive care unit stay. In more recent years, volume has increased, operative time has decreased, and the approach has become less invasive. It will be important for us to examine changes to patient outcomes and financial implications based on changes to standard processes over time.

The incidence of CSF leak in our study population of 3.9% was comparable to previous studies examining CSF leak following transsphenoidal surgery.^{7,19} Of the 20 preoperative variables studied between groups (CSF leak vs. nonleak), none were significantly more common in the CSF leak group (→ **Table 1**). This is in contrast to some previous studies. Ivan et al found that nonideal BMI was associated with postoperative CSF leak in a cohort of 98 consecutive patients operated on at the University of California, San Francisco (11 patients had CSF leaks).¹² It is important to recognize that in this study, a statistical significance was only found when these patients were grouped as either abnormal BMI (<18.5 or >25 kg/m²) or normal BMI (18.5–25 kg/m²). BMI itself was not an independent risk factor. In a study in Singapore, where obesity benchmarks are set at lower BMIs, BMI was found to be an independent risk factor for CSF leak: 27.0 kg/m² versus 24.6 kg/m²; *p* = 0.018.²⁰ There were three other studies that noted a correlation between BMI and CSF leak,^{11,13,14} although Lobatto et al conducted a systematic review and rated these studies with moderate-to-high bias.⁹ In our study, there was no significant difference between the groups. BMI in leak versus nonleak group was 32.62 versus 32.54, *p* = 0.97. In addition, multivariate analysis did not suggest an association between BMI and CSF leak. Potential confounders include demographic differences between the patient populations in these studies. For example, in the state of Michigan, where most our patients are from, the obesity rate was 31.6% in 2016.²¹ In California (where most of Ivan et al's cohort is likely from), the obesity rate was only 25.0%.

Similarly to the between-groups analysis that did not reveal significantly more common preoperative variables in the CSF leak group, multivariate analysis did not find significant contributions of these same variables to CSF leak. However, the association of prolactinomas with CSF approached significance (OR, 6.35; 95% CI, 0.71–40.45; *p* = 0.062). The association

between prolactinoma and postoperative CSF leak has not been clearly demonstrated before. In one literature review on spontaneous and medically induced CSF, the authors noted larger and invasive adenomas cause tumor expansion into surrounding dural structures, and some functional adenomas, such as growth hormone-secreting adenomas and prolactinomas, frequently invade sellar and infrasellar spaces. Nonfunctioning macroadenomas, meanwhile, have a tendency to invade the suprasellar space. In cases with additional invasion of the arachnoid and/or brain parenchyma, there is greater risk for development of a CSF fistula. As long as the tumor is large enough to occlude the opening, thereby “plugging” this opening, the escape of CSF can be prevented. However, reduction in tumor size, either due to medical or surgical therapy, can inadvertently lead to CSF escape.²² Nevertheless, a larger cohort study, especially one aimed at specifically investigating the effect of tumor type on postoperative CSF leak, will help elucidate further whether prolactinoma is an independent factor for postoperative CSF leak.

In regard to prior treatment, Nishioka et al found that previous transsphenoidal surgery and skull base radiation were each independent risk factors for postoperative CSF rhinorrhea.¹⁰ The rate of prior transsphenoidal surgery in our population was 23 versus 16% (*p* = 0.59) in the leak and nonleak group, respectively. History of prior radiation to the skull base in our patients was 15 versus 3% (*p* = 0.28) in the leak versus nonleak group, respectively. Although the difference is large, it did not reach significance in between-groups comparison. However, in multivariate analysis, previous skull base radiation was associated with greater odds of postoperative CSF leak, thereby supporting the findings in literature (OR, 8.67; 95% CI, 0.94–57.03; *p* = 0.031). Again, the CI included the null hypothesis. A larger cohort study is necessary to adequately delineate the significance of prior skull base radiation in future postoperative CSF leak complications.

DI was found to be significantly higher in the CSF group in between-group comparisons (Δ = 34%, *p* = 0.04). Conceptually, the increased frequency of DI in the leak group makes sense. A more aggressive surgical approach to resect pituitary tumor can affect the posterior pituitary, ultimately suppressing antidiuretic hormone secretion and leading to DI. However, in multivariate analysis, the association of CSF

leak and DI only approached significance (OR, 11.02; 95% CI, 0.45–102.7; $p = 0.064$).

For certain comorbidities, the prevalence within this population was too low to find significant association with CSF leak. We found that several comorbidities were actually associated with a decreased risk of CSF leak—history of CHF, stroke, liver disease, and immunosuppression. In fact, of the 13 patients that had CSF leak, none of them had any of these four comorbidities. To our knowledge, these variables' effects on CSF leak likelihood have not been specifically studied. Physiologically, we do not believe that any of these four conditions are related to CSF leak. Rather, we believe, again, that the significant differences between the leak and nonleak group are due to our cohort not being large enough to capture these rare comorbidities. To illustrate this, let us consider history of stroke, which was very rare in the nonleak group at 3%. If we assume that this rate is the same in the leak group of 13 patients, we expect less than one patient to have a history of stroke ($3\% \times 13 \text{ patients} = 0.6 \text{ patients}$). Not surprisingly, we have zero patients with a history of stroke in the leak group. Similarly, the rate of CHF, liver disease, and immunosuppression were all below 4.4% in the nonleak group and none were present in our 13 patients that had CSF leak. For this reason, we doubt that these variables are actually protective against CSF leak, despite the strong statistical significances shown in ►Table 5. As discussed previously, multivariate analysis showed that only history of radiation to the skull base had a p -value less than 0.5 out of all the preoperative variables, although the CI included the null hypothesis. We recommend that future studies examine larger cohorts to better identify if these variables, particularly history of radiation to the skull, are truly different in leak and nonleak patients.

There were three postoperative complications that appeared to be less common in the CSF leak group—VTE, decreased postoperative visual field, and respiratory failure. Again, we noticed that of the 13 patients with CSF leak, none of them had any of these three complications. A similar phenomenon explained in the paragraph above likely explains these findings, as well. All three of these complications were very rare in the nonleak group; the rates were all under 3%. Because of these very low rates in the nonleak, it should not be surprising that we do not see any of these complications in the CSF leak group despite the significant p -values when comparing the two groups.

Multivariate analysis, on the other hand, identified two significant postoperative complications—intracranial infection and intracerebral hemorrhage. Postoperative intracranial infection was found to have 28.73 times greater odds of also having postoperative CSF leak (OR, 28.73; 95% CI, 2.04–438.7; $p = 0.012$). It is known that CSF leak increases the risk of meningitis after skull base surgery, with one study demonstrating a relative risk of 14.6.²³ One explanation for this association may be attributed to lumbar drainage-related infection. Of note, the rate of intracranial infections in our CSF leak population was 15 versus 1% in the nonleak group, but between-groups analysis revealed a nonsignificant difference ($p = 0.18$). Given the small sample size and that 9 out of the 13 (69%) CSF leak patients had

lumbar drainage, it is difficult to compare the effect of lumbar drainage on the rate of intracranial infections in our population. Nevertheless, our analysis shows that the odds of developing intracranial infection are higher in CSF leak patients. These findings shed light on existing unanswered questions regarding management of CSF leak, such as whether patients with CSF leak should be treated with broad-spectrum antibiotics and whether surgeons should consider surgical repair over lumbar drain to treat CSF leak and prevent infection.

In between group comparisons, the CSF leak group had 8% postoperative intracerebral hemorrhage compared with 1% in the nonleak group, which was not significant ($p = 0.38$). However, in multivariate analysis, intracerebral hemorrhage was found to be significantly associated with CSF leak, although the CI included the null hypothesis (OR, 17.44; 95% CI, 0.65–275.3; $p = 0.044$). From our literature review, there is no study specifically comparing the association of hemorrhage with CSF leak. However, Lobatto et al's systematic review highlights the major risk factors for bleeding in transsphenoidal resection of pituitary adenomas—most notably internal carotid artery (ICA) injury. Risk factors for ICA injury are prior radiation and intraventricular extension.^{9,13} Interestingly, these variables have also been found to be risk factors for CSF leak.^{9,13,14} With these risk factors in mind, it would not be surprising to have a patient with both CSF leak and intracerebral hemorrhage.

Regarding the LOS, as expected, the CSF leak group had an average hospital stay after surgery that was 2.8 days longer. As shown in ►Table 7, univariate analysis also suggested this association (OR, 1.36; 95% CI, 1.16–1.60; $p < 0.001$). However, in multivariate analysis, the OR decreased to 1.0 (OR, 1.00; 95% CI, 1.09–1.67; $p = 0.005$), suggesting that the influence of CSF leak itself on LOS becomes less influential in the context of the additional variables studied here.

Readmission rates were also significantly less in leak patients (0 vs. 13%, $p < 0.001$), but this variable was not significantly associated with CSF leak in univariate analysis. It is possible that there are fewer readmissions in the leak group since they receive additional monitoring due to their longer initial hospital stay. Nonetheless, almost all postoperative leaks are found early, prior to discharge, thus not requiring readmission. Reoperation rate was unsurprisingly found to be significantly higher in the CSF leak group (54 vs. 2%, $p = 0.004$) in between-groups comparison, but this was not a significant variable in univariate analysis. Nevertheless, in our cohort, CSF leak was the most common reason for reoperation. We occasionally reoperate for residual functional adenoma and postoperative epistaxis

In the process of understanding and managing postoperative CSF leaks, it is important to consider not only the comorbidities, surgical complications, and clinical outcomes but also the financial consequences. Previous studies have noted that CSF leak patients had higher costs, primarily due to OR time and room & board—findings that are interesting but also not very surprising to most surgeons. The literature currently lacks depth to properly analyze the more stratified cost drivers and delves very little into charges and payments. Without these more granular analyses, it is difficult for a

hospital to identify an area of their practice to address to decrease costs and increase financial value.

It is worth briefly considering cost accounting. As noted in the methods section, financial data collected on our cohort included total hospital costs (what it costs internally to provide the care, allocated directly and indirectly), charges (the price tag to the payor/patient for hospital and professional services), and payments (what the hospital is actually paid for the care provided) for the hospital billing encounters associated with the pituitary operations. Charges and payments are relatively straightforward, but cost has many challenges in its assessment. Costs are assigned by the health system directly (patient-specific) or indirectly (“overhead”). Direct costs may be variable (e.g., the cost for an implant used for the specific patient; this is variable in its use and only charged when used, so the cost is essentially “all or none”) or fixed (e.g., the cost to provide nursing care to the patient on a specific unit for the day or session; the hospital will pay the entire day’s salary no matter how much nursing time is actually spent). Indirect costs comprise the costs to “keep the lights on,” and these are generally allocated evenly across patients or units. We have not evaluated here *how* these costs are specifically assigned by the health system and have only recorded the costs as provided to us.

We found that average total hospital costs and hospital charges for patients with CSF leak were significantly higher than those for patients without this complication, and hospital costs for CSF leak cases were primarily driven by room and board charges. While CSF leak resulted in increased hospital payments, the net profit (payment minus cost) from cases with CSF leak was a loss, compared with a gain in profit from those without CSF leak (►Table 8). One study examining cost of postoperative CSF leak across multiple neurosurgical procedures found that, on average, procedures involving a CSF leak cost 141% more than those that did not result in postoperative CSF leak.⁸ Another study examining costs associated with CSF leak in cerebellopontine angle surgery found that the median cost of a CSF leak was \$50,401.²⁴ Our data did not show as large of a difference between the two groups; however, the cost of leak was still 87% higher than the nonleak group.

The high cost of a CSF leak complication may be, in part, explained by associations with other complications and subsequent management. We found that postoperative intracranial infection, for example, was significantly associated with CSF leak, and indeed this is corroborated in literature.²³ Downstream complications past 30-days in the postoperative period were not followed in the present study. We suggest examining larger cohorts of patients in a longer postoperative window to better study the relationships of CSF leak with associated clinical complications and subsequent costs.

Examining financial drivers yielded primary contributions to total hospital charges, total hospital costs, hospital payments, and hospital direct variable costs by supplies/implants, room and board, and time-based OR subcategories. More generally, these subcategories can be attributed to

surgery and hospital stay. Not surprisingly, as evidenced by the longer LOS for patients with CSF leak, room and board were a primary cost driver. Interestingly, however, room and board charges were only the third largest contributor to hospital charges and payments, with time-based OR and supplies/implants as the first and second largest contributors, respectively, to both categories. Therefore, measures to decrease patient LOS, understandably, would reduce hospital costs and improve profits.

Our hypothesis in terms of management of CSF leaks was that early surgical intervention would actually be less costly overall than management with a lumbar drain, assuming that early surgical intervention would decrease LOS relative to prolonged lumbar drainage. Indeed, we found that surgical management of CSF leak decreases average LOS but raises hospital costs, although the latter difference was not significant (►Table 10). Surgery also increases hospital payments, with a positive net profit compared with lumbar drain alone (profit of \$9,612 vs. -7,927, $p < 0.05$). The method of treatment for postoperative CSF leak—lumbar drain versus surgery versus combination—was not significantly different in hospital charges, professional charges, or costs. However, surgical management of postoperative CSF leak required a shorter admission and received higher payments than lumbar drain, thus leading to a more favorable financial outcome for the hospital. If a patient can be clinically managed effectively with either surgery or lumbar drain, it may be prudent to opt for the surgical option.

Surgical decision-making during an operation, which can be dependent on factors such as surgeon experience, specialty training (neurosurgery vs. otolaryngology), and extensiveness of the operation, may have implications on the probability of CSF leak and ensuing financial burdens. When it comes to surgeon experience, more extensive experience with transsphenoidal surgery, defined by more than 500 previous operations, has been found, understandably, to be associated with a decreased percentage of postoperative complications, including CSF leak.¹⁹ As a result, we hypothesize that more advanced training and experience leads to lower costs associated with CSF leak. More research needs to be done to specifically explore this topic, and we will continue to follow our own outcomes over time.

Albeit retrospective, this study has, importantly, delineated cost drivers associated with CSF leak in transsphenoidal surgery for pituitary tumors. Previous studies have either focused on specific neurosurgical procedures or examined financial metrics without deconstructing the subcategories for which costs are attributed. Our study is limited as it was conducted in a single-center retrospective method. However, these data illuminate the importance of reducing LOS and conserving both time and resources in surgery without compromising treatment as they relate to economic savings for CSF leak. Financial considerations must be balanced with optimal clinical outcomes.

Conflict of Interest

None.

References

- 1 Dehdashti AR, Ganna A, Karabatsou K, Gentili F. Pure endoscopic endonasal approach for pituitary adenomas: early surgical results in 200 patients and comparison with previous microsurgical series. *Neurosurgery* 2008;62(05):1006–1015, discussion 1015–1017
- 2 Gondim JA, Schops M, de Almeida JP, et al. Endoscopic endonasal transsphenoidal surgery: surgical results of 228 pituitary adenomas treated in a pituitary center. *Pituitary* 2010;13(01):68–77
- 3 Zada G, Kelly DF, Cohan P, Wang C, Swerdloff R. Endonasal transsphenoidal approach for pituitary adenomas and other sellar lesions: an assessment of efficacy, safety, and patient impressions. *J Neurosurg* 2003;98(02):350–358
- 4 Amano K, Hori T, Kawamata T, Okada Y. Repair and prevention of cerebrospinal fluid leakage in transsphenoidal surgery: a sphenoid sinus mucosa technique. *Neurosurg Rev* 2016;39(01):123–131, discussion 131
- 5 Berker M, Hazer DB, Yücel T, et al. Complications of endoscopic surgery of the pituitary adenomas: analysis of 570 patients and review of the literature. *Pituitary* 2012;15(03):288–300
- 6 Cappabianca P, Cavallo LM, Colao A, de Divitiis E. Surgical complications associated with the endoscopic endonasal transsphenoidal approach for pituitary adenomas. *J Neurosurg* 2002;97(02):293–298
- 7 Piek J, Weber C, Kundt G, et al. Pharmacoeconomical consequences of postoperative CSF leaks after intracranial surgery - a prospective analysis. *J Neurol Surg A Cent Eur Neurosurg* 2012;73(01):25–28
- 8 Grotenhuis JA. Costs of postoperative cerebrospinal fluid leakage: 1-year, retrospective analysis of 412 consecutive nontrauma cases. *Surg Neurol* 2005;64(06):490–493, discussion 493–494
- 9 Lobatto DJ, de Vries F, Zamanipoor Najafabadi AH, et al. Preoperative risk factors for postoperative complications in endoscopic pituitary surgery: a systematic review. *Pituitary* 2018;21(01):84–97
- 10 Nishioka H, Haraoka J, Ikeda Y. Risk factors of cerebrospinal fluid rhinorrhea following transsphenoidal surgery. *Acta Neurochir (Wien)* 2005;147(11):1163–1166, discussion 1166
- 11 Dlouhy BJ, Madhavan K, Clinger JD, et al. Elevated body mass index and risk of postoperative CSF leak following transsphenoidal surgery. *J Neurosurg* 2012;116(06):1311–1317
- 12 Ivan ME, Iorgulescu JB, El-Sayed I, et al. Risk factors for postoperative cerebrospinal fluid leak and meningitis after expanded endoscopic endonasal surgery. *J Clin Neurosci* 2015;22(01):48–54
- 13 Boling CC, Karnezis TT, Baker AB, et al. Multi-institutional study of risk factors for perioperative morbidity following transnasal endoscopic pituitary adenoma surgery. *Int Forum Allergy Rhinol* 2016;6(01):101–107
- 14 Karnezis TT, Baker AB, Soler ZM, et al. Factors impacting cerebrospinal fluid leak rates in endoscopic sellar surgery. *Int Forum Allergy Rhinol* 2016;6(11):1117–1125
- 15 Senior BA, Ebert CS, Bednarski KK, et al. Minimally invasive pituitary surgery. *Laryngoscope* 2008;118(10):1842–1855
- 16 Jakimovski D, Bonci G, Attia M, et al. Incidence and significance of intraoperative cerebrospinal fluid leak in endoscopic pituitary surgery using intrathecal fluorescein. *World Neurosurg* 2014;82(3-4):e513–e523
- 17 McLaughlin N, Martin NA, Upadhyaya P, et al. Assessing the cost of contemporary pituitary care. *Neurosurg Focus* 2014;37(05):E7
- 18 Hanauer DA, Mei Q, Law J, Khanna R, Zheng K. Supporting information retrieval from electronic health records: a report of University of Michigan's nine-year experience in developing and using the Electronic Medical Record Search Engine (EMERSE). *J Biomed Inform* 2015;55:290–300
- 19 Ciric I, Ragin A, Baumgartner C, Pierce D. Complications of transsphenoidal surgery: results of a national survey, review of the literature, and personal experience. *Neurosurgery* 1997;40(02):225–236, discussion 236–237
- 20 Sun I, Lim JX, Goh CP, et al. Body mass index and the risk of postoperative cerebrospinal fluid leak following transsphenoidal surgery in an Asian population. *Singapore Med J* 2018;59(05):257–263
- 21 Lundeen EA, Park S, Pan L, O'Toole T, Matthews K, Blanck HM. Obesity prevalence among adults living in metropolitan and nonmetropolitan counties - United States, 2016. *MMWR Morb Mortal Wkly Rep* 2018;67(23):653–658
- 22 Lam G, Mehta V, Zada G. Spontaneous and medically induced cerebrospinal fluid leakage in the setting of pituitary adenomas: review of the literature. *Neurosurg Focus* 2012;32(06):E2
- 23 Horowitz G, Fliss DM, Margalit N, Wasserzug O, Gil Z. Association between cerebrospinal fluid leak and meningitis after skull base surgery. *Otolaryngol Head Neck Surg* 2011;145(04):689–693
- 24 Chern A, Hunter JB, Bennett ML. Cost analysis of cerebrospinal fluid leaks and cerebrospinal fluid leak prevention in patients undergoing cerebellopontine angle surgery. *Otol Neurotol* 2017;38(01):147–151