


Nickel(0)-Catalyzed [3+2] Cycloadditions of Bis(alkylidenecyclopropanes) with Diazenes: A Facile Synthesis of Functionalized Pyrazolidine-1,2-dicarboxylates

Bilash Kuila^aRayees Naikoo^aDinesh Mahajan^bPrabhpreet Singh^cGaurav Bhargava^{*a}

^a Department of Chemical Sciences, I. K. Gujral Punjab Technical University, Kapurthala, Punjab-144603, India
gaurav@ptu.ac.in
gauravorganic@gmail.com

^b Drug Discovery Research Centre (DDRC), Translational Health Sciences and Technology Institute (THSTI), Faridabad-121001, India

^c Department of Chemistry, Guru Nanak Dev University, Amritsar, Punjab 143005, India



Received: 08.09.2019

Accepted after revision: 07.11.2019

Published online: 26.11.2019

DOI: 10.1055/s-0039-1691502; Art ID: st-2019-d0474-1

Abstract A nickel(0)-catalyzed intermolecular [3+2] cycloaddition of bis(alkylidenecyclopropanes) with diazenes such as diethyl or diisopropyl azodicarboxylate gave pyrazolidine-1,2-dicarboxylates in moderate to good yields (61–72%).

Keywords bisalkylidenecyclopropanes, diazenes, pyrazolidine-dicarboxylates, [3+2] cycloaddition, nickel catalysis


Metal-catalyzed [*m+n*] cycloaddition reactions are effective tools for the synthesis of carbo- and heterocyclic systems.¹ Functionalized cyclopropanes have been exploited in metal-catalyzed [*m+n*] cycloadditions for the synthesis of monocyclic or condensed carbocycles.^{2–4} There are also reports on cycloadditions of activated cyclopropanes with heterodienophiles for the synthesis of monocyclic or condensed carbo- and heterocycles.^{2–4} De Meijere and co-workers explored the Lewis acid-catalyzed [3+2] cycloadditions of 2-arylcyclopropane-1,1-dicarboxylates with diazenes to afford functionalized pyrazolidines.⁵ However, [*m+n*] cycloadditions of functionalized nonactivated cyclopropanes, especially with heterodienophiles, have rarely been reported in the literature.^{2c,d,6,7}

Pyrazolidines have been evaluated as antibacterial, antifungal, anticancer, antidepressant, antiinflammatory, anti-tuberculosis, antioxidant, and antiviral agents in various pharmacological studies.⁸ Several pyrazolidine-based drugs have been marketed, including the antiinflammatory drug celecoxib, rimonabant for the treatment of obesity, fomepizole as an effective alcohol dehydrogenase inhibitor, and sildenafil as a phosphodiesterase inhibitor.⁹ Pyrazolidines are also useful as chiral auxiliaries and as synthetic re-

agents in multicomponent reactions.¹⁰ In addition, natural products containing pyrazolidine moieties have been shown to have pharmacological properties.¹¹

Conventional approaches, such as the condensation of 1,3-dicarbonyl compounds with hydrazines or [3+2] cycloadditions of 1,3-dipoles have been used in syntheses of simple pyrazolines.¹² However, there are few reports on synthesis of functionalized pyrazolidines. Chaudhry et al. recently reported acid-catalyzed cyclizations using allylic hydrazines for the synthesis of pyrazolidines.¹³

In view of these results and our ongoing interest in the cycloaddition chemistry of functionalized cyclopropanes, we wish to report an extension of our nickel(0)-catalyzed [3+2]-cycloadditions of bis(alkylidenecyclopropanes) to the preparation of pyrazolidines by using diazenes (Scheme 1).⁷



Scheme 1 [3+2]-Cycloadditions of bis(alkylidenecyclopropanes)

In the present work, we examined the [3+2] cycloadditions of bis(alkylidenecyclopropanes) with diazenes such as diethyl azodicarboxylate (DEAD) or diisopropyl azodicarboxylate (DIAD) as dienophiles (Scheme 1). The reaction


resulted in the formation of functionalized cyclopropane-tethered pyrazolidine-1,2-dicarboxylates in good yields.

The bis(alkylidene)cyclopropane reactants **1a–c** were synthesized by treating the appropriate dialkyl malonates with 1-vinylcyclopropyl tosylate by using the reported procedure.¹⁴ We examined the cycloaddition reactions of the bis(alkylidene)cyclopropanes **1a–c** with azodienophiles in the presence of nickel(0) complexes as catalysts. Importantly, the cycloaddition reactions with the diazenes DIAD and DEAD resulted in the formation of [3+2] cycloadducts, and no competitive [3+2+2] or [3+3+2] cycloadditions were observed.¹⁵ The [3+2] cycloadditions occurred by preferential ring opening at the allylic position of one of the methylenecyclopropane groups of the 2,2-bis(2-cyclopropylideneethyl)malonates **1a–c** to afford the corresponding pyrazolidinene-1,2-dicarboxylates **3a–f** in good yields.

Treatment of dimethyl bis(2-cyclopropylideneethyl)malonate (**1a**) with DEAD (**2a**) in toluene at various temperatures did not result in cycloaddition, and the starting material was recovered (Table 1, entries 1 and 2). The [3+2] cycloaddition of **1a** with DEAD (**2a**) was then examined in the presence of Ni(COD)₂ as catalyst under various conditions (entries 3–13). The reaction proceeded smoothly leading to the formation of diethyl (3*E*)-3-(5-cyclopropylidene-3,3-bis(methoxycarbonyl)pentylidene)pyrazolidine-1,2-dicarboxylate (**3a**) in good yields (entries 4–10). However, the [3+2] cycloaddition did not occur in the presence of the ligands PPh₃, DPPE, or P(OEt)₃ when Ni(COD)₂ was used as the catalyst (entries 11–13). The use of fewer equivalents of DEAD led to lower yields due to nonproductive decomposition of DEAD (entries 4 and 5). Poor yields

were observed in the polar aprotic solvent DMF, as well as in dichloroethane (DCE) or 1,4-dioxane (entries 8–10). Optimal conversion was obtained by using 10 equivalents of DEAD¹⁶ at 90 °C with toluene as solvent (entry 6).

After optimization of the reaction conditions, the [3+2]-cycloadditions of various dialkyl 2,2-bis(2-cyclopropylideneethyl)malonates with diazodienophiles using Ni(COD)₂ were explored (Scheme 2). These reactions resulted in the formation of functionalized pyrazolidine-1,2-dicarboxylates **3a–f** in moderate to good yields (Table 2).¹⁷ No significant change in the yield of the reaction was observed on changing the substrate or the dienophile.



The resulting [3+2] products were characterized by spectroscopic analysis.¹⁸ Diethyl (3*E*)-3-[5-cyclopropylidene-3,3-bis(isopropoxycarbonyl)pentylidene]pyrazolidine-1,2-dicarboxylate (**3c**), for example, showed an [M + H]⁺ ion at *m/z* 495.3 in its mass spectrum. The ¹H NMR (300 MHz) spectrum showed two characteristic multiplets at 5.61 and 5.57 ppm, corresponding to protons H₂ and H₃ re-

Table 1 Optimization of the Reaction Conditions for the [3+2] Cycloaddition of Dimethyl 2,2-Bis(2-cyclopropylideneethyl)malonate (**1a**) with Diethyl Azodicarboxylate (**2a**)

Entry	Catalyst	Ligand (1 equiv)	DEAD (equiv)	Solvent	Temp (°C)	Time (h)	Yield ^a (%)
1	–	–	10	toluene	40	24	0 ^b
2	–	–	10	toluene	120	24	0 ^b
3	Ni(COD) ₂	–	10	toluene	40	24 ^c	10
4	Ni(COD) ₂	–	1	toluene	90	16	15
5	Ni(COD) ₂	–	5	toluene	90	16	34
6	Ni(COD) ₂	–	10	toluene	90	16	72
7	Ni(COD) ₂	–	10	toluene	120	16	69
8	Ni(COD) ₂	–	10	1,4-dioxane	90	20	48
9	Ni(COD) ₂	–	10	DMF	110	24	22
10	Ni(COD) ₂	–	10	DCE	70	24	24
11	Ni(COD) ₂	PPh ₃	10	toluene	120	24	0
12	Ni(COD) ₂	DPPE	10	toluene	120	24	0
13	Ni(COD) ₂	P(OEt) ₃	10	toluene	120	24	0

^a Isolated yield after purification.

^b The starting material was recovered.


^c Incomplete reaction.

Table 2 Nickel-Catalyzed [3+2]-Cycloaddition Reactions of Bis-(alkylidenecyclopropanes) **1a–c** with Dialkyl Azodicarboxylates^a


Entry	R ¹	R ²	Product	Yield ^b (%)
1	Me	Et	3a	72
2	Et	Et	3b	70
3	<i>i</i> -Pr	Et	3c	61
4	Me	<i>i</i> -Pr	3d	68
5	Et	<i>i</i> -Pr	3e	67
6	<i>i</i> -Pr	<i>i</i> -Pr	3f	68

^a Reaction in toluene at 90 °C for 16 h.^b Isolated yield after purification.


spectively (Figure 1). A characteristic multiplet at 3.70 ppm corresponded to H₁ and two doublets of doublets at 1.05 ppm ($J = 4.5$ Hz) and 1.02 ppm ($J = 4.5$ Hz) were assigned to the H₄ and H₅ protons of the cyclopropyl ring, respectively. The ¹³C NMR spectrum showed the presence of two carbonyl carbons at 170.6 and 156.4 ppm corresponding to the isopropyl ester carbonyl and carbamate ester carbonyl, respectively. The ¹³C NMR spectrum also showed the presence of two olefinic carbons at 115.7 and 111.9 ppm, corresponding to C2 and C3, respectively, and two aliphatic carbons at 2.8 and 2 ppm, corresponding to C4 and C5, respectively (Figure 1).

**Figure 1** Diethyl (3E)-3-[5-cyclopropylidene-3,3-bis(isopropoxycarbonyl)pentylidene]pyrazolidine-1,2-dicarboxylate (**3c**)

A plausible mechanism for the metal-catalyzed cycloaddition involves an initial oxidative addition of the metal complex to the proximal bond of one of the methylene cyclopropanes of **1** to afford metallacyclobutane **4**. This is followed by oxidative insertion of the dialkyl azodicarboxylate to afford metallacycle **5**. The intermediates **4** and **5** are stabilized by coordination of the π -electrons of neighboring alkylidenecyclopropane moiety with the metal in a metallacyclobutane. The coordination of the metal in the metallacyclobutanes **4** and **5** with the neighboring cyclopropane alkene bond is deemed critical for the formation of functionalized pyrazolidine product **3**. Finally, intermediate **5**, upon reductive elimination, furnishes the [3+2]-cycloadduct product **3** (Scheme 3).

**Scheme 3** Plausible mechanism for the formation of **3**

Evidence that the presence of the second alkylidenecyclopropane is crucial for the success of these [3+2] cycloadditions came from the observation that the monoalkylidene compound **6** did not react in this manner, and no bisadducts were obtained (Scheme 4).

**Scheme 4**

In conclusion, we have developed an intermolecular [3+2] cycloaddition of previously unexplored bis-(alkylidenecyclopropanes) with diazenes mediated by a nickel(0) catalyst. The diazenes DIAD and DEAD were used in these intermolecular [3+2]-cycloaddition reactions, resulting in the formation of functionalized pyrazolidine-1,2-dicarboxylates in moderate to good yields. Further work exploring the application of transition metals in [m+n] cycloadditions of bis(alkylidenecyclopropanes) is in progress.

Funding Information

The Board of Research in Nuclear Sciences (BRNS), India, is thanked for Research Grant, Project No.2013/37C/11/BRNS/198. The Department of Science and Technology (DST), India, is also thanked for Research Grant, Project No. SB/FT/CS-079/2012.

Supporting Information

Supporting information for this article is available online at <https://doi.org/10.1055/s-0039-1691502>.

References and Notes

- (1) (a) Inglesby, P. A.; Evans, P. A. *Chem. Soc. Rev.* **2010**, *39*, 2791.
(b) Balme, G.; Bouyssi, G.; Monteiro, N. In *Multicomponent Reactions*; Zhu, J.; Bienaymé, H., Ed.; Wiley-VCH: Weinheim, **2005**,

- Chap. 8, 224. (c) Wang, Y.; Wang, J.; Su, J.; Huang, F.; Jiao, L.; Liang, Y.; Yang, D.; Zhang, S.; Wender, P. A.; Yu, Z.-X. *J. Am. Chem. Soc.* **2007**, *129*, 10060.
- (2) (a) Kulinkovich, O. G. *Cyclopropanes in Organic Synthesis*; Wiley: Hoboken, **2015**; and references cited therein. (b) Jiabin, L.; Liu, R.; Wei, Y.; Shi, M. *Trends Chem.* **2019**, *1*, 779. (c) Binger, P.; Büch, H. M. *Top. Curr. Chem.* **1987**, *135*, 77. (d) Trost, B. M. *Angew. Chem. Int. Ed.* **1986**, *25*, 1.
- (3) (a) Carson, C. A.; Kerr, M. A. *Chem. Soc. Rev.* **2009**, *38*, 3051. (b) Reissig, H.-U.; Zimmer, R. *Chem. Rev.* **2003**, *103*, 1151. (c) Schneider, T. F.; Kaschel, J.; Werz, D. B. *Angew. Chem. Int. Ed.* **2014**, *53*, 5504.
- (4) (a) Shao, L.-X.; Shi, M. *Curr. Org. Chem.* **2007**, *11*, 1135. (b) Waser, J.; De Simone, F. *Synthesis* **2009**, 3353. (c) Lautens, M.; Ren, Y. *J. Am. Chem. Soc.* **1996**, *118*, 9597.
- (5) (a) Korotkov, V. S.; Larionov, O. V.; Hofmeister, A.; Magull, J.; Meijere, A. D. *J. Org. Chem.* **2007**, *72*, 7504. (b) Zhao, L.; De Meijere, A. *Adv. Synth. Catal.* **2006**, *384*, 2484.
- (6) For metal-catalyzed $[m+n]$ -cycloaddition reactions, see: (a) Noyori, R.; Odagi, T.; Takaya, H. *J. Am. Chem. Soc.* **1970**, *92*, 5780. (b) Noyori, R.; Kumagai, Y.; Umeda, I.; Takaya, H. *J. Am. Chem. Soc.* **1972**, *94*, 4018. (c) Binger, P.; Doyle, M. J.; Benn, R. *Chem. Ber.* **1983**, *116*, 1. (d) Binger, P.; Brinkmann, A.; Wedemann, P. *Chem. Ber.* **1983**, *116*, 2920. (e) Binger, P.; Freund, A.; Wedemann, P. *Tetrahedron* **1989**, *45*, 2887. (f) Trost, B. M.; MacPherson, D. T. *J. Am. Chem. Soc.* **1987**, *109*, 3483. (g) Trillo, B.; López, F.; Gullias, M.; Castedo, L.; Mascareñas, J. L. *Angew. Chem. Int. Ed.* **2008**, *47*, 951. (h) Trost, B. M.; Hu, Y.; Horne, D. B. *J. Am. Chem. Soc.* **2007**, *129*, 11781.
- (7) (a) Kuila, B.; Mahajan, D.; Singh, P.; Bhargava, G. *Tetrahedron Lett.* **2015**, *56*, 1307. (b) Kuila, B.; Mahajan, D.; Singh, P.; Bhargava, G. *Eur. J. Org. Chem.* **2018**, 853.
- (8) (a) Ansari, A.; Ali, A.; Asif, M.; Shamsuzzaman, *New J. Chem.* **2017**, *41*, 16. (b) Karrouchi, K.; Radi, S.; Ramli, Y.; Taoufik, J.; Mabkhot, Y. N.; Al-aizari, F. A.; Ansar, M. *Molecules* **2018**, *23*, 134; and reference cited therein.
- (9) Mert, S.; Kasimogullari, R.; Ica, T.; Colak, F.; Altun, A.; Ok, S. *Eur. J. Med. Chem.* **2014**, *78*, 86.
- (10) (a) Tu, X.-J.; Hao, W.-J.; Ye, Q.; Wang, S.-S.; Jiang, B.; Li, G.; Tu, S.-J. *J. Org. Chem.* **2014**, *79*, 11110. (b) Castagnolo, D.; Schenone, S.; Botta, M. *Chem. Rev.* **2011**, *111*, 5247.
- (11) Kumar, V.; Kaur, K.; Gupta, G. K.; Sharma, A. K. *Eur. J. Med. Chem.* **2013**, *69*, 735.
- (12) (a) Elguero, J. In *Comprehensive Heterocyclic Chemistry*, Vol. 5; Katritzky, A. R.; Rees, C. W.; Potts, K. T., Ed.; Pergamon: Oxford, **1984**, Chap. 4.04, 167. (b) Elguero, J. *Comprehensive Heterocyclic Chemistry II*, Vol. 3; Katritzky, A. R.; Rees, C. W.; Scriven, E. F. V., Ed.; Pergamon: Oxford, **1996**, Chap. 3.01, 1. (c) Katritzky, A. R.; Wang, M.; Zhang, S.; Voronkov, M. V.; Steel, P. J. *J. Org. Chem.* **2001**, *66*, 6787. (d) Huang, Y. R. *Katzenellenbogen J. A. Org. Lett.* **2000**, *2*, 2833. (e) Wang, C.; Chen, Y.; Li, J.; Zhou, L.; Wang, B.; Xiao, Y.; Guo, H. *Org. Lett.* **2019**, *21*, 7519. (f) Liu, X.; Zhou, Y.; Song, Q. *Chem. Commun.* **2019**, *55*, 8943; and references cited therein.
- (13) (a) Yamazaki, S.; Maenaka, Y.; Fujinami, K.; Mikata, Y. *RSC Adv.* **2012**, *2*, 8095. (b) Chaudhry, F.; Kariuki, B. M.; Knight, D. W. *Tetrahedron Lett.* **2016**, *57*, 2833.
- (14) Stolle, A.; Ollivier, J.; Piras, P. P.; Salaiin, S.; De Meijere, A. *J. Am. Chem. Soc.* **1992**, *114*, 4051.
- (15) (a) Saya, L.; Bhargava, G.; Navarro, M. A.; Gullias, M.; López, F.; Fernández, I.; Castedo, L.; Mascareñas, J. L. *Angew. Chem. Int. Ed.* **2010**, *49*, 9886. (b) Saya, L.; Fernández, I.; López, F.; Mascareñas, J. L. *Org. Lett.* **2014**, *16*, 5008.
- (16) An excess of the azodienophile is used to compensate for its thermal decomposition during the reaction.
- (17) **[3+2]-Cycloadducts 3a–f; General Procedure**
Dialkyl azodicarboxylate **2** (2 mmol, 10 equiv) was added to a solution of the appropriate bis(alkylidene)cyclopropane **1** (0.2 mmol, 1 equiv) in toluene (4 mL), and the mixture was degassed for 10 min under argon. Ni(COD)₂ (5 mol%) was added, and the mixture was heated to 90 °C for 16 h. After completion of the reaction, the mixture was cooled to r.t., directly loaded onto a column without evaporation, and purified by flash chromatography [silica gel (100–200 mesh), EtOAc–hexanes].
- (18) **Diethyl (3E)-3-[5-Cyclopropylidene-3,3-bis(methoxycarbonyl)pentylidene]pyrazolidine-1,2-dicarboxylate (3a)**
Colorless liquid; yield: 119 mg (72%). ¹H NMR (300 MHz, CDCl₃): δ = 5.71 (m, 1 H), 5.48 (m, 1 H), 4.13–4.20 (m, 4 H), 3.71 (s, 6 H), 3.65 (m, 2 H), 2.75 (d, J = 9 Hz, 2 H), 2.49 (br s, 2 H), 2.01 (m, 2 H), 1.22–1.27 (m, 6 H), 0.99 (m, 4 H). ¹³C NMR (75 MHz, CDCl₃): δ = 171.1, 156.3, 137.7, 126.9, 116.0, 112.3, 61.5, 57.7, 52.1, 46.7, 37.0, 35.5, 29.3, 14.6, 2.6, 2.0. LRMS (ESI): m/z = 439.2 [M + H]⁺. HRMS (ESI): m/z [M + H]⁺ calcd for C₂₁H₃₁N₂O₈: 439.2080; found: 439.2067.
- Diethyl (3E)-3-[5-Cyclopropylidene-3,3-bis(isopropoxycarbonyl)pentylidene]pyrazolidine-1,2-dicarboxylate (3c)**
Pale-yellow liquid; yield: 94 mg (61%). ¹H NMR (300 MHz, CDCl₃): δ = 5.61 (m, 1 H), 5.57 (m, 1 H), 5.03 (m, 2 H), 4.19 (m, 4 H), 3.70 (m, 2 H), 2.76 (d, J = 7.2 Hz, 2 H), 2.66 (bs, 2 H), 1.78 (m, 2 H), 1.20–1.32 (m, 18 H), 1.05 (t, J = 4.5 Hz, 2 H), 1.02 (t, J = 4.5 Hz, 2 H). ¹³C NMR (75 MHz, CDCl₃): δ = 170.4, 155.6, 137.7, 126.2, 115.5, 112.1, 69.0, 62.4, 57.0, 46.1, 37.8, 35.8, 29.6, 21.5, 14.5, 2.8, 2.0. LRMS (ESI): m/z = 495.3 [M + H]⁺. HRMS (ESI): m/z [M + H]⁺ calcd for C₂₅H₃₉N₂O₈: 495.2706; found: 495.2718.