



The whistle-smile reflex: a missed sign

O reflexo do assobio-sorriso: um sinal esquecido

Mateus Damiani Monteiro¹ Giovanna Testa Brustolin¹ Eurípedes Gomes de Carvalho Neto¹
Carlos Roberto de Melo Rieder¹

¹Santa Casa de Misericórdia de Porto Alegre, Departamento de Neurologia, Porto Alegre RS, Brazil.

Address for correspondence Mateus Damiani Monteiro (e-mail: mateusdamiani@gmail.com).

Arq. Neuropsiquiatr. 2022;80(10):1075–1076.

Abstract

In this paper, we present a historical review of the whistle-smile reflex, a semiological sign missed in the literature and clinical practice.

Keywords

- Parkinson Disease
- Physical Examination

Resumo

Apresentamos neste artigo uma revisão histórica do reflexo do assobio-sorriso, um sinal semiológico esquecido na literatura e na prática clínica.

Palavras-chave

- Doença de Parkinson
- Exame Físico

INTRODUCTION

In 1943, Frederic Hanes described the whistle-smile reflex, also known as Hanes sign: when asked to whistle, a healthy person does so and then smiles after the unusual physician's request. Otherwise, a patient with parkinsonism performs the whistle but does not smile after whistling due to the facial bradykinetic movement¹ (► **Video 1** and ► **Figure 1**).

Dr. Hanes was a neurologist and professor of medicine at Duke University. During his many years in Durham, North Carolina, he explored the presence of hypomimia as a semiological tool for the diagnosis of Parkinson disease (PD); finding, during his evaluations, what he considers a helpful and reliable sign.

Lack of facial expression is a well-known feature of parkinsonian syndrome and an essential clue to diagnosing PD. Although James Parkinson did not mention hypomimia in his seminal work "An Essay on the Shaking Palsy"², it was recognized as an important semiological finding in a patient with PD in early works by authors such as Charcot, Gowers, and Wilson.³ The facial movement impairment may be not only voluntary but also spontaneous and emotional. The involvement of emotional expressions can be noticed with spontaneous smiling, reflected in reduced frequency and opening degree of the mouth.⁴

The whistle-smile reflex, alongside the Rolex sign⁵ and the floating door sign,⁶ lost its role in assessing a parkinsonian patient throughout the years. Although it is an interesting possibility to use the whistle-smile reflex in clinical

Video 1

The whistle-smile reflex. The patient and the healthy person gave us written consent to use their images. <https://www.arquivosdeneuropsiquiatria.org/wp-content/uploads/2022/08/ANP-2022.0120-Video.mov> Online content including video sequences viewable at: <https://www.thieme-connect.com/products/ejournals/html/10.1055/s-0042-1758396>.



received
June 3, 2022
accepted
July 24, 2022

DOI <https://doi.org/10.1055/s-0042-1758396>.
ISSN 0004-282X.

© 2022. Academia Brasileira de Neurologia. All rights reserved. This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>). Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil



Figure 1 Non-Parkinson disease patient's smiles after whistling (A-B). Parkinson disease patient shows no reaction after doing the same (C-D).

practice, this reflex has not been revised in detail over the years, and we still do not have established its sensitivity and specificity. Therefore, it deserves to be remembered, not only for its historical importance but for its possible semiological value in clinical practice, and further studies are needed for it

to become a routine tool in the assessment of a parkinsonian patient.

Authors' Contributions

MDM: writing, acquisition of data and literature review; GTB: image acquisition; EGCN: image editing, revision of manuscript; CRMR: critical revision of manuscript for intellectual content and study supervision.

Support

Carlos R. M. Rieder receives support from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) – Brazil (scholarship of research productivity). The other authors declare that there are no additional disclosures to report.

Conflict of Interest

The authors have no conflict of interests to declare.

References

- Hanes FM. Two Clinically Useful Signs: 1. The Whistle-Smile Reflex in the Parkinsonian Syndrome 2. The 'Nephritic Stare' in Chronic Nephritis. *JAMA* 1943;121(14):1152–1153
- Parkinson J. An essay on the shaking palsy. 1817. *J Neuropsychiatry Clin Neurosci* 2002;14(02):223–236, discussion 222. Doi: 10.1176/jnp.14.2.223
- Maycas-Cepeda T, López-Ruiz P, Feliz-Feliz C, et al. Hypomimia in Parkinson's Disease: What Is It Telling Us? *Front Neurol* 2021; 11:603582
- Bologna M, Fabbri G, Marsili L, Defazio G, Thompson PD, Berardelli A. Facial bradykinesia. *J Neurol Neurosurg Psychiatry* 2013;84(06):681–685
- Teive HA, Sá DS. The Rolex sign: first manifestation of Parkinson's disease: case report. *Arq Neuropsiquiatr* 2000;58(3A):724–725
- Kulkarni O, Lafaver K, Tarsy D. The "floating door sign" in Parkinson's disease. *Parkinsonism Relat Disord* 2013;19(09):825–826