

Address for correspondence Fernanda Rocha Rojas Ayala, MD, PhD,

Department of Medical Genetics, Unicamp University, Oncogenetics

Service at Clinics Hospital, Campinas, SP 13083-888, Brazil

(e-mail: fernotipogenotipo@gmail.com).

### THIEME OPEN ACCESS

# Replication Timing Aberration of *KIF14* and *MDM4*/*PI3KC*2 $\beta$ Alleles and Aneuploidy as Markers of Chromosomal Instability and Poor Treatment Response in Ewing Family Tumor Patients

Fernanda Rocha Rojas Ayala<sup>1</sup> Jeffrey William Martin<sup>2</sup> Carmen Silvia Bertuzzo<sup>3</sup>

Glob Med Genet 2023;10:54-62.

# Abstract

Replication timing of allelic gene pairs is strictly regulated according to expression, genome stability, and epigenetic changes, and tumorigenesis may be associated with changes in the allelic replication in various tumors. Our aim was to determine whether such alterations had a prognostic value in Ewing's family tumor (EFT) patients. The *KIF14* and *MDM4*/*PI3KC*2β and the centromeric satellite sequence of chromosomes 8 and 12 were used for replication timing assessments. Aneuploidy was assessed by enumerating the copy numbers of chromosomes 8 and 12. Replication timing and aneuploidy were detected cytogenetically using multicolors fluorescence in situ hybridization assay applied in 135 EFT. Patients with trisomy 8 presented an association with an asynchronous replication pattern (SD) of  $MDM4/PI3KC2\beta$  genes (p = 0.013). Trisomy 12 was associated with a synchronous pattern (DD) of KIF14 probe signals (p = 0.04). The DD synchronous replication pattern of KIF14 showed a correlation with age (p < 0.0001), and the SS synchronous replication pattern of the same locus showed a correlation with lung metastatic (p = 0.012). The subgroup of patients presenting with multiplet signals of MDM4/PI3KC2β showed an association with treatment response (p = 0.045) and age (p = 0.033). Replication pattern of KIF14 may, significantly, be associated with chromosomal instability as  $MDM4/PI3KC2\beta$  may be a considerably new marker of poor treatment response in EFT patients.

### Keywords

- replication timing
- ► aneuploidy
- chromosomal instability
- poor treatment response
- Ewing sarcoma

### Introduction

The temporal order of allelic replication, whereby both alleles of a gene expressed in a biallelic mode replicate synchronously or where the alleles of a gene undergoing monoallelic expression replicate asynchronously (allele-spe-

> DOI https://doi.org/ 10.1055/s-0043-1768238. ISSN 2699-9404.

cific replication), has the expressed allele replicating earlier than the silenced allele.<sup>1</sup> Imprinting, X-inactivation, and allelic exclusion may be the cause of the silencing of a single allele<sup>2</sup> where an unscheduled alteration in replication timing may delay the process of chromosome condensation, increasing genetic instability in several tumor cell lines and

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

<sup>&</sup>lt;sup>1</sup> Department of Medical Genetics, Unicamp University, Oncogenetics Service at Clinics Hospital, Campinas, SP, Brazil

<sup>&</sup>lt;sup>2</sup> Department of Medicine, Toronto University, Toronto, Ontario, Canada

<sup>&</sup>lt;sup>3</sup>Department of Genetics and Genomics, at Unicamp University, Oncogenetics Services at Clinics Hospital, Campinas, SP, Brazil

<sup>© 2023.</sup> The Author(s).

primary tumor samples. While the timing of replication has been investigating as an epigenetic marker demonstrating whether a particular gene is programmed to express biallelically or monoallelically,<sup>3,4</sup> others have established the asynchronous allelic replication of various human monoallelically expressed genes, such as those subjected to X inactivation, as well as numerous imprinted genes.<sup>5</sup> Previous published results in the peripheral blood lymphocytes of patients with hematological malignancies or solid tumors—such as renal cell carcinoma or prostate cancer and breast cancer<sup>6</sup> genes that replicate synchronously including tumor suppressor genes *TP53* and *RB1* and oncogenes *C-MYC* and *HER2* lose their synchronous mode and replicate asynchronously.

It is not yet completely understood how this sequential progression of replication in S-phase is regulated in human cells. Disruption of this process can be altered in cells with aneuploidy or a genetic predisposition to cancer<sup>2,7</sup> identifying as a possible marker for genetic instability related to cancer.<sup>8,9</sup> In this way, the allelic synchronization correlated with a higher frequency of aneuploidy is frequently lost in cells derived from patients with various cancers, such as leukemia,<sup>10</sup> in which aberrant chromosomes are known to bypass cell-cycle checkpoints and escape apoptosis<sup>7</sup>

Several studies have consistently shown trisomy of chromosomes 8 and 12 as the most common genetic abnormality in EFT patients.<sup>11-16</sup> Gains in chromosome number have been associated with more advanced disease in several solid tumors and are thought to reflect the loss of balanced control of mitotic disjunction.<sup>17</sup> Chromosomal aberrations acquired in addition to the EWS gene rearrangements have been postulated to serve as markers for more refractory disease. The 1q32.1 region investigated here encompasses two potential oncogenes, KIF14 and MDM4/PI3KC2B. KIF14 is an essential protein for cytokinesis and chromosome segregation,<sup>18</sup> and its overexpression is already reported to be associated with poor prognosis in lung cancer<sup>19</sup>, while increased MDM4 levels and the resulting inactivation of p53 contribute to the development of breast cancers.<sup>20</sup>

The interphase fluorescence in situ hybridization (iFISH) assay has been used to visualize specific genomic DNA sequences in interphase nuclei, making it possible to accurately examine the replication timing pattern of various loci within a single cell.<sup>2,21–23</sup> To the best of our knowledge, none has related the replication pattern of genes expressed in formalin-fixed paraffin-embedded (FFPE) tissue to clinical-pathological parameters of cancer patients so far.

In the current study, we estimated the replication patterns of *KIF14* and *MDM4/PI3KC*2 $\beta$  homologous loci in EFT patients and associated them with known significant aberrations such as polysomy 8 and 12 to estimate the influence of these specific aberrations on gene replication. We further examined the relationship between the clinical-pathological features of EFT patients and the behavior of the *KIF14* and *MDM4/PI3KC*2 $\beta$  loci.

# **Materials and Methods**

### **Patients and Tumors Specimens**

We retrospectively analyzed a total of 135 paraffin-embedded patient tumors retrieved from the pathological files between 1980 and 2008 at the Pathology Department of A. C. Camargo Hospital of Sao Paulo, Brazil. The diagnosis was given by the histopathology review, immunohistochemical staining for CD99 and FLI1 antibodies, and by iFISH for EWSR1/FLI1 fusion presence or not. In this series, confirmation of 22q12 rearrangements was present in 95.5% of the cases. Thus, the tissue microarray (TMA) slide included 135 cases in duplicate, of which 118 cases were primary tumors, while 17 cases were recurrent tumors. Retrospective clinical data gleaned from the corresponding patients' records included demographic characteristics, surgical resection data, and clinical outcome. Criteria for inclusion were initial treatment of the primary tumor in Hospital A.C. Camargo; clinical data available and FFPE tissue with material sufficient and with quality for the study; and at least 3 years of follow-up (except patients who died of the disease before 3 years). We excluded patients with a second synchronous tumor. Patients' treatment included multiagent chemotherapy, local control measures with radiotherapy and surgery, or a combination of these as appropriate for the period of the study. The most utilized drugs were ifosfamide, cyclofosfamide, adriamicine, etoposide, vincristine, and actinomicine. Radiotherapy was indicated for unresectable tumors, resectable, or partially resectable tumors in patients with metastatic disease and as palliative treatment.<sup>24</sup> Of 135 cases, 104 patients were treated with preoperative treatment and 29 patients were untreated; therapy status was unknown for two patients. The patient's clinical-pathological characteristics are depicted in **-Table 1**. This study has been reviewed and approved by the Ethical Review Board of A.C. Camargo Hospital.

### **Probe Selection**

The University of California Santa Cruz Human Genome Browser Database, March 2006 build (Greulich-Bode et al, 2008), was used to select the bacterial artificial chromosome (BAC) genomic name clones and their linear order (www.genome.ucsc.edu). The strategy for the assessment of copy number alterations (CNAs) in EFT by iFISH was designed using BAC clones (RP11-192G12, RP11-97B14/ RP11-433N15) covering approximately 4 Mb of the 1q32.1 cytoband (KIF14, MDM4/PI3KC2β) and BAC clones (RP11-54H15/RP11-196G18) for the 1q21 cytoband (pericentromeric region; **Fig. 1**). Subsequently, the BAC clones were obtained from the Centre for Applied Genomics, Sick Kids Hospital (Toronto, Ontario, Canada). Each test probe was paired with a reference probe in either the same chromosome arm or the chromosome centromere to control for numerical changes. Commercial probes (SpAqua -D8Z2 and SpOrange- D12Z3 - Abbott Molecular, IL, United States) were used for the interphase FISH enumeration of the whole chromosomes 8 and 12.

Categories	Patients N (%)
Male	82 (60.7)
Female	53 (39.3)
S15	74 (54.8)
>15	61 (45.2)
Osseous	121 (89.6)
Extraosseous	14 (10.4)
Inferior extremity	61 (45.2)
Pelvis	31 (23)
Superior extremity	28 (20.7)
Thorax	14 (10.4)
Others	1 (0.7)
Localized	97 (71.8)
Metastatic	38 (28.2)
Pain	30 (22.2)
Swelling	12 (8.9)
Pain + swelling	61 (45.2)
Others	24 (17 0.8)
Ignored	8 (5.9)
Treated	104 (77)
Untreated	29 (21.5)
Ignored	2(1.5)
Yes	69 (51.1)
No	65 (48.1)
Alive without disease	34 (25.2)
Alive with disease	17 (12.6)
Death due to disease	83 (61.5)
unknown	1 (0.7)
	Categories Male Male Male Female S15

 Table 1 Clinical-pathological characteristics of the study patients

**Probes Preparation and Fluorescent in situ Hybridization** BAC DNA (1 μg) was extracted by modification of the Qiagen mini-Kit protocol (Qiagen, Valencia, CA, United States) and labeled with SpectrumGreen-dUTP, SpectrumOrange-dUTP (Abbott Molecular), or SpectrumRed-dUTP (PerkinElmer Life and Analytical Sciences, Boston, MA, United States) using the Vysis Nick Translation Kit (Abbott Molecular). Labeling of probes was done as described previously.<sup>25,26</sup> The locus specificity and the correct chromosome location of all BAC clones were confirmed by polymerase chain reaction and metaphase FISH of normal peripheral lymphocytes, respectively.

FISH assays were performed using mixtures of the labeled BAC probes, according to previously published procedures,<sup>27</sup> with slight modifications, the incorporation of the micro-wave oven for pretreatment with citrate buffer and for crossed-link protein breakage to enhance the hybridization

efficiency as described elsewhere.<sup>28,29</sup> Briefly, TMA slides were deparaffinized, dehydrated, and air-dried. Slides were pretreated in sodium citrate (10 mM) using the microwave in three intervals of 5 minutes. Later, the slides were digested with 750 U/mL pepsin (Sigma-Aldrich) at 37°C for 8 minutes. The slides were then washed in  $2 \times$  saline-sodium citrate (SSC) and dehydrated in a graded ethanol series. Air-dried slides were warmed to 37°C inside the ThermoHybrite (Vysis-Abbott Laboratories), and then the probe mixture was applied to the target DNA area and a cover slip was sealed in place with rubber cement. Slides were placed in the microwave and heated at high power (920 W) for 70 seconds. Slides were codenatured at 80°C for 10 minutes and hybridized in a humidified chamber at 37°C for 16 to 20 hours (ThermoHybrite - Vysis-Abbott Laboratories). Following hybridization the slides were washed at 75°C in  $2 \times$  SCC/0.3% NP40 for 2 minutes, followed by 5 minutes in  $2 \times$  SCC at room temperature. Slides were mounted with 4',6-diamidino-2-phenylindole, dihydrochloride-Vectashield (Vector Laboratories, Inc. Burlingame, CA, United States) and, after 24 hours at -20°C they were examined blindly using an epifluorescence Zeiss Imager. M1 microscope was equipped with a digital camera AxioCam MRm and AxioVision 4.3 capturing software (Carl Zeiss Canada Ltd.).

### **Cytogenetic Analysis**

# KIF14 and MDM4/PI3KC2β Locus (1q32.1).

The replication patterns of genes in the 1q32.1 locus were assessed by the configuration of hybridization signals following three-color iFISH with a control probe for the pericentromeric region.<sup>30</sup> We analyzed the relative proportions of the replication pattern of cells in three groups: (1) all populations, (2) those with CNAs, and (3) those without CNAs. For each sample, at least 100 nonoverlapping, intact nuclei exhibiting distinct well-defined fluorescence hybridization signals were scored.

The examined cells were classified into four categories: (1) SS, for cells with two or more similarly shaped signals, or singlets, per locus; (2) DD, for cells with one or more doublets per locus; (3) SD, for cells with one singlet and one doublet signal per locus; and (4) multiplets, for cells with three or more signals spaced less than one signal diameter apart (Fig. 2). We recorded the frequency of SS, DD, SD, and multiplets in each cell population. Thus, based on the rate of the four replication configurations, it was possible to differentiate between three distinct replication patterns of homologous loci as previously described in Selig et al, (1992): (1) a synchronous replication pattern as revealed by two or more singlets (SS) or unreplicated loci in G1; (2) a synchronous replication pattern where cells have two or more doublets (DD) or replicated loci in S or G2; and (3) an asynchronous pattern with a relatively long time interval between early- and late-replicating loci, as indicated by two different signal configurations describing two nonsuccessive replication stages (SD), representing the S-phase where only one allele has been replicated. In cells that showed extragene copies (i.e., CNA), the scoring was done by counting the



**Fig. 1** FISH strategy analysis of *KIF14* or *MDM4*/*P/3KC28* genes replicating pattern in EFT cells at interphase. (A) The schematic chromosome 1illustrates the genomic localization and names of the BAC probes encompassing the pericentromeric region and the target genes at 1q32.1. (B) Representative FISH scheme are shown for EFT TMA applying the three-color FISH for the replication pattern analysis.



**Fig. 2** Schematic representation of timing replicating pattern of the *KIF14* or *MDM4*/*PI3KC2B* hybridization signals at interphase by FISH. (A) Configuration of signals in normal (no CNA) cells showing different replication patterns. (B) Guide for signal enumeration when the genes dots show up as doublets or multiplet. Note that in cells that showed gain of the genes, only the SS, DD, and M patterns were counted. Abbreviation: CNA\*: copy number alteration.

number of singlet (SS), doublet (DD), and multiplet (M) signals, probably representing an average of several different replication times within a limited range of S phase.<sup>21</sup> It has been demonstrated elsewhere that genes which replicate early in the cell cycle will show a high percentage of doublets, while for late-replicating genes, most nuclei will have singlet hybridization signals.<sup>21</sup> Therefore, it was assumed that the presence of a larger number of doublets represented an earlier time of replication.<sup>7,21</sup>

### **Detection of Aneusomy**

A dual-color iFISH assay was performed here to estimate the rate of aneusomy within cell populations. A conservative analysis of copy number enumeration was measured by counting the copies of chromosomes 8 and 12 based on the cut-off of 10% being level for the assay (mean  $\pm 2$  seconds in tumor diploid controls). Hence, we had two groups of genetic parameters – one normal (no CNA) and the other with CNA. For each cell, we recorded the frequencies of cells with one, two, and three or more signals.

### **Statistical Analysis**

The iFISH data for the gene replication patterns of the 1q32.1 locus were classified as continuous variables for all populations of EFT patients. The parameters included in the analysis were onset of symptoms to diagnosis, age ( $\leq$ 15 vs. >15 years), metastatic location at diagnosis (lung vs. nonlung), treatment response (Huvos grade), trisomy 8 and 12, and gains at 1q32.1. The nonparametric Mann–Whitney test was applied for testing differences between study groups for quantitative clinical parameters. The Spearman correlation was applied for testing the correlation between the different clinical parameters in the study group. A *p*-value of less than 0.05 was considered to indicate statistical significance. All tests were two sided. Statistical analysis was performed using Statistical Packages of Social Sciences (SPSS for Macintosh, Version 16.0, Inc, Chicago, IL, United States).

### Results

Multicolor iFISH was applied to EFT TMA comprising 135 patients by scoring for 100 cells in each tumor core. We examined whether there was a relationship between the replication behavior of genes and clinical-pathological characteristics as described below.

Different replication timing patterns were evaluated in 90/135 cases for *MDM4/PI3KC*2β and in 71/135 cases for *KIF14* in the total population. The proportion of cells with asynchronous (SD and M) replication and synchronous replication (SS and DD) patterns of *KIF14* alleles and of *MDM4/ PI3KC*2β alleles in the study groups are shown in – **Figs. 3**, **4**, and **5**. The most frequent replication pattern seen in all populations was the asynchronous stage (SD) either for *KIF14* or for *MDM4/PI3KC*2β, varying between 1.4 to 23.2% and 1.1 to 13.6%, respectively. For the group that showed gain at the 1q32.1 locus, there was an increase of SD, DD, and M patterns for *KIF14* alleles, whereas for *MDM4/PI3KC*2β signals, the SD and M replication patterns were increased when



**Fig. 3** Frequency of cells showing replication timing (SS, DD, SD, M) of *KIF14* and *MDM4/ P/3KC2B* alleles in the total EFT patients.

compared with the group that did not showed gain at that region. The higher frequency of the asynchronous pattern for MDM4/PI3KC2β indicates that, in EFT cells, its two loci differ largely in their replication timing. Additionally, in the group with trisomy 8, there was a slightly higher rate of the SS synchronous pattern for the KIF14 and MDM4/PI3KC2β alleles than in the group with trisomy 8. Furthermore, in the group with trisomy 12, there was a higher rate of DD synchronous pattern for the KIF14 alleles, whereas for the *MDM4*/**PI3KC**2β alleles, there was a slight increase of the SD pattern compared with the group without trisomy 12. Also, there were more cells with single dots in the studied groups, but without statistical significance. The difference in the proportion of cells with SD pattern in the groups studied (1q32.1, trisomy 8 and 12) did not show a statistically significant correlation for both probes analyzed. Interestingly, there was a significant negative correlation between SS and DD, SD, and M patterns for both hybridization signals for *KIF14* and *MDM4*/*PI3KC*2 $\beta$  (p < 0.05) in the total population.

### Statistical Analysis of Replication Pattern and Clinical-Pathological Features

The group of patients with trisomy chromosome 8 (27/124) presented a statistically significant association with cells that showed an asynchronous (SD) replication pattern of *MDM4/***PI3KC**2 $\beta$  genes (p = 0.013). Also, the subset of patients with trisomy chromosome 12 (18/124) showed a statistically significant association with a synchronous (DD) pattern of *KIF14* probe signals (p = 0.04) in all populations.

The DD synchronous replication pattern of *KIF14* showed a correlation with age (p < 0.0001), whereas the SS synchronous replication pattern of the same locus showed a correlation with the metastatic site (lung vs. nonlung; p = 0.012). Our results also showed a negative correlation between the DD synchronous replication of *KIF14* and the onset of symptoms (r = -0.260, p = 0.03). Additionally, the subgroup of patients presenting with the multiplets replication pattern of *MDM4*/*PI3KC*2 $\beta$  genes showed an association with clinical response (p = 0.045) in all populations and in the group with gain at 1q32.1 (15/128). Thus, patients who had poor treatment response (Huvos grade I, II<sup>31</sup>;) showed a lower mean



**Fig. 4** Distribution of cells showing replication timing (SS, DD, SD, M) of KIF14 alleles in EFT patients. The signals were evaluated in cells of EFT patients with CNA and with no CNA for chromosome gains at 1q32.1 and aneusomy of chromosomes 8 and 12.



**Fig. 5** Distribution of cells showing replication timing (SS, DD, SD, M) of *MDM4/P/3KC2B* alleles in EFT patients. The signals were evaluated in cells of EFT patients with CNA and with no CNA for chromosome gains at 1q32.1 and aneusomy of chromosomes 8 and 12.

(0.29 vs. 1.0) of multiplet signals per cell. This multiplet replication pattern of *MDM4*/**PI3KC**2 $\beta$  genes showed an association with age (p = 0.033) in all populations.

# Discussion

Ewing family tumors (EFT) are highly aggressive cancers for which drug resistance and relapse remain a constant clinical dilemma. Also, the cellular mechanism that underlies the difference between nonresponders and those who respond to treatment is still unclear. Here, we observed that the polysomy of chromosomes 8 and 12 and the clinical-pathological characteristics of EFT patients are related to the replication behavior of alleles of *KIF14* and *MDM4*/**PI3KC**2β loci.

The iFISH replication assay used here has already been proved sufficiently reliable for comparing the replication



**Fig. 6** Representation of FISH images of 1q32.1 region showing replication pattern in EFT TMA cells at interphase applying the three-color iFISH assay. *MDM4/P/3KC2B* signals hybridization are seen in green. *KIF14* signals hybridization are seen in orange and the control in *pink*. (A) cell with singlets (SS cells) in which neither allele has replicated; Synchronous pattern (B) cell with one singlet and one doublet (SD cells), which are Sphase cells in which one allele has replicated while its partner has not (*arrow*), (C) cell with doublets (DD cells) in which alleles have replicated; and (D) cell with multiplets pattern of signals (arrow).

status of one gene homolog relative to its counterpart in the same cell<sup>21,22,32,33</sup> ( $\succ$  Fig. 6).

We reported in the current study that the most frequent replication pattern seen in all cell populations was the asynchronous stage (SD), suggesting a loss of control of replication timing in EFT cells. Moreover, the two pairs of allelic KIF14 and MDM4/PI3KC2β loci replicated highly asynchronously in the group that showed gain than in the group with the absence of aberrations at 1q32.1. This higher frequency hemizygous replication pattern is probably a consequence of the genes' rearrangement in these EFT cells. Similar to, the asynchronous replication pattern of the c-MYC gene seen in lymphoma cells, in which the rearranged 5' end replicates later than the unaltered copy (Hayday et al, 1984; Calza et al, 1984). However, the allele-specific epigenetic alteration leading to functional hemizygosity of several genes within a single cell may also create a situation where a dominant oncogene is expressed in cells in which it is normally not expressed (Crossen, 1997), unlike the mechanism of LOH, which causes silencing of tumor suppressor genes. Histone acetylation and deacetylation states related to DNA methylation are also altered during gene regulation to affect transactivation of genes (Coolen et al., 2010). Yet, these genes may be located in the transition points between earlyand late-replicating domains where some fork barriers and repetitive elements are hot spots for DNA breaks and chromosome rearrangements<sup>34–36</sup> and, consequently, may be more prone to changes in gene sequence and in replication timing.

In the group that showed gain at 1g32.1, we noted that both KIF14 alleles had a rate of SS patterns lowered with a corresponding increase in the rate of SD, DD, and M patterns, meaning that at least one allele is early replicating compared with its normal replication schedule in some cells. For *MDM4*/**PI3KC**2β hybridization signals, we found a prevalence of the synchronous SS replication in the group without CNA but an increase in the SD and multiplet pattern with gain at 1q32.1 (**Fig. 2**), implying a loss of replication control in the cell cycle, which may lead to an overexpression of MDM4/ **PI3KC**2β in these cells. We demonstrated the existence of a high variability of cell-cycle phase among different EFT patient groups with CNA or without CNA. On the contrary, when the replication timing is highly synchronous (SS and DD), it has been indicated that a largely diploid chromosomal constitution is present (Selig et al., 1992).

Moreover, the polysomy of chromosome 8 showed significant statistical association with the SD replication pattern of *MDM4*/**PI3KC**2β, while the polysomy of chromosome 12 was statistically associated with the DD replication pattern of KIF14 for all populations. This may indicate a concentrationdependent requirement for KIF14 protein in chromosome disjunction, and failure of this association could contribute to cancer progression. Accordingly, other groups' data show that alteration of the replication pattern of chromosome regions may be one of the manifestations of genetic instability associated with increased risk for transformation.<sup>7,37</sup> We also observed a higher rate of asynchrony (SD) of the MDM4/ **PI3KC**2β locus in the trisomy 8 and 12 group compared with EFT patients without such aberrations, a finding which could reflect a correlation between the chromosomal aberration and gene replication leading to loss of cell-cycle control, as happen in chronic lymphocytic leukemia patients who presented with trisomy 12.<sup>7</sup>

It is clear that the EFT patients' cells show a high variability of replication pattern in normal (no CNA) and aberrant cells (with CNA) during the cell cycle, and the pattern that we detected in tumors with gain at 1q32.1 represents an average estimate among the cells. Furthermore, our results indicated that the relationship between replication pattern and clinical-pathological characteristics could be transmitted in the behavior of the genes' loci (1q32.1). Regarding our statistical results, there was a correlation between the synchronous (SS) KIF14 replication pattern and lung metastasis and age. Evidence has indicated that tissue-specific genes replicate early in the cell cycle and late in tissues in which they are silent. Although previous study indicates KIF14 as a molecular signature of poor prognosis in lung carcinoma, in our results linking the gene to lung metastasis, we demonstrate a late replication pattern, suggesting this gene locus must have undergone a switch in replication timing in an expressing cell type.<sup>36,38</sup> Besides, based on *in silico* analysis,<sup>39</sup> the entire *KIF14* gene was covered by a copy number variation region. In this manner, late replication due to inherent difficulties of replicating DNA could be a result of an accumulation of ssDNA that is more susceptible to damage and lost of control replication.<sup>35</sup> Additionally, in our findings, the early KIF14 allelic expression correlates with patients younger than

15 years of age. These younger patients had more DD replication pattern of KIF14 (mean 1.8 vs. 0.38) than patients older than 15. The DD synchronous pattern could be an indication of premutated genes.<sup>40</sup> Hence, we infer that in vounger patients premutations may be an early event in EFT tumorigenesis, as they may possess a higher actively replication mode needed for the endogenous events and cellcycle progression and more aggressive disease phenotype. Moreover, the negative correlation of time between onset of symptoms to diagnosis and synchronous replication pattern (DD) of the KIF14 locus suggests that patients who tolerate the symptoms longer have a lower rate of the DD pattern and potentially lower expression of KIF14, which is present during cell cycle. Noteworthy to say that no correlation was found between the treated or untreated group and the replication pattern of genes targeted at 1q32.1 locus. This is in line with a recent study that shows an abnormal pattern of replication (an epigenetic aberration) in lymphocytes with hematological disease at diagnosis is the same as that observed after chemotherapy but absent following allogeneic stem-cell transplantation. However, alteration in aneu-(a genetic aberration) that accompanies ploidy hematological malignancies, in contrast to epigenetic aberration, does not disappear following transplantation.<sup>6,41</sup> Conversely, within the treated group, we found different replication patterns between the treatment responders and nonresponders with the MDM4/PI3KC2β multiplet replication pattern (an asynchronous pattern). Patients who had poor treatment response showed lower frequency of the multiplet replication pattern of MDM4/PI3KC2ß genes. In non-Hodgkin lymphoma, asynchronous allelic replication is indicative of higher risk of relapse<sup>7</sup> and allelic-replication mode is a more accurate novel blood marker than the PSA antigen level in peripheral blood lymphocytes to differentiate between prostate cancer and benign prostate hyperplasia<sup>2</sup>. These findings led to the prospect of the MDM4/PI3KC2β multiplet replication pattern as a new genotoxic marker for different EFT nonresponder genotypes with a transcriptional and cell-cycle deregulation.

Following this line of thought, there is a notion that the association between the MDM4/PI3KC2β multiplet replication pattern and clinical response may be related to chemotherapy agents that can induce hypomethylation of DNA, which is the mechanism whereby genotoxic agents affect timing gene replication (Haaf, 1995) not discarding the possibility of being a reversible event after a stem cell transplantation, as described.<sup>41</sup> In this respect, it is possible to build a model that is consistent with our data-cytotoxic agents may bind to transcription factor DNA-binding domains, which attach to specific sequences of DNA adjacent to the genes that they regulate and leading to cell survival. Thus, this complex will control the aberrant replication timing and transcription for the transference of genetic information from DNA to mRNA. This is in line with the large amount of data showing correlation between allelicreplication timing and gene-expression profiles.<sup>1,23,33,36,42</sup> We suppose that in EFT patients, therapy resistance and eventual relapse can be triggered by aberrant regulation of the *MDM4*/*PI3KC*2 $\beta$  transcripts leading to escape from apoptosis, and therefore, tumor growth. Also, patients with worse treatment response have less multiplet hybridization signals<sup>42</sup>; however, they may have a more intricately disorientated molecular complex during the cell-cycle and DNA damage response. They may not only have deregulated activity of the DNA and RNA polymerases, but also may have more efficient DNA repair and cell-cycle checkpoint pathways leading to a deregulation of the DNA repair mechanism. Consequently, EFT tumors may survive due to generation of heterogeneous clonal cell populations.

Taken together, we have seen here a modified order of allelic replication in EFT patients at different genetic and clinical-pathological EFT groups, from which we suggest that the average of multiplet replication pattern of MDM4/ **PI3KC**2β may be an indicator of poor clinical response in EFT and the replication pattern of KIF14 may contribute to chromosomal instability in EFT patients. We feel that an cell survival could result from underlying epigenetic process in combination with genetic mechanisms that could arise from structural effects of drug/radiation therapy on DNA, deregulating replication, and transcription. Additionally, patients that respond poorly to treatment might have more efficient DNA damage repair pathways to escape from apoptosis than the responsive patients. Naturally, more experiments will be needed to confirm the intriguing correlations that we found between replication pattern and clinical-pathological features to extend the consistency of these data in EFT patients.

### Funding/Acknowledgments

Financial support was provided by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and Fundação de Apoio à Pesquisa do Estado de São Paulo (FAPESP).

# Conflict of Interest

None declared.

### References

- <sup>1</sup> Gunaratne PH, Nakao M, Ledbetter DH, Sutcliffe JS, Chinault AC. Tissue-specific and allele-specific replication timing control in the imprinted human Prader-Willi syndrome region. Genes Dev 1995;9(07):808–820
- 2 Dotan ZA, Dotan A, Ramon J, Avivi L. Altered mode of allelic replication accompanied by aneuploidy in peripheral blood lymphocytes of prostate cancer patients. Int J Cancer 2004;111(01): 60–66
- 3 Bergman Y, Cedar H. A stepwise epigenetic process controls immunoglobulin allelic exclusion. Nat Rev Immunol 2004;4 (10):753-761
- 4 Gimelbrant AA, Ensminger AW, Qi P, Zucker J, Chess A. Monoallelic expression and asynchronous replication of p120 catenin in mouse and human cells. J Biol Chem 2005;280(02):1354–1359
- 5 Squire JA, Li M, Perlikowski S, et al. Alterations of H19 imprinting and IGF2 replication timing are infrequent in Beckwith-Wiedemann syndrome. Genomics 2000;65(03):234–242
- 6 Grinberg-Rashi H, Cytron S, Gelman-Kohan Z, Litmanovitch T, Avivi L. Replication timing aberrations and aneuploidy in peripheral blood lymphocytes of breast cancer patients. Neoplasia 2010; 12(08):668–674

- 7 Amiel A, Levi E, Reish O, Sharony R, Fejgin MD. Replication status as a possible marker for genomic instability in cells originating from genotypes with balanced rearrangements. Chromosome Res 2001;9(08):611–616
- 8 Lengauer C, Kinzler KW, Vogelstein B. Genetic instability in colorectal cancers. Nature 1997;386(6625):623-627
- 9 Schvartzman JM, Sotillo R, Benezra R. Mitotic chromosomal instability and cancer: mouse modelling of the human disease. Nat Rev Cancer 2010;10(02):102–115
- 10 Korenstein-Ilan A, Amiel A, Lalezari S, Lishner M, Avivi L. Allelespecific replication associated with aneuploidy in blood cells of patients with hematologic malignancies. Cancer Genet Cytogenet 2002;139(02):97–103
- 11 Armengol G, Tarkkanen M, Virolainen M, et al. Recurrent gains of 1q, 8 and 12 in the Ewing family of tumours by comparative genomic hybridization. Br J Cancer 1997;75(10):1403–1409
- 12 Mugneret F, Lizard S, Aurias A, Turc-Carel C. Chromosomes in Ewing's sarcoma. II. Nonrandom additional changes, trisomy 8 and der(16)t(1;16). Cancer Genet Cytogenet 1988;32(02): 239–245
- 13 Kullendorff CM, Mertens F, Donnér M, Wiebe T, Akerman M, Mandahl N. Cytogenetic aberrations in Ewing sarcoma: are secondary changes associated with clinical outcome? Med Pediatr Oncol 1999;32(02):79–83
- 14 Zielenska M, Zhang ZM, Ng K, et al. Acquisition of secondary structural chromosomal changes in pediatric Ewing sarcoma is a probable prognostic factor for tumor response and clinical outcome. Cancer 2001;91(11):2156–2164
- 15 Hattinger CM, Rumpler S, Ambros IM, et al. Demonstration of the translocation der(16)t(1;16)(q12;q11.2) in interphase nuclei of Ewing tumors. Genes Chromosomes Cancer 1996;17(03): 141–150
- 16 Corson TW, Gallie BL. KIF14 mRNA expression is a predictor of grade and outcome in breast cancer. Int J Cancer 2006;119(05): 1088–1094
- 17 Cahill DP, Lengauer C, Yu J, et al. Mutations of mitotic checkpoint genes in human cancers. Nature 1998;392(6673):300–303
- 18 Carleton M, Mao M, Biery M, et al. RNA interference-mediated silencing of mitotic kinesin KIF14 disrupts cell cycle progression and induces cytokinesis failure. Mol Cell Biol 2006;26(10): 3853–3863
- 19 Corson TW, Huang A, Tsao MS, Gallie BL. KIF14 is a candidate oncogene in the 1q minimal region of genomic gain in multiple cancers. Oncogene 2005;24(30):4741–4753
- 20 Abdel-Fatah TM, Powe DG, Agboola J, et al. The biological, clinical and prognostic implications of p53 transcriptional pathways in breast cancers. J Pathol 2010;220(04):419–434
- 21 Selig S, Okumura K, Ward DC, Cedar H. Delineation of DNA replication time zones by fluorescence in situ hybridization. EMBO J 1992;11(03):1217–1225
- 22 Haaf T. The effects of 5-azacytidine and 5-azadeoxycytidine on chromosome structure and function: implications for methylation-associated cellular processes. Pharmacol Ther 1995;65(01): 19–46Review
- 23 Méndez J. Temporal regulation of DNA replication in mammalian cells. Crit Rev Biochem Mol Biol 2009;44(05):343–351Review
- 24 Costa CML, Rondinelli P, Campbell B. Tumor neuroectodérmico primitivo da infância: Relato de 13 casos e Revisão de Literatura. J Oncol 2000;46:293–298
- 25 Merscher S, Marondel I, Pedeutour F, Gaudray P, Kucherlapati R, Turc-Carel C. Identification of new translocation breakpoints at 12q13 in lipomas. Genomics 1997;46(01):70–77

- 26 Sirvent N, Forus A, Lescaut W, et al. Characterization of centromere alterations in liposarcomas. Genes Chromosomes Cancer 2000;29(02):117–129
- 27 Selvarajah S, Yoshimoto M, Ludkovski O, et al. Genomic signatures of chromosomal instability and osteosarcoma progression detected by high resolution array CGH and interphase FISH. Cytogenet Genome Res 2008;122(01):5–15
- 28 Ko E, Rademaker A, Martin R. Microwave decondensation and codenaturation: a new methodology to maximize FISH data from donors with very low concentrations of sperm. Cytogenet Cell Genet 2001;95(3-4):143–145
- 29 Sugimura H. Detection of chromosome changes in pathology archives: an application of microwave-assisted fluorescence in situ hybridization to human carcinogenesis studies. Carcinogenesis 2008;29(04):681–687Review
- 30 Mukherjee AB, Murty VV, Chaganti RS. Detection of cell-cycle stage by fluorescence in situ hybridization: its application in human interphase cytogenetics. Cytogenet Cell Genet 1992;61 (02):91–94
- 31 Böhling T, Bacchini P, Bertoni F, et al. Diagnosis and tumor response in osteosarcoma and Ewing's sarcoma. Acta Orthop Scand Suppl 2004;75(311):72–76
- 32 Knoll JH, Cheng SD, Lalande M. Allele specificity of DNA replication timing in the Angelman/Prader-Willi syndrome imprinted chromosomal region. Nat Genet 1994;6(01):41–46
- 33 White LM, Rogan PK, Nicholls RD, Wu BL, Korf B, Knoll JH. Allelespecific replication of 15q11-q13 loci: a diagnostic test for detection of uniparental disomy. Am J Hum Genet 1996;59(02): 423–430
- 34 Rothstein R, Michel B, Gangloff S. Replication fork pausing and recombination or "gimme a break". Genes Dev 2000;14(01): 1–10Review
- 35 Watanabe G, Shimizu K. DNA sequence analysis of long PCR amplified products at the D1S80 locus. Leg Med (Tokyo) 2002;4 (01):37–39
- 36 Cohen SM, Furey TS, Doggett NA, Kaufman DG. Genome-wide sequence and functional analysis of early replicating DNA in normal human fibroblasts. BMC Genomics 2006;7:301
- 37 Ekholm-Reed S, Spruck CH, Sangfelt O, et al. Mutation of hCDC4 leads to cell cycle deregulation of cyclin E in cancer. Cancer Res 2004;64(03):795–800
- 38 Reish O, Orlovski A, Mashevitz M, et al. Modified allelic replication in lymphocytes of patients with neurofibromatosis type 1. Cancer Genet Cytogenet 2003;143(02):133–139
- 39 Ferreira BI, Alonso J, Carrillo J, et al. Array CGH and gene-expression profiling reveals distinct genomic instability patterns associated with DNA repair and cell-cycle checkpoint pathways in Ewing's sarcoma. Oncogene 2008;27(14):2084–2090
- 40 Yeshaya J, Shalgi R, Shohat M, Avivi L. FISH-detected delay in replication timing of mutated FMR1 alleles on both active and inactive X-chromosomes. Hum Genet 1999;105(1-2):86–97
- 41 Nagler A, Cytron S, Mashevich M, Korenstein-Ilan A, Avivi L. The aberrant asynchronous replication—characterizing lymphocytes of cancer patients—is erased following stem cell transplantation. BMC Cancer 2010;10:230
- 42 Hamdan SM, van Oijen AM. Timing, coordination, and rhythm: acrobatics at the DNA replication fork. J Biol Chem 2010;285(25): 18979–18983
- 43 Shetty D, Jain H, Rohil Y, et al. Role of cytogenetic abnormalities detected by fluorescence *in situ* hybridization as a prognostic marker: pathogenesis & clinical course in patients with B-chronic lymphocytic leukaemia. Indian J Med Res 2021;153(04):475–483