



Surgical treatment of temporal lobe epilepsy: comparative results of selective amygdalohippocampectomy versus anterior temporal lobectomy from a referral center in Brazil

Tratamento cirúrgico da epilepsia do lobo temporal: resultados comparativos da amigdalohipocampectomia seletiva versus lobectomia temporal anterior em um centro de referência no Brasil

Laryssa Crystinne Azevedo Almeida¹ Vanessa Alves Lobato¹ Maria do Carmo Vasconcelos Santos¹
Aline Curcio de Moraes¹ Bruno Silva Costa¹

¹Santa Casa de Misericórdia de Belo Horizonte, Departamento de Neurologia e Neurocirurgia, Belo Horizonte MG, Brazil.

Address for correspondence Laryssa C. Azevedo Almeida (email: laryssazevedo@hotmail.com).

Arq. Neuropsiquiatr. 2023;81:647–655.

Abstract

Background Temporal lobe epilepsy (TLE) is a high prevalence neurological disorder. Surgery has emerged as a promising treatment.

Objective The objective of this work is to compare the surgical results of anterior temporal lobectomy (ATL) versus selective amygdalohippocampectomy (SAH) in a cohort of 132 patients.

Methods We performed a retrospective study of 146 patients operated for TLE from 2008 to 2019. Initially, 13 patients were excluded from the study due to insufficient medical record data or follow-up loss. One patient was excluded from the analysis of the results due to death in the first postoperative week. We used the ILAE scale to classify seizure control after surgery. In patients with left hippocampal sclerosis, SAH was performed and in right temporal lobe epilepsy, ATL was the approach of choice.

Results The mean follow-up time after surgery was 57.2 months (12–137). In our data analysis, we found that the group of patients undergoing ATL had a higher prevalence of being completely seizure-free (ILAE I) (57.1% versus 31%) and a higher rate of satisfactory seizure control (88.6% versus 69.3%) $p = 0,006$, when compared with patients undergoing SAH.

Conclusion The literature is still controversial about seizure control concerning the technique used due to the lack of a robust methodology. Our data analysis identified the superiority of ATL over SAH in seizure outcomes. ATL may be the best option for

Keywords

- ▶ Epilepsy, Temporal Lobe
- ▶ Hippocampal Sclerosis
- ▶ Anterior Temporal Lobectomy
- ▶ Epilepsies, Partial

received
February 4, 2023
received in its final form
April 5, 2023
accepted
April 12, 2023

DOI <https://doi.org/10.1055/s-0043-1771172>.
ISSN 0004-282X.

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>).

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

Resumo**Palavras-chave**

- ▶ Epilepsia do Lobo Temporal
- ▶ Esclerose Hipocampal
- ▶ Lobectomia Temporal Anterior
- ▶ Epilepsias Parciais

adequately controlling seizures with minimal additional morbidity in countries with a cost limitation for extended propaedeutics.

Antecedentes A epilepsia do lobo temporal (TLE) é uma desordem neurológica de alta prevalência. A cirurgia surgiu como um tratamento promissor.

Objetivo O objetivo deste trabalho é comparar os resultados da lobectomia temporal anterior (ATL) versus amigdalohipocampectomia seletiva (SAH) em uma coorte de 132 pacientes.

Métodos Realizamos um estudo retrospectivo de 146 pacientes operados por TLE de 2008 a 2019. Inicialmente, 13 pacientes foram excluídos por insuficiência de dados em prontuário ou perda de seguimento. Um paciente foi excluído da análise por óbito na primeira semana de pós-operatório. Usamos a escala ILAE para classificar o controle das crises após a cirurgia. Em pacientes com esclerose hipocampal à esquerda, foi realizada a SAH, e na epilepsia do lobo temporal à direita, a ATL foi a abordagem de escolha.

Resultados O tempo médio de seguimento após a cirurgia foi de 57,2 meses (12–137). Em nossa avaliação, encontramos que o grupo de pacientes submetidos à ATL apresentou maior prevalência de ausência total de crises (ILAE I) (57,1% versus 31%) e maior taxa de controle satisfatório da epilepsia (88,6% versus 69,3%) $p=0,006$, quando comparado ao grupo submetido à SAH.

Conclusão A literatura ainda é controversa em relação à redução das crises de acordo com a técnica utilizada devido a falta de uma metodologia robusta. Nosso estudo identificou superioridade da ATL sobre a SAH nos desfechos convulsivos. ATL pode ser a melhor opção para controlar adequadamente as convulsões com morbidade adicional mínima em países com limitação de custo para propedêutica estendida.

INTRODUCTION

Temporal lobe epilepsy (TLE) is the most common human epileptic syndrome¹ being a disabling and progressive entity.² In addition to seizures, which already represent clinical management difficulties, they may also be associated with cognitive, language, or psychiatric disorders.^{3,4} A tendency to drug refractoriness characterizes it, and up to a third of patients are drug-resistant.⁵ In recent years, surgery has proven to be a therapeutic option with good results, with a controlled clinical trial demonstrating its superiority to drug treatment alone.⁶

The hippocampal sclerosis etiology is multifactorial, typically caused by inflammatory, infectious insults, trauma, or febrile seizures.^{7,8}

There are technical variations in TLE surgery, and there is no consensus on the best surgical approach. The most common techniques are Anterior temporal lobectomy (ATL) and selective amigdalohipocampectomia (SAH). Foerster pioneered subtotal temporal lobectomy in 1925.⁹ Falconer developed en bloc resection of the temporal lobe and mesial structures in 1953.¹⁰ Morris, in 1956 used the term standard temporal lobectomy for a 6.5 cm resection of the temporal lobe,^{11,12} Spencer refined this technique with 4.5 cm in the non-dominant cortex and 3 cm in the dominant cortex. Niemeyer, in 1958, described a transventricular selective access to mesial temporal structures through an

incision in the medial temporal lobe.¹³ Wieser and Yasargil proposed a transsylvian approach to the amygdala and hippocampus.¹⁴

There is still controversy about the best surgical approach for mesial temporal epilepsy.¹⁵ Elseways, selective resections of mesial structures could have less cognitive effects, whereas an anterior temporal lobectomy has better seizure control.

In a review in 2008, Schramm¹⁶ cites eight studies that compared selective surgery against temporal lobectomy concerning seizure control. In six of these studies, the authors found no difference in seizure control despite the surgical approach. ATL was more effective in two papers, one in children.^{17,18} Josephson¹⁹ compared ATL and SAH in a meta-analysis of 13 articles and 1203 patients, showing better control of seizures in ATL.

METHODS**Ethical statement**

All patients signed an informed consent, and the study was conducted following the Declaration of Helsinki. The national ethics board approved the study.

Participants and evaluation

A retrospective study was performed based on the medical records of 146 patients operated on for temporal lobe

Table 1 ILAE outcome classification

Outcome classification	Definition
1	Completely seizure free; no auras
2	Only auras; no other seizures
3	One to three seizure days per year; \pm auras
4	Four seizure days per year to 50% reduction of baseline seizure days; \pm auras
5	Less than 50% reduction of baseline seizure days to 100% increase of baseline seizure days; \pm auras
6	More than 100% increase of baseline seizure days; \pm auras

epilepsy from 2008 to 2019. The ILAE classification²⁰ (**Table 1**) was used to determine the degree of seizure control, and we compared the descriptive results according to the technique used.

The preoperative evaluation of these patients included neuropsychological testing, video-EEG, and high-resolution MRI. In cases where the video-EEG with scalp electrodes failed to define the temporal lobe as an epileptogenic source, a foramen ovale electrode was implanted as a complementary method. We included only patients with unilateral hippocampal sclerosis on MRI and concordant epileptic onset on video-EEG. Non-invasive options such as functional MRI would add additional costs and time, being a method not

exempt from clinical differences.²¹ Postoperatively, an MRI was requested for all patients to assess whether the resection area was satisfactory.

Approach and selection of groups

The same surgeon performed all surgeries. In right-sided hippocampal sclerosis, a temporal lobectomy and hippocampectomy was performed using the Spencer technique,¹² resecting at least 3.5 cm of the anterior border of the temporal lobe (**Figures 1** and **2**). In left hippocampal sclerosis, a selective amygdalohippocampectomy was used as described by Niemeyer (**Figures 3** and **4**).¹³

The surgical procedure of hippocampal resection is accompanied by the removal of other mesial structures, including the uncus, amygdala, and parahippocampal gyrus. The resection of the hippocampus and parahippocampal gyrus should be performed as posteriorly as possible, extending at least up to the level of the lateral mesencephalic sulcus.

The rationale for the choice of surgical access

The choice between ATL and SAH based on the sclerosis side followed the following rationale:

- All patients have typical temporal lobe seizures;
- The Video-EEG showed seizures with semiology and a typical electrographic pattern;
- Volumetric MRI showed no lesions other than unilateral hippocampal sclerosis.

In Brazil and several other developing countries, the cost of performing surgeries is often a major obstacle. The use of




Figure 1 MRI (Coronal, axial and sagittal) pre and postoperative of surgery with Anterior temporal lobectomy for right hippocampal sclerosis. (A, B, C) Preoperative MRI showing right hippocampal sclerosis. (D, E, F) Postoperative MRI showing excision of the neocortex (3,5 cm) and mesial structures.




Figure 2 Surgical photos of anterior temporal lobectomy for right hippocampal sclerosis. (A) Surgical view after anterior temporal lobectomy. Mesial structures not yet resected. (B) Resection of mesial structures. Hippocampus and parahippocampal gyrus already disconnected.




Figure 3 MRI (Coronal IR and sagittal T1 volumetric) pre and postoperative of surgery with selective access for left hippocampal sclerosis. (A) Preoperative coronal MRI. Collateral groove shows limit of resection of mesial structures. (B) Preoperative sagittal MRI. Arrows show the hippocampus and amygdala. (C and D) Details of the resection of the mesial structures in coronal and sagittal sections.

invasive research to individualize access points based on electrophysiological details can significantly increase expenses, rendering procedures impossible to perform. The cost of public healthcare for the evaluation and surgical treatment of temporal lobe epilepsy, including medical fees and hospital costs, is roughly equivalent to 1200 US dollars. In this context, the addition of extensive invasive monitoring is not feasible.

On the other hand, there is a chronic shortage of epilepsy medication in the public health system, leading to uncertain clinical treatment. This further underscores the value of safe surgical interventions that can produce positive outcomes, even in the context of limited resources.

The literature shows that both accesses have excellent results in seizure control and neuropsychological outcome.^{16,17} Despite overall good results, some studies show




Figure 4 Surgical photos of selective access for left hippocampal sclerosis. (A) Initial view of the hippocampus after corticectomy. In dotted lines, the collateral eminence, lateral limit of the resection of the mesial structures. (B) After lateral disconnection at the level of the collateral and medial eminence in the ambient cistern, we visualize the basal vein of Rosenthal and the fimbria. (C) After resection of the mesial structures, we visualize the vessels of the peduncular cistern, third nerve and peduncle protected by the arachnoidal plane. (D) final view with the corticectomy area.

a worse language performance in patients operated with left ATL. Similarly, SAH would have a worse outcome in epilepsy control.^{17,18,22–24}

Statistical analysis

We utilized contingency tables to compare the two groups, analyzing the results in relation to the type of surgery performed (ATL or SAH). Additionally, we conducted a Kaplan-Meier survival analysis to examine the occurrence of seizures, the patient's ILAE 1 status, and good outcome (ILAE 1 to 3).

The data were analyzed with the IBM SPSS Statistics Software. A *p*-value of less than 0.05 was considered significant.

RESULTS

Our database included 146 patients who underwent surgery for temporal lobe epilepsy secondary to hippocampal sclerosis between 2008 and 2019. Thirteen patients were excluded from our initial analysis due to incomplete medical records or follow-up loss. One patient died from pulmonary thromboembolism in the immediate postoperative period, resulting in a mortality rate of 0.06% and their exclusion from the analysis. The remaining 132 patients, comprising 72 females (53%) and 60 males (47%), were evaluated. The mean age at the time of surgery was 37.85 years (range: 9–65 years).

Hippocampal sclerosis was present on the right side in 70 patients (53%) and on the left side in 62 patients (47%). All the surgical cases underwent anatomopathological and immunohistochemical analysis of the resected tissues, which revealed no abnormalities or neuronal depopulation suggestive of hippocampal sclerosis. There was no statistical difference between age, sex, and follow-up time in the two groups (ATL and SAH), as shown in ►Table 2. The mean follow-up time after surgery was 57.2 months (range: 12–137 months).


In the follow-up period, 66 patients (50%) had at least one seizure (excluding events within the first 30 days after surgery). At the end of the follow-up period, 105 patients (79.5%) had achieved an ILAE score of 1–3, indicating a good result. Of the patients who underwent ATL, 62 (88.6%) obtained a good result, compared with 43 (69.3%) in the SAH group (*p* = 0.006). Only 58 patients (43.9%) were on ILAE 1 at the end of the follow-up period, with 40 (57.1%) in the ATL group and 18 (31.0%) in the SAH group (*p* = 0.001), as summarized in ►Table 2.

The average interval until an epileptic event was 22.8 months (range: 1–86.1 months). Patients who underwent ATL had a mean time to the first seizure of 23.04 months, compared with 21.86 months in those who underwent SAH, with no statistical difference (*p* = 0.82), as shown in ►Figure 5.

Table 2 Frequencies comparing anterior temporal lobectomy (ATL) and selective amygdalo hippocampectomy (SAH) groups for the treatment of hippocampal sclerosis (HS)

		Total	ATL Right HS	SAH Left HS	p
n		132	70 (53.0%)	62 (47.0%)	
Gender	M	60 (47.0%)	30 (42.9%)	30 (48.4%)	
	F	72 (53.0%)	40 (57.1%)	32 (51.6%)	
Mean age in years		37.85	36.94	38.87	
Mean follow up in months		57.2	55.7	58.8	
ILAE	1	58 (43.9%)	40 (57.1%)	18 (31.0%)	p = 0.001
	2	2 (1.5%)	0 (0.0%)	2 (3.2%)	
	3	45 (34.1%)	22 (31.4%)	23 (37.1%)	
	1 to 3*	105 (79.5%)	62 (88.6%)	43 (69.3%)	p = 0.006
	4	16 (12.1%)	5 (7.1%)	11 (17.7%)	
	5	11 (8.3%)	3 (4.3%)	8 (12.9%)	
	6	0 (0.0%)	0 (0.0%)	0 (0.0%)	

Note: * ILAE 1 to 3 are considered as good result.




Abbreviations: ATL, anterior temporal lobectomy; SAH, selective amygdalohippocampectomy. Notes: Student *t*-test between surgery groups: $p = 0.82$. ATL mean 23.04 months; SAH mean 21.86 months. **Figure 5** Event free interval comparing the ATL and SAH groups.

The Kaplan-Meier mortality curves (► **Figure 6**), using the Log Rank (Mantel-Cox) statistical analysis with seizure as the event, showed a significant difference between the ATL and SAH groups ($p = 0.024$).

Our surgical morbidity rate was 11.8% (17/143), and mortality was 0.6% (1/143), consistent with the results described in the literature. The patient who died in the first postoperative week due to PTE was excluded from the analysis of results regarding epilepsy control. Other complications are summarized in ► **Table 3**.

DISCUSSION

The primary purpose of surgery is to control seizures. Maintaining a good functional status of patients is also mandatory. The search for a more selective resection is based



Abbreviations: ATL, anterior temporal lobectomy; SAH, selective amygdalohippocampectomy. Statistical analysis using Mantel-Cox Log Rank. $p = 0.024$.

Figure 6 Kaplan-Meier survival analysis with seizure as event, comparing ATL and SAH.

on not worsening memory and language deficits, especially in the dominant hemisphere. It has been shown in several articles that the selective resection of mesial structures has a benefit, even if marginal, in the cognitive assessment of patients.²⁴⁻²⁷ Other authors do not show differences between ATL and SAH regarding neuropsychological prognosis.^{27,28} Helmstaedter²³ postulates that the cognitive deficit after eloquent temporal resection could be more linked to perioperative cortical injury, visible in post-surgical MRI, than to the type of resection.

The only multicenter randomized study showing the outcome of surgery in terms of seizure control was

Table 3 Complications reported in our case series

Age, sex	Complication
29, Male	Cardiorespiratory arrest in anesthetic induction
41, Male	Wound infection, cranioplasty, chronic headache
25, Female	Wound infection
43, Female	Wound infection
20, Female	Wound infection
20, Female	Wound infection
40, Female	Memory impairment
41, Female	Memory impairment
30, Female	Memory impairment
48, Male	CSF leak submitted to external lumbar shunt, acute subdural hematoma, decompressive craniectomy, wound infection, debridement, cranioplasty.
46, Female	Postoperative hemiparesis with subsequent recovery
48, Female	Visual field disorder
41, Female	Visual field disorder
51, Female	Visual field disorder
27, Male	Visual field disorder
65, Male	Intraparenchymal hematoma
39, Female	Deep vein thrombosis

Abbreviation: CSF, cerebrospinal fluid.

performed by Wiebe et al. in 2001. In the group of patients operated on, always by ATL, 38% were completely free of seizures (Engel 1). Concerning the control of epilepsy, there are variable results in the literature when comparing SAH and ATL. Several authors show equality in seizure control,^{29–32} while others show better results in ATL.^{18,19} Clusmann,¹⁹ in a series of 89 children and adolescents, found a worse result in seizure control in SAH when compared with ATL. Also, patients with left hippocampal sclerosis had a worse result despite the surgical technique. In a meta-analysis including 13 studies and 1203 patients, Josephson¹⁹ found better control of seizures in ATL than in SAH.

Our evaluation sought to compare the two types of surgery, not only in terms of good surgical outcomes but also in the occurrence of seizures and the complete control of seizures (ILAE 1). The statistical analysis show a better outcome in occurrence of seizures ($p = 0.005$), ILAE 1 final result ($p = 0.001$) and good result ($p = 0.006$) in favor of ATL. We performed Kaplan Meyer's survival analysis having a seizure as the target event. The curves showed better results in ATL over SAH. The Log Rank statistical analysis significantly favors the ATL group ($p = 0.029$). The Kaplan Meier graphs show a downward curve with a progressive worsening of the results over time, consistent with the findings described in the literature. The results are consistent with a better surgi-

cal prognosis in resections that include the temporal neocortex over more selective resections.

Interestingly, the interval until the first epileptic event was similar when comparing the two groups. However, patients in the SAH group had seizures more often after this initial event (–**Figure 5**). This finding could be related to the maintenance of an altered neuronal network in patients with more selective resection, which quickly resumes the pattern of seizures after a first ictal event.²² The same difficulty of resources that makes it difficult to carry out a more individualized approach to cases makes the surgery attractive from a cost-benefit point of view.

As the procedure is safe with very low morbidity and mortality, surgery proves to be an effective procedure for our reality. Clinical treatment has a significant cost and, in most cases, it is paid for by the state, which cannot maintain this treatment without interruptions due to lack of funds.

Limitations of the study

We did not assess neuropsychological, speech, and language differences between groups, as all surgeries in the left hemisphere were SAH. In our series, preoperative tests to determine hemispheric dominance for language were not performed. By convention, all cases on the left (predominantly dominant hemisphere in humans^{33,34}) were submitted to SAH, and those on the right to ATL, the interpretation of our results is limited. It is essential to mention that Clusmann,¹⁹ in 2004, observed that surgeries in the dominant hemisphere had worse results despite the technique used. It is a possible bias in our work. Despite these limitations, we believe that the data obtained adds relevance to the discussion about the difference in results regarding the technique used.

We excluded all patients who had MRI lesions other than HS to avoid a worse result in selective surgeries for patients who had lesions in the temporal neocortex. However, MRI is not always able to clearly show small dysplastic lesions.

The postoperative evaluation time was long enough to show the differences between the two techniques. As there is a tendency for the results to progressively worsen over time, as demonstrated in our survival curves, a longer follow-up perhaps showed that the two techniques tend to match up with a longer follow-up.

In conclusion, there is still controversy about the influence of more selective procedures in surgery for ATL epilepsy. Our work has shown better results concerning the control of epilepsy when we use ATL compared with SAH. Despite the varied results of articles on the subject, our data show that performing ATL may be more effective in controlling epilepsy, emphasizing the importance of assessing language and memory before and after surgery to define the standard of comparison between the two surgical techniques. Whenever possible, the choice of access route should be made individually for each patient, based on neurophysiological and imaging findings. In countries with a cost limitation for extended propaedeutics, ATL may be the best option for the proper control of seizures with minimal additional morbidity.

Authors' Contributions

LCAA: conceptualization, data curation, formal analysis, investigation, methodology, writing – original draft, writing – review & editing; VAL: conceptualization, data curation, formal analysis, investigation, writing – original draft; MCVS: conceptualization, data curation, investigation, supervision; ACM: conceptualization, data curation, investigation; BSC: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, supervision, validation, writing – review & editing.

Conflict of Interest

There is no conflict of interest to declare.

References

- Mohan M, Keller S, Nicolson A, et al. The long-term outcomes of epilepsy surgery. *PLoS One* 2018;13(05):e0196274. Doi: 10.1371/journal.pone.0196274
- Cascino GD. Temporal lobe epilepsy is a progressive neurologic disorder: Time means neurons!. *Neurology* 2009;72(20):1718–1719. Doi: 10.1212/wnl.0b013e3181a4e465
- Shahani L, Cervenka G. Impact of surgical intervention on seizure and psychiatric symptoms in patients with temporal lobe epilepsy. *BMJ Case Rep* 2019;12(07):e229242. Doi: 10.1136/bcr-2019-229242
- Engel J. Jr PT. (1998). *Epilepsy: A Comprehensive Textbook*. 2nd ed. Archives of Neurology.; Philadelphia: Lippincott Williams & Wilkins; 2008 1373–1374 pp. Doi: 10.1001/archneur.55.10.1373
- Pascual MR. Temporal lobe epilepsy: clinical semiology and neurophysiological studies. *Semin Ultrasound CT MR* 2007;28(06):416–423. Doi: 10.1053/j.sult.2007.09.004
- Wiebe S, Blume WT, Girvin JP, Eliasziw M Effectiveness and Efficiency of Surgery for Temporal Lobe Epilepsy Study Group. A randomized, controlled trial of surgery for temporal-lobe epilepsy. *N Engl J Med* 2001;345(05):311–318
- Costa BS, Santos MCV, Rosa DV, Schutze M, Miranda DM, Romano-Silva MA. Automated evaluation of hippocampal subfields volumes in mesial temporal lobe epilepsy and its relationship to the surgical outcome. *Epilepsy Res* 2019;154:152–156. Doi: 10.1016/j.eplepsyres.2019.05.011
- Rosa DV, Rezende VB, Costa BS, et al. Circulating CD4 and CD8 T cells expressing pro-inflammatory cytokines in a cohort of mesial temporal lobe epilepsy patients with hippocampal sclerosis. *Epilepsy Res* 2016;120:1–6. Doi: 10.1016/j.eplepsyres.2015.11.011
- Feindel W, Leblanc R, de Almeida AN. Epilepsy surgery: historical highlights 1909–2009. *Epilepsia* 2009;50(Suppl 3):131–151. Doi: 10.1111/j.1528-1167.2009.02043.x
- Hill D, Falconer MA, Pampiglione G, Liddell DW. Discussion on the surgery of temporal lobe epilepsy. *Proc R Soc Med* 1953;46(11):965–976. Doi: 10.1177/00359157530460112
- Morris AA. Temporal lobectomy with removal of uncus, hippocampus, and amygdala; results for psychomotor epilepsy three to nine years after operation. *AMA Arch Neurol Psychiatry* 1956;76(05):479–496
- Spencer DD, Spencer SS, Mattson RH, Williamson PD, Novelly RA. Access to the posterior medial temporal lobe structures in the surgical treatment of temporal lobe epilepsy. *Neurosurgery* 1984;15(05):667–671. Doi: 10.1097/00006123-198411000-00005
- Niemeyer P. (1958). The transventricular amygdalohippocampectomy in temporal lobe epilepsy. *Temporal Lobe Epilepsy*
- Wieser HG, Yaşargil MG. Selective amygdalohippocampectomy as a surgical treatment of mesial limbic epilepsy. *Surg Neurol* 1982;17(06):445–457. Doi: 10.1016/s0090-3019(82)80016-5
- Mansouri A, Fallah A, McAndrews MP, et al. Neurocognitive and Seizure Outcomes of Selective Amygdalohippocampectomy versus Anterior Temporal Lobectomy for Mesial Temporal Lobe Epilepsy. *Epilepsy Res Treat* 2014;2014(306382):306382. Doi: 10.1155/2014/306382
- Schramm J. Temporal lobe epilepsy surgery and the quest for optimal extent of resection: a review. *Epilepsia* 2008;49(08):1296–1307. Doi: 10.1111/j.1528-1167.2008.01604.x
- Moshir Estekhahreh SS, Saghebdoost S, Zare R, Hakak MA, Hashemabadi BAG. Memory and executive functioning outcomes of selective amygdalohippocampectomy in patients with hippocampal sclerosis: A preliminary study in a developing country. *Surg Neurol Int* 2022;13:161. Doi: 10.25259/SNI_49_2022
- Bate H, Eldridge P, Varma T, Wiesmann UC. The seizure outcome after amygdalohippocampectomy and temporal lobectomy. *Eur J Neurol* 2007;14(01):90–94. Doi: 10.1111/j.1468-1331.2006.01565.x
- Clusmann H, Kral T, Gleissner U, et al. Analysis of different types of resection for pediatric patients with temporal lobe epilepsy. *Neurosurgery* 2004;54(04):847–859, discussion 859–860. Doi: 10.1227/01.neu.0000114141.37640.37
- Wieser HG, Blume WT, Fish D, et al; Commission on Neurosurgery of the International League Against Epilepsy (ILAE) ILAE Commission Report. Proposal for a new classification of outcome with respect to epileptic seizures following epilepsy surgery. *Epilepsia* 2001;42(02):282–286
- Omisade A, O'Grady C, Sadler RM. Divergence between functional magnetic resonance imaging and clinical indicators of language dominance in preoperative language mapping. *Hum Brain Mapp* 2020;41(14):3867–3877. Doi: 10.1002/hbm.25092
- Josephson CB, Dykeman J, Fiest KM, et al. Systematic review and meta-analysis of standard vs selective temporal lobe epilepsy surgery. *Neurology* 2013;80(18):1669–1676. Doi: 10.1212/wnl.0b013e3182904f82
- Helmstaedter C, Van Roost D, Clusmann H, Urbach H, Elger CE, Schramm J. Collateral brain damage, a potential source of cognitive impairment after selective surgery for control of mesial temporal lobe epilepsy. *J Neurol Neurosurg Psychiatry* 2004;75(02):323–326
- Clusmann H, Schramm J, Kral T, et al. Prognostic factors and outcome after different types of resection for temporal lobe epilepsy. *J Neurosurg* 2002;97(05):1131–1141. Doi: 10.3171/jns.2002.97.5.1131
- Rydenhag B, Silander HC. Complications of epilepsy surgery after 654 procedures in Sweden, September 1990–1995: a multicenter study based on the Swedish National Epilepsy Surgery Register. *Neurosurgery* 2001;49(01):51–56, discussion 56–57. Doi: 10.1227/00006123-200107000-00007
- Helmstaedter C, Reuber M, Elger CC. Interaction of cognitive aging and memory deficits related to epilepsy surgery. *Ann Neurol* 2002;52(01):89–94. Doi: 10.1002/ana.10260
- Roessler K, Kasper BS, Shawarba J, et al. Operative variations in temporal lobe epilepsy surgery and seizure and memory outcome in 226 patients suffering from hippocampal sclerosis. *Neurol Res* 2021;43(11):884–893. Doi: 10.1080/01616412.2021.1942407
- Nascimento FA, Gatto LA, Silvado C, Mäder-Joaquim MJ, Moro MS, Araujo JC. Anterior temporal lobectomy versus selective amygdalohippocampectomy in patients with mesial temporal lobe epilepsy. *Arq Neuropsiquiatr* 2016;74(01):35–43. Doi: 10.1590/0004-282x20150188
- Wolf RL, Ivnik RJ, Hirschorn KA, Sharbrough FW, Cascino GD, Marsh WR. Neurocognitive efficiency following left temporal lobectomy: standard versus limited resection. *J Neurosurg* 1993;79(01):76–83. Doi: 10.3171/jns.1993.79.1.0076

- 30 Paglioli E, Palmi A, Portuquez M, et al. Seizure and memory outcome following temporal lobe surgery: selective compared with nonselective approaches for hippocampal sclerosis. *J Neurosurg* 2006;104(01):70–78. Doi: 10.3171/jns.2006.104.1.70
- 31 Arruda F, Cendes F, Andermann F, et al. Mesial atrophy and outcome after amygdalohippocampectomy or temporal lobe removal. *Ann Neurol* 1996;40(03):446–450. Doi: 10.1002/ana.410400314
- 32 Tanriverdi T, Olivier A. Cognitive changes after unilateral cortico-amygdalohippocampectomy unilateral selective-amygdalohippocampectomy mesial temporal lobe epilepsy. *Turk Neurosurg* 2007;17(02):91–99
- 33 Carey DP, Johnstone LT. Quantifying cerebral asymmetries for language in dextrals and adextrals with random-effects meta analysis. *Front Psychol* 2014;5(1128):1128. Doi: 10.3389/fpsyg.2014.01128
- 34 Tzourio N, Crivello F, Mellet E, Nkanga-Ngila B, Mazoyer B. Functional anatomy of dominance for speech comprehension in left handers vs right handers. *Neuroimage* 1998;8(01):1–16. Doi: 10.1006/nimg.1998.0343