



Evaluation of Knowledge about Epilepsy among Undergraduate Health Students: A Cross-Sectional Study

Abdulrahman M. Alshahrani¹ Nemer Alotaibi² Mohammad Azhar Rashikh³

¹Department of Internal Medicine (Neurology), College of Medicine at Shaqra, Shaqra University, Shaqra, Saudi Arabia

²Department of Paediatrics, College of Medicine at Dawadmi, Shaqra University, Dawadmi, Saudi Arabia

³Department of Pharmacology, College of Medicine at Dawadmi, Shaqra University, Dawadmi, Saudi Arabia

Address for correspondence: Mohammad Azhar Rashikh, PhD, Department of Pharmacology, College of Medicine at Dawadmi, Shaqra University, Dawadmi, Saudi Arabia (e-mail: mrashikh@su.edu.sa).

Int J Ep 2024;10:20–27.

Abstract

Background Greater knowledge of undergraduate health students will provide quality care to patients with epilepsy. Most previous studies have targeted teachers, health care providers, and the general public. The purpose of this study was to assess the knowledge gaps of undergraduate health students on basic subjects of epilepsy, such as etiologies, symptoms, and treatment options.

Materials and Methods Our study conducted a cross-sectional survey among undergraduate health students. We collected data using a descriptive information form. The chi-square test was employed to compare knowledge of seizure between demographic and academic variables.

Results Of the total, 84.4% of students had read or heard about epilepsy. However, only 2.8% of students displayed good knowledge scores toward epilepsy. About 18.2 and 32% of students incorrectly identified that epilepsy could be due to possession by evil spirits and blood disorder, respectively. More than half of the students considered epilepsy to be nontreatable (52.1%) and a contagious disease (55.4%). Moreover, students in the advanced age group, medicine, and clinical years displayed significantly better knowledge scores than their peers ($p < 0.05$). All dental college students revealed poor scores in terms of knowledge about epilepsy's causes, symptoms, and treatment.

Conclusion Our findings concluded that most undergraduate health students demonstrated poor knowledge scores regarding epilepsy. Notably, students studying in dental college and preclinical years had a lower understanding of epilepsy than their peers. This finding suggests that introducing comprehensive epilepsy education in all health-related courses could improve patient care.

Keywords

- ▶ epilepsy
- ▶ knowledge
- ▶ Riyadh province

article published online
September 10, 2024

DOI <https://doi.org/10.1055/s-0044-1790287>.
ISSN 2213-6320.

© 2024. Indian Epilepsy Society. All rights reserved.
This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

Introduction

Epilepsy is a chronic neurological disease distressing more than 70 million people globally.^{1,2} It has been reported that approximately 80% of epileptic cases are found in the populations of developing countries.³ The prevalence of epilepsy in Saudi Arabia is projected at 6.54 per thousand inhabitants.^{4,5} One study has discovered that the occurrence of seizures is higher in Saudi Arabia than in other Arab nations.⁶

Epilepsy affects people of all ages and genders and inflicts high physical, mental, financial, and social burdens on these individuals. In particular, social stigma is a significant burden that exposes patients to unnecessary social discrimination and hinders appropriate treatment.⁴ In traditional Arab societies, for instance, women may choose to hide their seizure condition to preserve their marriage prospects, which could potentially contribute to the higher prevalence of seizures among females.⁴ Furthermore, the belief that evil spirits cause seizures is another factor that shapes the perception of seizures. As a result, individuals affected by epilepsy may turn to religious healers or assistance rather than seeking medical care.⁷

Several studies have shown a lack of awareness regarding epilepsy among the general population and health care professionals.^{8–10} There is a considerable demand for inclusive education and proper actions in response to seizures. Academic training of health care professionals, including health sciences students, is paramount and is part of the mission of the International League Against Epilepsy to understand, diagnose, and treat individuals with epilepsy in numerous settings.¹¹ Undergraduate health students may play a dynamic role in providing quality care to patients with epilepsy and are essential to any successful social health care program.¹² Higher knowledge of undergraduate health students will help prevent the adverse consequences of seizures in the community. However, only a few studies with a similar aim have been conducted among undergraduate health students. Therefore, the purpose of this study was to assess the knowledge gaps of undergraduate health students on basic subjects of epilepsy, such as causes, symptoms, and treatment options that could be useful in preparing training material for health sciences students.

Materials and Methods

Study Design and Participants

The cross-sectional study was conducted from May 16, 2023, to May 31, 2023, in Riyadh province, Saudi Arabia. The study participants included undergraduate health students, mainly from Shaqra University. The inclusion criteria were: (1) students of the undergraduate health colleges and (2) age \geq 18 years. The exclusion criteria were as follows: (1) nonundergraduate students, (2) undergraduate students from courses other than health courses, (3) students in the preparatory year, and (4) age $<$ 18 years.

Sample Size

The study assumes a prevalence of 50% of undergraduate health students having inadequate knowledge about epilepsy.¹³ We can calculate the sample size as follows:

$$n = \frac{z^2 pq}{d^2}$$

where n is the sample size, Z is the level of confidence (95% confidence interval = 1.96), p is the prevalence (0.5), $q = 1 - p$ (0.5), d is the margin of error (0.05), and the minimum sample size (n) is calculated as 384.

Data Collection Tools

The questionnaire tool used in this research is based on formerly similar studies.^{14–20} The questionnaire was translated into Arabic. The Google Forms link was used to accumulate data about epilepsy from undergraduate health students through WhatsApp, Telegram, and email. The cover letter of the questionnaire tool emphasized the study's objectives and inclusion criteria. The questionnaire was structured into two sections. Section one encompassed demographic data of undergraduate health care students and included four questions such as gender, age, college of the study, and study academic year. Section two explored the knowledge of epilepsy etiology, symptoms, and treatment. This section was comprised of eight knowledge questions, and the total scores were 16. One point was awarded for the correct option and zero for the incorrect one, except for Q3 and Q4. Five points are awarded to Q3 and five to Q4 because there are five correct answers. An individual score \leq 80% (1–12 score) and 81 to 100% (13–16 score) were considered poor and good knowledge, respectively.

Data Analysis

Data were analyzed using IBM SPSS (Statistical Package for the Social Sciences) software version 25.0.0.0 (SPSS Inc., Chicago, Illinois, United States). Descriptive statistics were used to describe the sociodemographic characteristics, presented as numbers and percentages for categorical variables. We used the chi-square test to compare demographic variables with knowledge of epilepsy. The p -values $<$ 0.05 were considered statistically significant.

Ethical Approval

The study protocol obtained ethical consent from the Institutional Ethics Committee at Shaqra University (HAPO-01-R-128) with approval number ERC_SU_20230023.

Results

Characteristics of the Respondents (– Table 1)

A total of 612 questionnaires were circulated, of which 528 (86.3% response rate) participants completed the online survey. However, 493 participants who fulfilled the inclusion criteria were included. Most participants were male (58%), studied applied medical sciences, and were in their second year.

Table 1 Characteristics of the study population (n = 493)

Variables		% (n)
Gender	Male	58 (286)
	Female	42 (207)
Age (y)	18 -20	27.2 (134)
	21–23	55.6 (274)
	24–26	17.2 (85)
College of the study	Medicine	13 (64)
	Dental	21.9 (108)
	Pharmacy	26.6 (131)
	Applied medical sciences	38.5 (190)
Year of the study	The first year	21.9 (108)
	The second year	38.9 (192)
	The third year	22.7 (112)
	The fourth year	8.7 (43)
	The fifth year	4.9 (26)
	Intern	2.4 (12)

Note: n, frequency; %, percentage.

Knowledge Scores of Study Participants (→ Fig. 1)

Only 2.8% of students had good knowledge of epilepsy etiology, symptoms, and treatment, and 97.2% had poor knowledge.

Comparison of Knowledge Scores according to Demographic Variables (→ Table 2)

Regarding good knowledge of epilepsy, advanced age group, medicine, and clinical year students displayed significantly better knowledge scores than their peers (p < 0.05). However, all the students of dental college revealed poor knowledge scores about epilepsy causes, symptoms, and treatment.

Students' Knowledge Level about Epilepsy by Different Colleges (→ Table 3)

In the present study, 84.4% of the students reported that they had heard or read about seizures. However, 15.6% of the




Fig. 1 Knowledge scores of study participants. Knowledge score ≤ 80% = poor knowledge, 81–100% score = good knowledge.

participants had never heard of epilepsy. Furthermore, a more significant proportion of medical students had heard about epilepsy than their peers. However, no statistically significant difference was observed between colleges (p > 0.05).

Of the total study participants, 35.7% of students correctly identified epilepsy as a neurological disorder and medicine students demonstrated a higher knowledge level than their peers. In contrast, 39.9, 13.8, and 9.9% of the students incorrectly identified epilepsy as a metabolic disorder, psychological disorder, and infectious disease, respectively. Moreover, a statistically significant difference in knowledge level was observed between participants from different colleges (p < 0.004).

Most of the students believed that epilepsy originated due to neurological problems (42%), genetic factors (39.7%), use of certain medications (38.9%), and brain infection (36.1%). On the other hand, 32 and 18.2% of students incorrectly identified blood disorders and possession by evil spirits are also causes of epilepsy, respectively. Furthermore, applied medical science students exhibited the lowest knowledge level with mental or emotional stress (10.5%) and hereditary (33.2%) than their peers. A statistically significant difference

Table 2 Comparison of knowledge scores according to sociodemographic variables

Variables		Poor knowledge % (n)	Good knowledge % (n)	Chi-square	p-Value
Age (y)	18 -20	99.3 (133)	0.7 (1)	7.693	0.021
	21–23	97.4 (267)	2.6 (7)		
	24–26	92.9 (79)	7.1 (6)		
College of the study	Medicine	81.2 (52)	18.8 (12)	67.902	0.000
	Dental	100 (108)	0		
	Pharmacy	100 (131)	0		
	Applied medical sciences	98.9 (188)	1.1 (2)		
Stages of the study	Preclinical year students (1st, 2nd, and 3rd year)	98.5 (406)	1.5 (6)	17.395	0.000
	Clinical year students (4th, 5th, and intern)	90.1 (73)	9.9 (8)		

Note: n, frequency; %, percentage; p-values were generated using the Persian chi-square test for comparing two groups. Significant at p-value ≤ 0.05. An individual score ≤ 80% (1–12 score) and 81–100% (13–16 score) were considered poor and good knowledge, respectively.

Table 3 Students knowledge level of epilepsy by different colleges

Variables	Total % (n)	Medicine % (n)	Dental % (n)	Pharmacy % (n)	Applied medical sciences % (n)	p-Value
Q1. Have you heard or read about epilepsy?	Yes	84.4 (416)	92.2 (59)	85.2 (92)	84.7 (111)	0.202
	No	15.6 (77)	7.8 (5)	14.8 (16)	15.3 (20)	
Q2. What is epilepsy?	Neurological disease ^a	35.7 (176)	60.9 (39)	27.8 (30)	32.8 (43)	0.004
	Metabolic disease	39.9 (197)	26.5 (17)	46.3 (50)	39.7 (52)	
	Psychological disease	13.8 (68)	7.8 (5)	14.8 (16)	12.2 (16)	
	Infectious disease	9.9 (49)	4.7 (3)	10.2 (11)	15.3 (20)	
	I do not know	0.6 (3)	0	0.9 (1)	0	
Q3. What are the etiologies of epilepsy?	Mental or emotional stress ^a	14.6 (72)	28.1 (18)	13.9 (15)	14.5 (19)	0.008
	Genetic factors ^a	39.7 (196)	43.7 (28)	48.1 (52)	40.5 (53)	0.069
	Neurological problems ^a	42.0 (207)	62.5 (40)	34.3 (37)	41.9 (55)	0.003
	Brain infection ^a	36.1 (178)	32.8 (21)	43.5 (47)	33.6 (44)	0.336
	Use of certain medications ^a	38.9 (192)	43.7 (28)	44.4 (48)	35.1 (46)	0.368
	Blood disorder	32.0 (158)	20.3 (13)	36.1 (39)	30.5 (40)	0.131
	Possession by evil spirits	18.2 (90)	14.1 (9)	20.4 (22)	17.5 (23)	0.757
	Loss of consciousness ^a	20.5 (101)	45.3 (29)	10.2 (11)	19.1 (25)	0.000
Q4. What are the symptoms of epilepsy?	Secretion from the mouth ^a	43.2 (213)	54.7 (35)	37.9 (41)	38.2 (50)	0.089
	Tongue biting ^a	42.8 (211)	51.5 (33)	33.3 (36)	48.1 (63)	0.056
	Convulsion of the body ^a	44.8 (221)	53.1 (34)	39.8 (43)	42.7 (56)	0.351
	Urine incontinence ^a	17.8 (88)	40.6 (26)	14.8 (16)	9.9 (13)	0.000
	Medical treatment ^a	25.3 (125)	57.8 (37)	9.2 (10)	23.7 (31)	0.000
Q5. What are the usual treatments for epilepsy?	Herbal medicine	38.9 (192)	26.5 (17)	50.9 (55)	42.0 (55)	0.000
	Cauterization	23.9 (118)	9.4 (6)	26.8 (29)	24.4 (32)	
	Surgical intervention ^a	6.9 (34)	1.5 (1)	7.4 (8)	6.9 (9)	
	No need for treatment	2.6 (13)	0	5.5 (6)	2.3 (3)	
	I do not know	2.2 (11)	4.7 (3)	0	0.7 (1)	
Q6. Can epilepsy be treated? Yes ^a	47.9 (236)	57.8 (37)	48.1 (52)	43.5 (57)	47.4 (90)	0.314
Q7. Is epilepsy a long duration treatment course? Yes ^a	49.5 (244)	68.7 (44)	46.3 (50)	43.5 (57)	48.9 (93)	0.008
Q8. Do you think epilepsy is contagious? No ^a	55.4 (273)	67.2 (43)	50.0 (54)	51.9 (68)	56.8 (108)	0.128

Note: n, frequency; %, percentage; p-values were generated using the Persian chi-square test for comparing two groups. Significant at p-value ≤ 0.05.

^aDenotes correct option.

in knowledge level was observed between the participants of different colleges ($p < 0.05$), only with neurological problems and emotional stress.

Note that 44.8% of students claimed that convulsion of the body was the most common symptom of epilepsy, followed by secretion from the mouth (43.2%) and tongue biting (42.8%).

Note that 25.3% of the students believed that medical treatment was a mainstay approach for epilepsy, followed by surgical intervention (6.9%). However, 38.9% of the students incorrectly identified that epilepsy can be treated by herbal medicine, followed by cauterization (23.9%). Furthermore, dental students exhibited the lowest knowledge of medical treatment (9.2%) compared with their peers. However, medical students demonstrated the lowest knowledge level with surgical intervention (1.5%). A statistically significant difference was observed between participants from different colleges ($p < 0.001$).

Nearly half of the students acknowledged that epilepsy can be treated (47.9%) and believed in a long-duration treatment course (49.5%). More than half of the students (55.4%) correctly affirmed that epilepsy is not contagious. Additionally, medicine students displayed higher knowledge levels than their peers.

Students' Knowledge Level of Epilepsy by Different Study Stages (–Table 4)

Over two-thirds of clinical year students correctly identified neurological problems as the leading cause of epilepsy. However, 58% of the basic year students correctly identified genetic factors as a primary cause of epilepsy. In contrast, a higher number of basic year students incorrectly believed that blood disorders (37.1% vs. 18.5%) and possession by evil spirits (19.4% vs. 12.3%) are a cause of epilepsy than clinical year students. However, a statistically significant difference in knowledge level was observed between participants of different study stages ($p < 0.05$) only with the blood disorders, neurological problems, and emotional stress.

Basic year students displayed significantly lower knowledge levels than clinical year students toward epilepsy symptoms such as loss of consciousness (14.6% vs. 50.6%; $p < 0.001$), convulsion of the body (40.8% vs. 69.1%, $p < 0.001$), and urine incontinence (14.3% vs. 34.8%, $p < 0.001$). However, more clinical students correctly identified tongue biting as a symptom of epilepsy than basic year students (64.2% vs. 39.3%, $p < 0.001$).

Clinical year students correctly identified medical treatment as a mainstay approach for epilepsy and exhibited significantly higher knowledge levels than basic year students (61.7% vs. 18.2%, $p < 0.001$). Additionally, clinical year students demonstrated better knowledge than basic year students about epilepsy, which is treatable (66.7% vs. 44.2%, $p < 0.001$), long-duration treatment course (65.4% vs. 46.4%, $p < 0.01$), and not contagious (66.7% vs. 53.1%, $p < 0.05$), and a statistically significant difference in knowledge level was observed between participants of different study stages ($p < 0.05$).

Discussion

The incidence of seizures in Saudi Arabia has been reported as substantial, highlighting the urgent need for comprehensive education on seizures.⁶ High awareness and knowledge about seizures are of the utmost importance in preventing many harmful measures that could affect the community.²¹ However, there is a lack of knowledge about how to deal with someone having a seizure. Therefore, this study aimed to explore undergraduate health students' awareness and knowledge of epilepsy, including causes, symptoms, and treatment options.

In our study, 84.4% of students had heard or read about epilepsy. This figure is comparable to the findings of the previous study of Saudi Arabia²² (80%) and Sudan²³ (85.4%). However, we found that our figure was lower than Germany¹⁹ (96.7%), China²⁴ (94.7%), and India²⁵ (92.5%). On the other hand, the percentage of clinical students who reported awareness of epilepsy was significantly higher than that of preclinical students (93.8% vs. 82.5%, $p < 0.01$), which is consistent with previous studies in Saudi Arabia.^{26,27}

Regarding the overall knowledge of seizure etiology, symptoms, and treatment, only 3.2% of the students demonstrated good knowledge scores, much lower than studies conducted in China^{24,28} and India.²⁹ This difference is primarily attributed to study setting variations, population, and educational curriculum.

Regarding the students' knowledge scores, our study displayed a significant difference in age group, consistent with a study conducted in Turkey¹⁶ and Croatian.³⁰ Similarly, our study observed a significant difference in knowledge scores between the students of different colleges and study stages. Moreover, we found a high percentage of good knowledge in medicine and clinical year students, which matched the Palestine study.^{31,32} The possible reason for the better knowledge of medicine students is that they are trained with new teaching models, such as problem-based learning methods, and spend more time with patients in clinical settings than other student groups.^{20,33}

In the current study, around 35.7% of the students believed that epilepsy is a neurological disease. This percentage is much lower than that of previous studies of Saudi Arabia 86.5³⁴ and 82.7%.³⁵ The difference might be due to the study on a different population. Moreover, our findings have revealed that the most common causes of epilepsy were neurological problems (42%), hereditary (39.7%), and use of certain medications (38.9%). However, medical students from Sudan and previous studies in Saudi Arabia reported that brain diseases, genetic factors, and mental or emotional stress disorders were the most common causes of epilepsy.^{8,22,25} Contrarily, epilepsy has been supposed to be a disease that results from evil spirits.⁹ In the present study, 18.2% of students believed that the evil spirit could cause epilepsy. This percentage is similar to a previous study of Riyadh³⁶ (15%) and Sudan²³ (15.7%). However, a higher percentage was reported in Jordan³³ (31.5%), Libya³⁷ (37.5%), and Kuwait³⁸ (24.6%). These findings suggest an

Table 4 Students' knowledge level of epilepsy by different study stages

Variables	Total % (n) 100 (493)	Preclinical year students (1st, 2nd, and 3rd year) % (n) 83.6 (412)	Clinical year students (4th, 5th, and interns) % (n) 16.4 (81)	p-Value
Q1. Have you heard or read about epilepsy?	Yes	84.4 (416)	93.8 (76)	0.010
	No	15.6 (77)	6.2 (5)	
Q2. What is epilepsy?	Neurological disease ^a	35.7 (176)	29.4 (121)	0.000
	Metabolic disease	39.9 (197)	44.4 (183)	
	Psychological disease	13.8 (68)	14.1 (58)	
	Infectious disease	9.9 (49)	11.4 (47)	
	I do not know	0.6 (3)	0.6 (3)	
	Mental or emotional stress ^a	14.6 (72)	12.4 (51)	
Q3. What are the etiologies of epilepsy?	Neurological problems ^a	40.5 (200)	34.9 (144)	0.000
	Genetic factors ^a	56.2 (277)	58.0 (239)	
	Use of certain medication ^a	39.7 (196)	38.8 (160)	
	Brain infection ^a	29.0 (143)	30.3 (125)	
	Blood disorder	34.1 (168)	37.1 (153)	
	Possession by evil spirits	18.3 (90)	19.4 (80)	
	Loss of consciousness ^a	20.5 (101)	14.6 (60)	
	Secretion from the mouth ^a	43.2 (213)	42.0 (173)	
Q4. What are the symptoms of epilepsy?	Tongue biting ^a	43.4 (214)	39.3 (162)	0.000
	Convulsion of the body ^a	45.4 (224)	40.8 (168)	
	Urine incontinence ^a	17.8 (88)	14.3 (59)	
	Medical treatment ^a	25.3 (125)	18.2 (75)	
	Herbal medicine	38.9 (192)	43.4 (179)	
	Cauterization	23.9 (118)	26.9 (111)	
	Surgical intervention ^a	6.9 (34)	6.8 (28)	
	No need for treatment	2.6 (13)	2.7 (11)	
Q5. What are the usual treatments for epilepsy?	I do not know	2.2 (11)	1.9 (8)	0.000
	Medical treatment ^a	25.3 (125)	18.2 (75)	
Q6. Can epilepsy be treated? Yes ^a	Herbal medicine	38.9 (192)	43.4 (179)	0.000
	Cauterization	23.9 (118)	26.9 (111)	
Q7. Is epilepsy a long duration treatment course? Yes ^a	Surgical intervention ^a	6.9 (34)	6.8 (28)	0.000
	No need for treatment	2.6 (13)	2.7 (11)	
Q8. Do you think epilepsy is contagious? No ^a	I do not know	2.2 (11)	1.9 (8)	0.025
	Medical treatment ^a	25.3 (125)	18.2 (75)	
Q6. Can epilepsy be treated? Yes ^a	Herbal medicine	38.9 (192)	43.4 (179)	0.000
	Cauterization	23.9 (118)	26.9 (111)	
Q7. Is epilepsy a long duration treatment course? Yes ^a	Surgical intervention ^a	6.9 (34)	6.8 (28)	0.002
	No need for treatment	2.6 (13)	2.7 (11)	
Q8. Do you think epilepsy is contagious? No ^a	I do not know	2.2 (11)	1.9 (8)	0.025
	Medical treatment ^a	25.3 (125)	18.2 (75)	

Note: n, frequency; %, percentage; p-values were generated using the Persian chi-square test for comparing two groups. Significant at p-value ≤ 0.05.

^aDenotes correct option.

improvement in the knowledge and awareness among undergraduate health students.

In our study, 44.8% of the participants indicated that convulsions are the foremost manifestations of epilepsy. The present finding result is much lower than the previous study of Taif²¹ (97%), Riyadh²⁵ (92.3%), and China²⁷ (97%). However, the clinical year students who knew the symptoms and causes of epilepsy displayed significantly higher knowledge than preclinical year students, which is consistent with previous studies in Saudi Arabia.²⁶

The seizures of most people can be adequately controlled with appropriate medication regimens. However, 20 to 30% of people with epilepsy are refractory to all forms of medical therapy and may require surgical interventions.³⁹ Note that 25.3% of students believed that medical treatment is the mainstay approach for epilepsy. This finding is similar to Libya³⁷ (31.8%) but lower than the Taif²¹ (77.7%). On the other hand, only 6.9% of students think that surgical intervention could be a treatment choice for epilepsy. This percentage is much lower than the previous study of Saudi Arabia⁸ (44%), Turkey⁴⁰ (25.2%), and Slovenia⁴¹ (74.5%). Moreover, nearly half of the students (47.9%) acknowledged that epilepsy can be treated. This finding is higher than the previous study of Riyadh²⁵ (36.6%) and Southern China⁴² (37.7%) but lower than Sudan⁴³ (90.4%). Of note, a significantly higher number of clinical students believed that epilepsy is a treatable disorder compared with preclinical students, which is consistent with the Palestine study⁴⁴ and a previous study of Saudi Arabia.²⁶

Additionally, 49.5% of students believed that epilepsy treatment is typically a long-duration process, consistent with the Sudan study conducted on first-year medical students.²² Moreover, 55.4% of the students correctly affirmed that epilepsy is not contagious. This percentage is much lower than the previous study of Makkah³⁵ (94.9%) and Sudan²² (98.3%). However, our study result is better than that of the Aseer region of Saudi Arabia; 95.8% of the participants acknowledged that epilepsy is a contagious illness.⁴⁵ The difference might be due to the study conducted on different populations.

The present study has certain limitations. First, our study used convenience sampling, which may only be representative of some populations of undergraduate health college students. Second, this study utilizes a cross-sectional design, which could disclose the associations between variables but does not institute causal relationships. Third, this study did not include questions such as paroxysmal events with and without impaired awareness, brief events, and stereotypes.

Conclusion

The results of our study revealed that most undergraduate health students in Riyadh province had inadequate knowledge about epilepsy, evidenced by the deficient percentage of good knowledge of epilepsy. Furthermore, it was observed that the advanced age group, students studying in medical college, and clinical years, had a better understanding of epilepsy than younger age, other undergraduate health

students, and preclinical students. The study recommends introducing epilepsy education from the primary training stage in all health-related courses, which is crucial to enhancing students' knowledge of epilepsy. It emphasizes the importance of increasing epilepsy awareness among the general public, orienting medical students to emergency neurological conditions at the early stages of their education, and educating epilepsy in applied medical sciences and pharmacy. Further comparison study on the knowledge of seizure between undergraduates and postgraduate health students is recommended.

Availability of Data and Materials

All the data supporting this study's findings are available in the manuscript.

Authors' Contributions

A.M.A.: Supervision, conceptualization, designed the study and questionnaire, collected the data, and critically reviewed the manuscript; N.A.: Designed the study and critically reviewed and edited the manuscript. M.A.R.: Collected the data, analyzed the data, interpreted the results, and prepared the original draft of the manuscript.

Funding

None.

Conflict of Interest

None declared.

Acknowledgment

The authors are incredibly thankful to the Deanship of Scientific Research at Shaqra University, Kingdom of Saudi Arabia, for supporting this study.

References

- 1 Thijs RD, Surges R, O'Brien TJ, Sander JW. Epilepsy in adults. *Lancet* 2019;393(10172):689–701
- 2 Fiest KM, Sauro KM, Wiebe S, et al. Prevalence and incidence of epilepsy: a systematic review and meta-analysis of international studies. *Neurology* 2017;88(03):296–303
- 3 Birbeck GL. Epilepsy care in developing countries: part I of II. *Epilepsy Curr* 2010;10(04):75–79
- 4 Al Rajeh S, Awada A, Bademosi O, Ogunniyi A. The prevalence of epilepsy and other seizure disorders in an Arab population: a community-based study. *Seizure* 2001;10(06):410–414
- 5 MOH Health Days 2020—International Epilepsy Day. Accessed May 18, 2024 at: <https://www.moh.gov.sa/en/HealthAwareness/healthDay/2020/Pages/HealthDay-2020-02-10.aspx>
- 6 Benamer HT, Grosset DG. A systematic review of the epidemiology of epilepsy in Arab countries. *Epilepsia* 2009;50(10):2301–2304
- 7 Obeid T, Abulaban A, Al-Ghatani F, Al-Malki AR, Al-Ghamdi A. Possession by 'Jinn' as a cause of epilepsy (Saraa): a study from Saudi Arabia. *Seizure* 2012;21(04):245–249
- 8 Alaqeel A, Alebdi F, Sabbagh AJ. Epilepsy: what do health-care professionals in Riyadh know? *Epilepsy Behav* 2013;29(01):234–237
- 9 Chomba EN, Haworth A, Atadzhanov M, Mbewe E, Birbeck GL. Zambian health care workers' knowledge, attitudes, beliefs, and

- practices regarding epilepsy. *Epilepsy Behav* 2007;10(01):111–119
- 10 Vancini RL, Benedito-Silva AA, Sousa BS, et al. Knowledge about epilepsy among health professionals: a cross-sectional survey in Sao Paulo, Brazil. *BMJ Open* 2012;2(02):e000919
 - 11 O'Hara KA. First aid for seizures: the importance of education and appropriate response. *J Child Neurol* 2007;22(05):305–375
 - 12 Blümcke I, Arzimanoglou A, Beniczky S, Wiebe S. Roadmap for a competency based educational curriculum in epileptology: report of the epilepsy education task force of the International League Against Epilepsy. *Epileptic Disord* 2019;21(02):129–140
 - 13 Elhassan MA, Alemairy AA, Amara ZM, Hamadelneel AA, Mohamed AH, Elaimeri AA. Epilepsy: knowledge, attitude, and practice among secondary school teachers in Khartoum State. *Neurol Ther* 2017;6(02):225–235
 - 14 Ozer Z, Turan GB, Kose S. Are nursing students ready to provide quality care to patients with epilepsy? A comparative cross sectional study. *Am J Nurs Stud* 2023;3(01):1018
 - 15 Erdem RZ, Yilmaz C, Uygun LA. Evaluation of the attitude and knowledge level of dentistry undergraduate and specialist students towards epilepsy patients. *Epilepsy Behav* 2023;149:109497
 - 16 Aksoy M, Büyükbayram Z. Evaluating the knowledge of and attitudes toward epilepsy among Turkish undergraduate nursing students: a cross-sectional study. *Epilepsy Behav* 2022;126:108477
 - 17 Shawahna R, Jaber M. Assessing knowledge and attitudes of Palestinian undergraduate nursing students toward epilepsy and patients with epilepsy: a cross-sectional study. *Epilepsy Behav* 2020;102:106811
 - 18 Hakami T, Mahfouz M, Najmi H, et al. Knowledge of and attitude towards epilepsy among university students in Saudi Arabia: misconceptions of the next generation. *Epilepsy Behav Rep* 2021;16:100450
 - 19 Mewes S, Jeschke S, Bertsche T, Neining MP, Bertsche A. Knowledge of and attitudes towards epilepsy among first- and second-year students at a German university. *Epilepsy Behav* 2020;112:107490
 - 20 Alhalaiqa F, Al Omari O, Batiha AM, et al. Knowledge and attitudes of Jordanian university students toward epilepsy: a cross-sectional comparison study. *Int Q Community Health Educ* 2018;38(02):75–82
 - 21 England MJ, Austin JK, Beck V, Escoffery C, Hesdorffer DC. Erasing epilepsy stigma: eight key messages. *Health Promot Pract* 2014;15(03):313–318
 - 22 Kabel AM, Algethami SA, Algethami BS, Alzahrani AS, Almutairi SK, Almutairi AS. Knowledge, perceptions, and attitudes of students of health-related science colleges towards epilepsy in Taif, Saudi Arabia. *J Family Med Prim Care* 2020;9(05):2394–2399
 - 23 Ibrahim ME, Hasabo EA, Hsabo EA, Alhadi AS. Awareness and attitude of first-year medical students toward epilepsy in University of Khartoum, Sudan. *Brain Behav* 2022;12(01):e2461
 - 24 Zhao T, Zhang X, Cui X, et al. Awareness, attitudes and first aid knowledge of epilepsy among university students - a cross-sectional study in Henan Province, China. *Epilepsy Res* 2024;201:107315
 - 25 Panda SB, Prabhu K, Rao S, et al. Evaluation of knowledge of and attitudes toward epilepsy among the health science students of Manipal University. *Epilepsy Behav* 2011;20(03):447–449
 - 26 Bin Abdulrahman KA, Alharbi MG, Alzeer AM. Awareness and attitudes toward epilepsy among medical students and interns in Riyadh, Saudi Arabia. *Prim Health Care Res Dev* 2022;23:e70
 - 27 Alomar S, Kadi M, Alabbas D, et al. Awareness and attitudes toward epilepsy among medical and allied healthcare students - a survey study in a teaching hospital in Jeddah. *Epilepsy Behav* 2020;102:106815
 - 28 Tu J, Sun Y, Tian D, et al. Impact of metacognition on attitudes toward epilepsy in medical students. *Epilepsy Behav* 2023;143:109243
 - 29 Murthy MKS, Govindappa L, Marimuthu P, Dasgupta M. Exploring knowledge, attitude, and practices in relation to epilepsy among undergraduates for effective health promotion: Initial evaluation. *J Educ Health Promot* 2019;8:122
 - 30 Žuvela T, Filipović-Grčić B, Rušić D, et al. Knowledge and attitudes towards epilepsy of Croatian general student population and biomedical students: a cross-sectional study. *Healthcare (Basel)* 2023;11(18):2550
 - 31 Shawahna R, Jaber M, Maqboul I, et al. Are medical students adequately prepared to provide quality care for patients with epilepsy? A cross-sectional study of their knowledge and attitude. *Epilepsy Behav* 2021;120:107976
 - 32 Vodougnon C, Gérard D, Bruand PE, et al. Knowledge, attitudes, and practices of health sciences students regarding epilepsy at the end of their curriculum in Benin. *Epilepsy Behav* 2019;92:165–170
 - 33 Hijazeen JK, Abu-Helalah MA, Alshraideh HA, et al. Knowledge, attitudes, and beliefs about epilepsy and their predictors among university students in Jordan. *Epilepsy Behav* 2014;41:238–243
 - 34 Alsulami H, Alhadhrani S, Alshareef B, Alqurashi R, Alzahrani A, Alkhotani A. Assessment of knowledge and attitude regarding epilepsy and seizure first aid among male teachers in Mecca region, Saudi Arabia: a cross-sectional study. *Cureus* 2022;14(10):e30945
 - 35 Alkhotani AM, Alkhotani AM. Epilepsy stigma among university students in Makkah: a cross-sectional study. *Neuropsychiatr Dis Treat* 2022;18:1047–1056
 - 36 Muthaffar OY, Jan MM. Public awareness and attitudes toward epilepsy in Saudi Arabia is improving. *Neurosciences (Riyadh)* 2014;19(02):124–126
 - 37 Alhagamhmad MH, Shembesh NM. Investigating the awareness, behavior, and attitude toward epilepsy among university students in Benghazi, Libya. *Epilepsy Behav* 2018;83:22–27
 - 38 Al-Rashed H, Al-Yahya D, Al-Kandari A, Shehab A, Al-Sabah R, Al-Taiar A. Knowledge of, perceptions of, and attitudes toward epilepsy among university students in Kuwait. *Epilepsy Behav* 2009;14(02):367–371
 - 39 Kobau R, Zahran H, Thurman DJ, et al. Centers for Disease Control and Prevention (CDC) Epilepsy surveillance among adults—19 States, Behavioral Risk Factor Surveillance System, 2005. *MMWR Surveill Summ* 2008;57(06):1–20
 - 40 Kartal A. Knowledge of, perceptions of, attitudes and practices regarding epilepsy among medical students in Turkey. *Epilepsy Behav* 2016;58:115–118
 - 41 Zupan G, Lorber B. Knowledge and awareness of epilepsy surgery among medical students. *J Epilepsy Res* 2017;7(01):50–53
 - 42 Yang K, He Y, Xiao B, Wang J, Feng L. Knowledge, attitudes and practice towards epilepsy among medical staff in Southern China: does the level of hospitals make a difference? *Seizure* 2019;69:221–227
 - 43 Mustafa MN, Wedaa Alla LG, Ahmad TK. Knowledge and attitude toward epilepsy among the final and semi-final years medical students at University of Bahri, Sudan. *Epilepsia Open* 2022;7(01):144–150
 - 44 Abuawad M, Rjoub A, Abu Zahra W. Assessment of epilepsy awareness, knowledge, and attitudes among preclinical and clinical medical students in the West Bank of Palestine. *Epilepsy Behav* 2023;145:109317
 - 45 Abdel Ghaffar NF, Asiri RN, Al-Eitan LN, et al. Improving public stigma, sociocultural beliefs, and social identity for people with epilepsy in the Aseer region of Saudi Arabia. *Epilepsy Behav Rep* 2021;16:100442