


**Figure 1** A healthy 59-year-old man underwent colonoscopy because of a positive fecal blood test. A yellowish-white lobulated polyp, 15-mm in size with a “baby’s hand”-like morphology, was seen. Endoscopic polypectomy for suspected leiomyoma was performed without complication.



**Figure 2** The edematous stroma was filled with enlarged serpiginous veins and arterioles, leading to a diagnosis of angiodysplasia. Angiodysplasia is one of the major causes of lower gastrointestinal bleeding, often encountered during emergency colonoscopy. The typical endoscopic appearance is often reported to be a

slightly elevated reddish lesion, reflecting the dilated, tortuous veins in the submucosa. Lesions having a polypoid morphology are extremely rare. This case suggests us the possibility of angiodysplasia as a polypoid lesion, and the need for care when performing biopsy or endoscopic polypectomy.

**N. Kakushima<sup>1</sup>, M. Fujishiro<sup>1</sup>, N. Yahagi<sup>1</sup>, M. Oka<sup>1</sup>, K. Kobayashi<sup>1</sup>, T. Hashimoto<sup>1</sup>, H. Miyoshi<sup>1</sup>, Y. Moriyama<sup>1</sup>, S. Enomoto<sup>2</sup>, M. Iguchi<sup>2</sup>, M. Ichinose<sup>2</sup>, M. Omata<sup>1</sup>**

<sup>1</sup> Department of Gastroenterology, Faculty of Medicine, University of Tokyo, Tokyo, Japan

<sup>2</sup> Second Department of Internal Medicine, Wakayama Medical College, Wakayama, Japan

## Corresponding Author

**N. Yahagi, M.D.**

Department of Gastroenterology  
Faculty of Medicine  
University of Tokyo  
7-3-1 Hongo  
Bunkyo-ku  
Tokyo  
Japan

Fax: +81-3-5800-8806

E-mail: Yahagi-tky@umin.ac.jp