

Comparison of Bone Mineral Density, T-Scores and Serum Zinc between Diabetic and Non Diabetic Postmenopausal Women with Osteoporosis

Priyanka R Siddapur, Anuradha B Patil, Varsha S Borde¹

Departments of Biochemistry and ¹Physiology, Jawaharlal Nehru Medical College, KLE University, Belgaum, Karnataka, India

Address for correspondence: Dr. Priyanka Ramappa Siddapur, E-mail: priyankasiddapur@gmail.com

ABSTRACT

Context: Postmenopausal osteoporosis is a public health problem. Diabetics are at increased risk of osteoporosis-related fractures. Zinc (Zn) has a role in collagen metabolism, and its levels are altered in diabetes.

Aims: The aim was to compare bone mineral density (BMD), T-score and serum Zn between diabetic and nondiabetic postmenopausal women with osteoporosis to see if they influence increased fracture risk in diabetes.

Settings and Design: It is a cross-sectional study conducted at Department of Biochemistry, Jawaharlal Nehru Medical College, Belgaum.

Materials and Methods: Thirty type 2 diabetic and 30 age-matched (aged 45-75 years) nondiabetic Dual energy X-ray absorptiometry (DEXA) confirmed postmenopausal osteoporotics were included from January 2011 to March 2012. Serum Zn was analyzed by atomic absorption spectrophotometry.

Statistical Analysis Used: Mean and standard deviation of the parameters of the two groups were computed and compared by unpaired Student's *t*-test. Relationship between variables was measured by Karl Pearson's correlation co-efficient. A statistical significance is set at 5% level of significance ($P < 0.05$).

Results: T-score was significantly higher in diabetics compared with nondiabetics (-2.84 ± 0.42 vs. -3.22 ± 0.74) $P < 0.05$. BMD and serum Zn of diabetics showed a significant positive correlation with body mass index (BMI).

Conclusions: Type 2 diabetic postmenopausal osteoporotics have a higher T-score than the nondiabetics. High BMI in type-2 diabetes mellitus (T2DM) may contribute to high BMD and may be a protective factor against zincuria. Increased fracture risk in T2DM could be due to other factors like poor bone quality due to hyperglycemia rather than BMD. Strict glycemic control is of paramount importance.

Key words: Bone mineral density, diabetes, menopause, osteoporosis, serum zinc, T-score

INTRODUCTION

The word “osteoporosis” literally means “porous bone.”^[1] By definition, Osteoporosis is a disease characterized by low bone mass and deterioration of bone microarchitecture with a consequent increase in bone fragility and hence fracture risk.^[2] It has been called as a silent epidemic as bone

loss is silent and progressive often asymptomatic until the first fracture occurs.^[1] Dual X-ray absorptiometry has become the standard for measurement of low bone mineral density (BMD) associated with osteoporosis. The definition of Osteoporosis according to World Health Organization (WHO) is a fall in BMD 2.5 standard deviations (SDs) below the mean for young healthy adults of the same gender which is also referred to as a T-score of -2.5 .^[3]

In their lifetime 30-50% of women and 15-30% of men are at risk for osteoporosis and its related fractures all over the world.^[4] Conservative estimates suggest that around 25 million of Indian population would be affected by 2015.^[5] Women are predominantly affected by this disease.^[1] Menopausal period is

Access this article online	
Quick Response Code: 	Website: www.jlponline.org
	DOI: 10.4103/0974-2727.151681

associated with increased bone loss due to estrogen deficiency and ageing.^[3] Although osteoporosis has not been traditionally listed as a complication of diabetes, patients with Diabetes are at increased risk for this disease.^[6] Osteoclasts, the bone resorbing cells, require glucose for energy. Hence, enhanced osteoclast activity may be seen in diabetes. Hyperglycemia also leads to glycation of various bone proteins including type I collagen, which may impair bone quality.^[7]

Zinc (Zn) an essential trace element is necessary for normal collagen synthesis and mineralization of bone.^[8] It acts as a cofactor for alkaline phosphatase, bone forming metalloenzyme.^[9] Reduced serum or plasma Zn concentrations and increased urinary Zn excretion have been reported in women with osteoporosis.^[10] Zn also plays an important role in insulin action with several investigators showing the perturbation of Zn metabolism in diabetics.^[11] It has not been clearly elucidated whether Zn deficiency is a consequence of hyperglycemia or whether Zn deficiency contributes to the pathogenesis of diabetes.^[12]

Studies have shown increased risk of fractures and bone abnormalities in diabetes mellitus patients.^[6,7] Hence, this study is undertaken to compare the BMD, T-scores and serum Zn between diabetic and nondiabetic postmenopausal women with osteoporosis to see if they are altered in diabetes contributing to increased fracture risk in diabetes.

MATERIALS AND METHODS

The study is cross-sectional in design. It was conducted in Department of Biochemistry, KLES Dr. Prabhakar Kore Hospital and Medical Research Centre, Belgaum between January 2011 to March 2012. It comprised of 30 diabetic and age-matched 30 nondiabetic clinically, diagnostically confirmed cases of Postmenopausal Osteoporosis attending the orthopedic unit of KLES Dr. Prabhakar Kore Hospital and Medical Research Centre, Belgaum. Permission to conduct the study was obtained from the Institutional ethics committee on human subject's research of Jawaharlal Nehru medical college, Belgaum. All the cases were evaluated and selected by simple random technique after fulfilling selection criteria. The cases of Osteoporosis reported to the Department of Orthopedics, KLES Dr. Prabhakar Kore Hospital and Medical Research Center were screened. After finding the suitability as per selection criteria, they were requested to participate in the study and briefed about the nature of the study and interventions used.

A written informed consent was obtained. The consented patients were enrolled in the study. Further descriptive data of the participants like name, age, sex, detailed history, were obtained by interviewing the participants and were recorded on a predesigned and pretested proforma.

The women were characterized as postmenopausal if they had not menstruated for at least 12 months. The study population belonged to the age group of 45-75 years. Dual energy X-ray absorptiometry (DEXA) scan 'T' score < -2.5 according to the WHO criteria^[3] were considered to be osteoporotic. The diabetic group comprised of known cases of diabetes with at least 1 month duration. Women were identified as nondiabetic based on fasting plasma glucose levels < 100 mg/dL according to the WHO criteria for diagnosis of Diabetes.^[13] Women with surgical menopause, hypertensives, renal disorders, smokers, alcoholics and those being treated with biphosphonates, calcitonin, anabolic steroids, hormone replacement therapy, calcium, Vitamin D previously at any point since the beginning of menopause were excluded from the study. Diabetic women taking insulin preparations containing Zn were excluded.

It was intended to include both Type 1 and Type 2 diabetic women. However, the diabetic women who met the inclusion and exclusion criteria in this study belonged to type 2 DM category (T2DM).

According to previous hospital records, the number of postmenopausal DEXA confirmed osteoporotic cases from Orthopedic department in KLE hospital has been around 80 cases/year. The number of Diabetic patients with DEXA confirmed Osteoporosis was unknown. For the study to be statistically significant minimum sample size of 30 Diabetic and 30 nondiabetic postmenopausal women with osteoporosis were taken.

Bone mineral density assessment

Bone mineral density of the patient was determined using whole body densitometer, DEXA Scan (GE Healthcare Lunar prodigy advance, scanner serial no. PA + 302343, software version— ENCORE 2008 version 12.2, Germany) in KLE hospital.

Bone mineral density values and T-scores of L1-L4 Lumbar spine were noted. Height and weight were measured at the time of DEXA measurement and body mass index (BMI) was calculated as the weight divided by the square of the height (kg/m^2).

Collection and storage of the blood sample

Five milliliter of blood was collected in the fasting state in the morning before breakfast in a plain nonvacuum tube and was allowed to stand in room temperature till clot was formed. Serum was separated within 1 h of venipuncture by centrifuging the tubes at 3,000 r.p.m for 10 min. Serum was pipetted out into sterile Eppendorf tubes. Serum glucose was estimated immediately by Trinder's Method^[14] using ERBA reagent kit in ERBA Chem 5 semiautoanalyzer in KLEs Charitable Hospital laboratory. All samples were stored in nonvacuum sterile tubes at -20°C till further analysis.

Zinc estimation

Serum Zn estimation of the stored samples was done using Atomic Absorption Spectrophotometer (AAS) Perkin Elmer Analyst 300, University Science Instrument Centre, Shivaji University, Kolhapur. Serum sample was diluted to 1:10 or 1:5 with deionized water. The dilution ratio was adjusted so that the concentrations fall within a suitable absorbance range.

Standard

Zinc stock solution provided by the manufacturer was diluted, and appropriate standards were prepared.

Blank

Deionized water was used as blank solution.

Analysis

Instrument was set in standard condition for the Zn analysis. Blank was aspirated first followed by a suitable standard and then the sample. Zn standard was read after every five samples. The readings were recorded in parts per million (p.p.m) and then converted to $\mu\text{g}/\text{dL}$.

Statistical analysis

The data were analyzed by SPSS Inc. Released 2007. SPSS for Windows, Version 16.0. Chicago, SPSS Inc.

Mean and SD of BMD, T-score and serum Zn of the two groups were computed and compared by unpaired Student's *t*-test. Results are expressed as Mean \pm SD. Relationship between variables was measured by Karl Pearson's correlation co-efficient. A statistical significance is set at 5% level of significance ($P < 0.05$).

RESULTS

Bone mineral density values and T-scores in 30 type 2 diabetic and 30 age-matched nondiabetic postmenopausal women with osteoporosis were compared.

The data obtained from the study was compiled, tabulated and subjected to statistical analysis. As the two groups were age-matched, there was no significant difference in the mean age in both the groups ($P = 0.9669$). Diabetics had a significantly higher mean weight and BMI than nondiabetic participants. Diabetics had a higher mean BMD than nondiabetics, but it was not statistically significant ($P = 0.0661$). However, the T-scores were significantly higher in case of diabetic women compared to nondiabetics ($P = 0.0177$). There was no significant difference in the mean serum Zn levels between diabetics and nondiabetic postmenopausal women with osteoporosis ($P = 0.1042$) [Table 1].

BMD levels showed a significant positive correlation with serum Zn in both diabetic $P < 0.001$ and nondiabetic participants $P < 0.002$ [Table 2, Graphs 1 and 2]. BMI of diabetics showed a significant positive correlation with Zn ($P = 0.021$) and BMD ($P = 0.019$) [Table 2, Graphs 3 and 4].

DISCUSSION

30 Type 2 diabetic and 30 nondiabetic postmenopausal women with DEXA confirmed osteoporosis was chosen and their BMD levels, T-scores and serum Zn levels were compared.

Table 1: Distribution of respondents (diabetics and nondiabetics)


Parameter	Diabetic		Nondiabetic		P
	Mean	SD	Mean	SD	
Age (years)	59.44	7.42	59.36	7.44	0.9669
BMI (kg/m^2)	26.79	3.88	23.56	3.76	0.0018*
Height (cm)	152.95	5.29	151.03	6.45	0.2133
Weight (kg)	62.6	9.04	53.73	9.28	0.0004*
BMD (g/cm^2)	0.83	0.06	0.79	0.09	0.0661
T-score	-2.84	0.42	-3.22	0.74	0.0177*
Fasting serum glucose (mg/dL)	164.5	62.19	87.6	7.29	<0.001*
Serum zinc ($\mu\text{g}/\text{dL}$)	62.4	13.35	68.2	13.86	0.1042

* $P < 0.05$, SD: Standard deviation, BMI: Body mass index, BMD: Bone mineral density


Table 2: Relationship between BMD, serum zinc and BMI in diabetic and nondiabetic participants

Correlation between parameters	Diabetics	Nondiabetics
BMD and serum zinc	$r = 0.613, P < 0.001^*$	$r = 0.544, P < 0.002^*$
BMI and serum zinc	$r = 0.419, P = 0.021^*$	$r = 0.122, P = 0.522$
BMI with BMD	$r = 0.427, P = 0.019^*$	$r = 0.051, P = 0.791$


* $P < 0.05$, r = Pearson's correlation coefficient. BMI: Body mass index, BMD: Bone mineral density




Graph 1: Correlation of bone mineral density with serum zinc in diabetics



Graph 2: Correlation of bone mineral density with serum zinc in nondiabetics



Graph 3: Correlation of body mass index with serum zinc in diabetics



Graph 4: Correlation of body mass index with bone mineral density in diabetics

This study found higher BMD levels in diabetic postmenopausal women with osteoporosis than nondiabetics though not statistically significant, while the T-scores were significantly higher in the diabetic group.

The pathogenic complexity of diabetes mellitus has led to conflicting results on bone involvement in this condition. One of the chronic complications of diabetes is bone loss. Hyperglycemia generates a high concentration of Advanced Glycation End products in collagen. They decrease bone strength, promote osteoblast apoptosis and increase osteoclast-mediated bone resorption. Several clinical trials have supported that new bone formation, bone microarchitecture and thus bone quality are altered in both types of diabetes resulting in increased fracture risk. T2DM is associated with low bone turnover, which slows bone loss that may lead to higher BMD levels.^[15] Hyperinsulinemia is an important feature of T2DM. Insulin is structurally homologous to IGF-1 whose signaling pathway is crucial for bone acquisition. By interacting with the IGF-1 receptor present on osteoblasts, insulin shows an anabolic effect on bone. Hence, increased insulin levels in T2DM may also lead to high BMD levels.^[16] Despite high BMD, there could be increased bone fragility in T2DM patients through the accumulation of fatigue damage.^[15] Moreover, they have increased the risk of falling increasing the fracture risk.^[16]

Our study showed no significant difference in the mean serum Zn levels between the two groups though they were lower in Diabetics. This is in accordance with Zargar *et al.*^[17] and Diwan *et al.*^[18] who similarly found no significant difference in the serum Zn levels between type 2 Diabetics and controls. In contrast Emeribe *et al.*^[19] found significantly higher serum Zn levels in type 2 diabetics compared to controls that they thought could be due to the imbalanced rate of absorption to excretion of Zn.

Urinary discharge of Zn has been shown to be increased in postmenopausal osteoporotic women and the degree of zincuria correlates with the severity of osteoporosis.^[20,21] Zincuria as a feature of diabetes has been demonstrated by several studies.^[22,23] Walter *et al.*^[23] noted a six-fold increase in urinary Zn excretion in type 1 and type 2 diabetics compared with controls. Urinary Zn did not correlate significantly with plasma glucose concentrations, urinary glucose concentrations, HbA1c, proteinuria, or diuretic use.

Diabetic patients with renal diseases were excluded in this study. Hence, there must have been no significant zincuria in

the chosen diabetic group. Moreover the T2DM participants were on regular oral hypoglycemic drugs. They were on one or the other sulfonylurea drugs. Sulfonylureas act by stimulating pancreatic β cells to secrete more insulin.^[13] In addition, four participants were also on insulin therapy that did not contain Zn. Besides having anabolic effect on bone, insulin may reduce hyperzincuria accompanying diabetes.^[22] Insulin and sulfonylureas which increase insulin secretion might have also been responsible for the insignificant difference in the serum Zn levels between diabetic women as compared to the nondiabetics in our study.

None of the diabetic patients was taking thiazolidinediones. These drugs activate PPAR γ that induces adipogenesis over osteoblastogenesis in pluripotent cells.^[15]

Obesity is another important feature of T2DM. Diabetic women in this study were obese and had a significantly higher BMI than the nondiabetics ($P < 0.01$). This is in accordance with Zargar *et al.*^[17] The high BMI in this study was attributable to significantly higher weight of the diabetic women ($P < 0.001$). BMI of diabetics showed a significant positive correlation with Zn ($P < 0.05$). High BMI in T2DM may have a protective influence in maintaining serum Zn levels and to compensate for the decrease in Zn attributed to zincuria seen in DM.

Bone mineral density showed a significant positive correlation with serum Zn in both diabetics and nondiabetics. Serum Zn levels may serve to be important predictors of BMD.

Body mass index showed a significant positive correlation with BMD in the diabetic women ($P < 0.05$). This is in accordance with Brown *et al.*^[24] Low BMI (<20) has been considered as an important risk factor for osteoporosis.^[25] In obesity, there is increased conversion of androgens to estrogens by adipose tissue aromatase which have protective effects on bone.^[19] Adipose tissue also releases a wide variety of adipokines that have been implicated in the regulation of bone remodeling. Body fatness may have an impact on the accuracy of DEXA based BMD measurement. However, such a measurement error is negligible.^[16] No significant correlation between BMI and BMD in nondiabetics was found. There was not a major variation in the BMI among the nondiabetic participants, unlike diabetics. Most of them were in the normal BMI range. This could be the reason for the lack of correlation between BMI and BMD in nondiabetics.

In our study out of the 30 diabetic women, six were having FSG levels >200 mg/dL out of which three diabetic

participants were recently diagnosed with diabetes with duration of 1 month. Severe Hyperglycemia in these women can be a confounding factor. The rest three diabetic women may not have been under strict glycemic control.

CONCLUSION

T-scores are significantly higher in type 2 diabetic postmenopausal women with osteoporosis compared with nondiabetics. Serum Zn levels are not significantly altered in T2DM postmenopausal women with osteoporosis when patients with overt nephropathy are excluded. Medications with sulfonylureas and/or insulin may reduce the zincuria that may accompany diabetes. Serum Zn levels may serve to be important predictors of BMD. High BMI in T2DM may contribute to high BMD and may be a protective factor against zincuria seen in diabetes mellitus. However to what extent these factors that have helped in maintaining serum Zn levels prevent diabetes-related fractures cannot be told as fractures in T2DM are attributable to poor bone quality rather than BMD. Strict glycemic control is recommended. More studies are required in this arena on the role of Zn in postmenopausal women with diabetic osteoporosis.

Limitations of the study

- Both newly diagnosed and old cases of diabetes were included
- Patients with overt diabetic nephropathy were excluded, but testing for microalbuminuria in diabetics that is an early predictor of nephropathy was not done.

REFERENCES

1. Agrawal VK, Gupta DK. Recent update on osteoporosis. *Int J Med Sci Public Health* 2013;2:164-8.
2. Odabasi E, Turan M, Aydin A, Akay C, Kutlu M. Magnesium, zinc, copper, manganese, and selenium levels in postmenopausal women with osteoporosis. Can magnesium play a key role in osteoporosis? *Ann Acad Med Singapore* 2008;37:564-7.
3. Lindsay R, Cosman F. Osteoporosis. In: Longo DL, editor. *Harrison's Principles of Internal Medicine*. 18thed. United States of America: McGraw-Hill Publishers; 2012. p. 3120-35.
4. Damodaran P, Subramaniam R, Omar SZ, Nadkarni P, Paramsothy M. Profile of a menopause clinic in an urban population in Malaysia. *Singapore Med J* 2000;41:431-5.
5. Malhotra N, Mithal A. Osteoporosis in Indians. *Indian J Med Res* 2008;127:263-8.
6. Chau D, Edelman S. Osteoporosis and diabetes. *Clin Diabetes* 2002;20:153-7.
7. Hofbauer LC, Brueck CC, Singh SK, Dobnig H. Osteoporosis in patients with diabetes mellitus. *J Bone Miner Res* 2007;22:1317-28.
8. Hyun TH, Barrett-Connor E, Milne DB. Zinc intakes and plasma concentrations in men with osteoporosis: The Rancho Bernardo Study. *Am J Clin Nutr* 2004;80:715-21.
9. Molokwu CO, Li YV. Zinc homeostasis and bone mineral density. *Ohio Res Clin Rev* 2006;6:6-15.

10. Gupta N, Arora K. The status of trace elements after menopause: A comparative study. *J Clin Diagn Res* 2011;5:795-7.
11. Oh HM, Yoon JS. Glycemic control of type 2 diabetic patients after short-term zinc supplementation. *Nutr Res Pract* 2008;2:283-8.
12. Gunasekara P, Hettiarachchi M, Liyanage C, Lekamwasam S. Blood Sugar lowering effect of zinc and multi vitamin/mineral supplementation is dependent on initial fasting blood glucose. *J Diabetol* 2011;1:1-13.
13. Powers AC. Diabetes mellitus. In: Longo DL, editor. *Harrison's Principles of Internal Medicine*. 18thed. United States of America: McGraw-Hill Publishers; 2012. p. 2968-3003.
14. Trinder P. Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor. *Ann Clin Biochem* 1969;6:24-7.
15. Räkäl A, Sheehy O, Rahme E, LeLorier J. Osteoporosis among patients with type 1 and type 2 diabetes. *Diabetes Metab* 2008;34:193-205.
16. Ma L, Oei L, Jiang L, Estrada K, Chen H, Wang Z, *et al.* Association between bone mineral density and type 2 diabetes mellitus: A meta-analysis of observational studies. *Eur J Epidemiol* 2012;27:319-32.
17. Zargar AH, Shah NA, Masoodi SR, Laway BA, Dar FA, Khan AR, *et al.* Copper, zinc, and magnesium levels in non-insulin dependent diabetes mellitus. *Postgrad Med J* 1998;74:665-8.
18. Diwan AG, PradhanAB, LingojwarD, Krishna KK, Singh P, Almelkar SI. Serum zinc, chromium and magnesium levels in Type 2 diabetes. *Int J Diabetes Dev Ctries* 2006;26:122-3.
19. Emeribe AU, Akpotuzor JO, Etukudo MH. Serum chromium, zinc and testosterone levels in diabetics in university of Calabar Teaching Hospital Calabar Nigeria. *J Biol Agric Healthc* 2012;2:93-100.
20. Yanagisawa H. Zinc deficiency and clinical practice. *Jpn Med Assoc J* 2004;47:359-64.
21. Maser RE, Stabley JN, Lenhard MJ, Owusu-Griffin P, Provost-Craig MA, Farquhar WB. Zinc intake and biochemical markers of bone turnover in type 1 diabetes. *Diabetes Care* 2008;31:2279-80.
22. Chausmer AB. Zinc, insulin and diabetes. *J Am Coll Nutr* 1998;17:109-15.
23. Walter RM Jr, Uriu-Hare JY, Olin KL, Oster MH, Anawalt BD, Critchfield JW, *et al.* Copper, zinc, manganese, and magnesium status and complications of diabetes mellitus. *Diabetes Care* 1991;14:1050-6.
24. Brown SA, Sharpless JL. Osteoporosis: An under-appreciated complication of diabetes. *Clin Diabetes* 2004;22:10-20.
25. Florence R, Allen S, Benedict L, Compo R, Jensen A, Kalogeropoulou D, *et al.* In: *Diagnosis and Treatment of Osteoporosis*. 8thed. Bloomington: Institute for Clinical Systems Improvement; 2013. p. 1-87.

How to cite this article: Siddapur PR, Patil AB, Borde VS. Comparison of bone mineral density, T-scores and serum zinc between Diabetic and non diabetic postmenopausal women with osteoporosis. *J Lab Physicians* 2015;7:43-8.

Source of Support: Nil, **Conflict of Interest:** None declared.

Author Help: Reference checking facility

The manuscript system (www.journalonweb.com) allows the authors to check and verify the accuracy and style of references. The tool checks the references with PubMed as per a predefined style. Authors are encouraged to use this facility, before submitting articles to the journal.

- The style as well as bibliographic elements should be 100% accurate, to help get the references verified from the system. Even a single spelling error or addition of issue number/month of publication will lead to an error when verifying the reference.
- Example of a correct style
Sheahan P, O'leary G, Lee G, Fitzgibbon J. Cystic cervical metastases: Incidence and diagnosis using fine needle aspiration biopsy. *Otolaryngol Head Neck Surg* 2002;127:294-8.
- Only the references from journals indexed in PubMed will be checked.
- Enter each reference in new line, without a serial number.
- Add up to a maximum of 15 references at a time.
- If the reference is correct for its bibliographic elements and punctuations, it will be shown as CORRECT and a link to the correct article in PubMed will be given.
- If any of the bibliographic elements are missing, incorrect or extra (such as issue number), it will be shown as INCORRECT and link to possible articles in PubMed will be given.