


New techniques in endocytoscopy: submucosal injection heightens the visibility


Endocytoscopy involves the use of a light contact microscope capable of magnifying endoscopic images up to $\times 520$, allowing optical biopsy to the cellular level [1, 2].

A 69-year-old patient had an 8-mm 0-IIa lesion located in the transverse colon. We performed colonoscopy using endocytoscopy (CF-H290ECI; Olympus, Tokyo, Japan) without distal attachment (▶ **Video 1**).

Magnifying narrow-band imaging (NBI) showed Japan NBI Expert Team classification type 2A around the whole surface of the lesion (▶ **Fig. 1 a**) [3]. We then stained the lesion with 0.05% crystal violet. Magnifying chromoendoscopy revealed type IIIIL pit pattern (▶ **Fig. 1 b**) based on Kudo's classification [3].



▶ **Video 1** Endocytoscopy for an 8-mm 0-IIa lesion, located in the transverse colon. Because of the lack of light, we performed submucosal injection. Submucosal injection increased visualization of the lesion at the cellular level. EC, endocytoscopy.



▶ **Fig. 1** Endoscopic findings, treatment, and pathology. **a** An 8-mm 0-IIa lesion was located in the transverse colon. **b** Magnifying chromoendoscopy using crystal violet staining identified a type IIIIL pit pattern. **c** The lesion was stained with methylene blue and endocytoscopic examination was performed ($\times 520$). **d** Submucosal injection was performed and endocytoscopic examination was repeated. **e** The lesion was pathologically diagnosed as low grade adenoma.

Subsequently, the lesion was stained with 1% methylene blue (► **Fig. 1 c**, ×520). Diagnosis was made on the basis of Kudo's endocytoscopy classification [4]. However, it was difficult to evaluate lumina and nuclei, and we considered the lack of light to be the cause. Therefore, we performed submucosal injection of 0.9% saline under the lesion. In addition to the light reflected from the scope into the submucosal layer, submucosal injection led to a thickening of the submucosal layer, increasing the amount of light from the surroundings. The increased light intensity provided a clearer contrast to the background stroma and improved the visibility of the lumina and nuclei. The endocytoscopy image showed slit-like smooth lumina, and uniform and roundish nuclei (► **Fig. 1 d**). Furthermore, because we did not find irregular and rough lumina and a large number of roundish nuclei, we diagnosed the lesion as EC2. Endoscopic mucosal resection was performed and histopathological findings showed low grade adenoma (► **Fig. 1 e**).

Submucosal injection improved cellular visualization because swelling of the submucosal layer with saline strengthened the contrast between stroma, lumina, and nuclei. This method may improve the diagnostic ability of endocytoscopy. Further assessment is necessary to evaluate the effectiveness.

Endoscopy_UCTN_Code_TTT_1AQ_2A]

Competing interests

The authors declare that they have no conflict of interest.

The authors

Yoshiaki Kimoto¹, Ken Ohata¹, Maiko Takita¹, Yohei Minato¹, Takashi Muramoto¹, Yoshiyuki Kayashima^{2,3}, Ikuro Koba²

- 1 Department of Gastrointestinal Endoscopy, NTT Medical Center Tokyo, Tokyo, Japan
- 2 Department of Gastroenterology, Yamaga Chuo Hospital, Kumamoto, Japan
- 3 Department of Gastroenterology, Faculty of Medicine, Fukuoka University, Fukuoka, Japan

Corresponding author

Ken Ohata, MD, PhD

Department of Gastrointestinal Endoscopy, NTT Medical Center Tokyo, 5-9-22 Higashi-gotanda, Shinagawa-ku, Tokyo 141-8625, Japan
 Fax: +81-3-34486541
 ken.ohata1974@gmail.com

References

- [1] Inoue H, Kazawa T, Sato Y et al. In vivo observation of living cancer cells in the esophagus, stomach, and colon using catheter-type contact endoscope, "Endo-Cytoscopy system". *Gastrointest Endosc Clin N Am* 2004; 14: 589–594
- [2] Kudo T, Suzuki K, Kudo SE et al. Endocytoscopy for the differential diagnosis of colorectal low-grade adenoma: a novel possibility for the "resect and discard" strategy. *Gastrointest Endosc* 2020; 91: 676–683
- [3] Sano Y, Tanaka S, Kudo SE et al. Narrow-band imaging (NBI) magnifying endoscopic classification of colorectal tumors proposed by the Japan NBI Expert Team. *Dig Endosc* 2016; 28: 526–533
- [4] Kudo SE, Misawa M, Wada Y et al. Endocytoscopic microvasculature evaluation is a reliable new diagnostic method for colorectal lesions (with video). *Gastrointest Endosc* 2015; 82: 912–923

Bibliography

Endoscopy 2021; 53: E245–E246
 DOI 10.1055/a-1252-1963
 ISSN 0013-726X
 published online 23.9.2020
 © 2020. Thieme. All rights reserved.
 Georg Thieme Verlag KG, Rüdigerstraße 14,
 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS
<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>