

# Profuse diarrhea after percutaneous change of a computed tomography-guided gastrostomy: endoscopic closure of a colocutaneous fistula



A 71-year-old patient was admitted after the accidental dislodgement of a computed tomography (CT)-guided percutaneous gastrostomy, placed 1 year prior. The initial procedure was uneventful and facilitated rapid weight gain. A percutaneous MIC-KEY tube (Avanos, Alpharetta, Georgia, USA) was rapidly placed along the previous route. As soon as nutrition was restarted, “on and off” profuse diarrhea occurred, ceasing as the nutrition was stopped and resuming immediately upon resumption. Opacification of the MIC-KEY tube revealed a fistulous tract between the transverse colon and the skin (► **Video 1**). Conservative treatment was decided, involving the endoscopic closure of the residual colocutaneous fistula, achieved by placing an Ovesco clip (Ovesco Endoscopy, Tübingen, Germany) during a colonoscopy [1]. In conclusion, a transcolonic perforation may go unnoticed during the months [2] after a CT-guided percutaneous gastrostomy. Colocutaneous fistulous tract can be safely closed using an Ovesco clip during a colonoscopy [1, 3], to avoid the need for surgery.

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AK

## Acknowledgement

We address our acknowledgments to Mr. Charles Duboc for the creation of the animated graphic illustrations.

## Conflict of Interest

The authors declare that they have no conflict of interest.



► **Video 1** Management of a silent colonic transfixation after computed tomography-guided gastrostomy. The colocutaneous fistula was revealed by diarrhea and could be safely closed with an Ovesco clip (Ovesco Endoscopy, Tübingen, Germany).

## The authors

**Amani Al Khatib**<sup>1,2</sup>, **Anaïs Bertrand**<sup>1,2</sup>, **Henri Duboc**<sup>1,2</sup>, **Olivier Bonsack**<sup>1,2</sup>, **Benoît Coffin**<sup>1,2</sup>, **Heithem Soliman**<sup>1,2</sup>

- 1 Centre de Recherche sur l'Inflammation, Inserm UMRS 1149, Université Paris Cité, Paris, France
- 2 Hépatogastro-entérologie, Hôpital Louis Mourier, APHP, Colombes, France

## Corresponding author

**Henri Duboc, MD, PhD**  
Hépatogastro-entérologie, Hôpital Louis Mourier, 178 rue des Renouillers, Colombes 92700, France  
henri.duboc@aphp.fr

## References

- [1] Rodrigues-Pinto E, Santos AL, Macedo G. Endoscopic closure of a colocutaneous fistula after placement of percutaneous endoscopic gastrostomy. *Endoscopy* 2020; 52: 187–188. doi:10.1055/a-1063-6276
- [2] Van Gossum A, DesMarez B, Cremer M. A colo-cutaneous-gastric fistula: a silent and unusual complication of percutaneous endoscopic gastrostomy. *Endoscopy* 1988; 20: 161. doi:10.1055/s-2007-1018166
- [3] Friedmann R, Feldman H, Sonnenblick M. Misplacement of percutaneously inserted gastrostomy tube into the colon: report of 6 cases and review of the literature. *JPEN J Parenter Enteral Nutr* 2007; 31: 469–476. doi:10.1177/0148607107031006469

## Bibliography

*Endoscopy* 2024; 56: E584  
DOI 10.1055/a-2358-1166  
ISSN 0013-726X  
© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited.  
(<https://creativecommons.org/licenses/by/4.0/>)  
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

