

Use of Simini Protect Lavage as an Adjuvant in the Antiseptic Protocol for Revision Surgeries Involving Total Hip Replacement

Ida Forzisi¹ Luca Vezzoni¹ Milo Bozzerla² Aldo Vezzoni¹

 ¹ Surgical Department, Clinica Veterinaria Vezzoni, Cremona, Italy
² Surgical Deparment, AniCura - Clinica Veterinaria Malpensa, Samarate, Italy Address for correspondence Ida Forzisi, DVM, Clinica Veterinaria Vezzoni, Via delle Vigne 190, Cremona 26100, Italy (e-mail: idaforzisi@gmail.com).

VCOT Open 2025;8:e13-e20.

Abstract

Objectives Despite advances in surgical asepsis and improvements in sterile surgical techniques, surgical site infection occurs in 0.8 to 15.8% of surgical wounds. Most infections are caused by microorganisms forming biofilms that adhere to implants or dead bone and can evade host immunity and antimicrobial agents. Simini Protect Lavage (SPL) is a hypertonic aqueous solution designed to break cross-links within the extracellular polymeric substance of biofilms produced by bacteria. This study aimed to report the outcome of our total hip replacement (THR) revision cases using an SPL in addition to our standard antiseptic protocol.

Methods The medical records of patients at Vezzoni Veterinary Clinic were searched for animals undergoing a revision orthopaedic surgery involving a THR that included the use of SPL. In all patients, a minimum of 1-year follow-up and culture and a sensitivity test pre- and post-Simini Lavage were required.

Results SPL was used in 36 cases. At the beginning of surgery, culture and sensitive tests were positive in eight patients, and at the end of surgery, seven dogs were still positive for bacterial infection. Only one case, which experienced three revision surgeries, underwent explantation due to persistent, clinically evident infection. The other dogs showed neither clinical nor X-ray signs of infection over a long period.

Keywords ► infections

- ► surgical site infection
- revision surgery
- intraoperative antiseptic lavage

other dogs showed neither clinical nor X-ray signs of infection over a long period. **Clinical Significance** The use of SPL can be considered as an adjuvant in the antiseptic protocol for the one-stage revision THR surgeries without a plan of prosthesis explanation. Nevertheless, further studies are needed to objectively evaluate the efficacy of SPL in eradicating the postoperative infections.

Introduction

Surgical site infection (SSI) is costly and associated with poor patient outcomes, increased morbidity, and high reoperation rates. The complications posed by SSI are particularly challenging to resolve, especially in orthopaedic cases. Despite advances in surgical asepsis, the rate of SSI ranges from 0.8 to 18.1%, depending on patient characteristics, the degree of

received June 30, 2024 accepted after revision December 15, 2024 DOI https://doi.org/ 10.1055/a-2506-0553. ISSN 2625-2325. wound contamination, the surgical environment, and the type of surgical procedure.^{1,2} Factors increasing the risk of SSI include prolonged anaesthesia and surgery times, body weight, the number of people in the operating suite, hospitalization time, the physical status score according to the American Society of Anaesthesiologists physical status classification system, hypotension, type of surgery, and open fractures.^{3–7}

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Oswald-Hesse-Straße 50, 70469 Stuttgart, Germany

^{© 2025.} The Author(s).

The management of periprosthetic joint infection is a significant challenge, with irrigation and debridement playing a pivotal role in addressing SSI. Therefore, meticulous mechanical and chemical debridement of nonviable tissue and judicious selection of irrigation solutions are imperative in minimizing the presence of infected debris.

Total hip replacement (THR) is considered the gold standard treatment for some of the common canine coxofemoral joint diseases like end-stage osteoarthritis and severe hip dysplasia.^{8–11} Predisposing factors for infection in dogs following THR include revision surgery, preexisting hip infection, prolonged surgical time, and hematogenous spread secondary to an infection in a remote area.^{12,13}

In both veterinary and human medicine, the gold standard for treating SSI in orthopaedic surgery involves the removal of implants, antibiotic therapy, and, if required, a two-stage surgical revision procedure.^{14,15}

The most difficult infections to treat in orthopaedic surgery are those caused by microorganisms that adhere to implants or dead bone (sequestra), forming a biofilm that makes them resistant to host defence mechanisms and most antimicrobial agents.^{16–18} Curing an existing infection is difficult because, although antibiotics can kill planktonic or free-living bacteria, they are usually ineffective against bacteria in biofilms.¹⁹

Costerton et al first proposed that communities of bacteria embedded in a highly hydrated polysaccharide matrix called a "biofilm," mediated adhesion to solid-liquid interfaces. These functionally heterogeneous microcolonies or single-cell aggregates are enclosed in a matrix of self-produced extracellular polymers, which may adhere to biotic or abiotic surfaces.^{20,21} In the 1980s, it was shown that most bacteria and fungi can form a biofilm, which protect them from biotic and abiotic stresses, providing a survival mechanism in hostile environments.^{22–25} It has been reported that biofilm-producing bacteria encapsulated in an extracellular polymeric substance (EPS), which may also contain components derived from the host, are more tolerant to antibiotics and antiseptics, resisting antimicrobial concentrations 100 to 1,000 times higher than those effective against their planktonic counterparts.^{26–29}

The bacteria most frequently involved in SSI in veterinary medicine are *Staphylococcus* spp., ³⁰ *Streptococcus* spp., and *Escherichia coli*,³¹ with *Staphylococcus pseudintermedius* being the most common in dogs. Based on recent studies, approximately 50% of bacteria isolated from implant-associated osteomyelitis were methicillin-resistant *Staphylococcus* spp. strains.^{3,32–34}

Various strategies are utilized to mitigate the risk of SSIs. These encompass perioperative antibiotic prophylaxis, applying specific antiseptic solutions for skin preparation, intraoperative topical adjuvants, antiseptic irrigation, meticulous tissue handling, ensuring effective hemostasis, and adhering to rigorous aseptic techniques.³⁵ Intraoperative irrigation is a commonly employed technique advocated in all surgical disciplines before incision closure. Irrigation hydrates the surgical bed, facilitates examination of the area immediately before closure, removes blood clots and

Povidone–iodine has been recommended as an irrigation solution due to its potent oxidizing properties, which inactivate cell membranes and intracellular constituents in a concentration-dependent manner. In a recent study evaluating largely index THR surgeries, no postoperative infections developed after 3-minute lavage with 0.35% povidone–io-dine solution, followed by rinsing with sterile saline before closing the incision in 97 primary THR and five revisions for luxation.⁴⁰

Recent studies have indicated that a 0.35% solution of povidone–iodine is effective in eradicating methicillin-susceptible *Staphylococcus aureus* and *E. coli*, also showing the ability to remove periprosthetic joint infection bacteria biofilm on orthopaedic material, with negligible cytotoxic effects on osteoblasts, chondrocytes, and fibroblasts.^{36,38,41–46}

Chlorhexidine gluconate (CHG) is a cationic bisbiguanide that binds to the negatively charged cell wall of bacteria, disrupting their osmotic equilibrium. Chlorhexidine gluconate has a broad spectrum of activity and is highly effective against various microorganisms responsible for SSI, including methicillin-susceptible S. aureus, methicillin-resistant S. aureus, coagulase-negative Staphylococcus, Gram-negative bacteria, fungi, and mycobacteria. The bactericidal effect of CHG is almost immediate, with maximum uptake occurring within 20 seconds of exposure. The duration of effect is directly related to the duration of exposure and concentration. The antimicrobial activity of CHG may last for an extended period, preventing the attachment and growth of other bacteria on surfaces. Although the optimal CHG dilution for intraoperative irrigation is debated, concentrations of 0.05 to 0.5% may effectively reduce bacterial load without being cytotoxic.47,48

In human medicine, a surgical lavage solution called Bactisure Wound Lavage (Zimmer Biomet) has recently been shown to reduce the bacterial load within the surgical site. Bactisure is a preformulated combination solution of ethanol, acetic acid, sodium acetate, and benzalkonium chloride in water. The acetic acid chelates the metal ions in the exopolysaccharide matrix, and the sodium acetate acts as a buffer to maintain an ideal pH. Benzalkonium chloride is a surfactant that reduces the surface tension of the biofilm and lyses bacterial cells. This lavage solution deconstructs EPS, which facilitates the exposure of the bacteria to antibiotics and the innate immune system, potentially leading to their subsequent eradication through lavage.49,50 An identical solution was developed for use in veterinary medicine: Simini Protect Lavage (SPL). It is a hypertonic aqueous solution composed of ethanol (solvent), acetic acid (pH modifier), sodium acetate (buffer), benzalkonium chloride 0.13% (surfactant), and water. SPL is designed to break crosslinks within the EPS of biofilms. Its effects become effective within 1 minute and must be removed by rinsing with an equal or greater volume of saline solution.

To the authors' knowledge, available information about the most effective treatment for SSI following orthopaedic surgery, which would also allow for one-stage revision surgery, is limited. Therefore, the purpose of this study was to report the short- and long-term clinical outcome of our THR revision case series while using the SPL added to our standard antiseptic protocol.

Methods

Case Selection

The medical records of dogs that underwent a revision orthopaedic surgery involving Zurich Cementless THRs (Kyon, CH) from November 2019 to December 2022, where SPL was used at Vezzoni Veterinary Clinic (Cremona, Italy), were retrospectively evaluated. To be included in the study, animals were required to have a clinical and radiographic follow-up evaluation a minimum of 1 year postoperatively, along with culture and sensitivity tests both before and after Simini lavage. Data collected included signalment, indication for the surgery, number of revisions surgeries, culture and antimicrobial susceptibility testing results before and after SPL, type and class of bacteria isolated, and outcome at the last reevaluation.

Surgical Procedure

All surgeries were performed by one of the authors (A.V.). The animals underwent a preoperative antibiotic prophylaxis protocol that included 20 mg/kg cefazolin sodium (Cefazolina 1 g, Teva, Italy) administered intravenous 30 minutes before the skin was incised and then every 90 minutes until the end of surgery. After the induction of general anaesthesia, the hair was clipped according to the procedure required, and the animals were moved to the operating theatre and positioned on the surgical table. Preoperative skin asepsis was achieved using a solution of 4.0% chlorhexidine digluconate and 2.6% isopropyl alcohol, followed by applying a solution of 0.5% CHG and 66.0% denatured ethyl alcohol. This procedure was repeated at least four times with a contact time of 8 minutes. Before draping, a coloured propanol-based

skin antiseptic solution (Cutasept G; propan 2 of 72% + benzalkonium chloride; Bode-Chemie, Germany) was applied with a contact time of 2 minutes. SPL was used after debridement and implant removal and again after new implant positioning immediately before closing. SPL was employed in every surgical revision surgery and consistently in dogs undergoing multiple revision surgeries. The volume of SPL fluid used in each case was sufficient to cover the surgical wound. It was allowed to remain in contact with the wound for 1 minute, after which it was removed by rinsing with an equal or greater volume of saline solution (**Figs. 1**, **2**). Two samples were collected for culture: the first at the beginning of the surgical procedure, from the synovial membrane, and the second from the joint before its closure. In all cases, a gentamicin-impregnated collagen sponge (Genta-coll Resorb) was applied after SPL and after the culture sample before suturing of the deep tissues (**Fig. 2**).

All animals received 20 mg/kg of amoxicillin/clavulanic acid (Konclav, Fatro, Italy) orally, three times a day, until the culture and antimicrobial susceptibility testing results were available. Antibiotic therapy was then discontinued or adjusted based on the test results. In case with a positive culture at the end of surgery, antibiotic therapy was continued for 10 days.

Given on the retrospective observational nature of the study, ethical approval was not required; however, signed informed owner consent was obtained before all surgeries.

Results

Simini Protect Lavage was used in 36 cases of THR revision surgery between November 2019 and December 2022. The indication for surgery included dislocation of the prosthetic head (n = 21), cup loosening (n = 5), periprosthetic femoral fracture following THR (n = 2), stem breakage (n = 4), revision with THR of a capital physeal fracture previously treated with internal fixation (n = 1), revision of double pelvic osteotomy to implant the THR (n = 1), revision of extraarticular iliofemoral suture to implant the THR (n = 1), revision of infected THR (n = 1).



Fig. 1 Intraoperative images showing collection of a sample for microbial culture after debridement (A) and application of Simini Protect lavage fluid (B) using enough lavage fluid to cover the surgical wound (C).



Fig. 2 Intraoperative images showing that Simini Protect Lavage fluid becomes black during 1 minute of contact with the surgical wound (left) and application of a Genta-coll Resorb sponge before closure (right).

Five dogs experienced three revision surgeries in a short period (cases: 1, 6, 7, 11, 21), 8 had two revision surgeries (cases: 2, 3, 4, 5, 13, 30, 32, 33), and the remaining 23 underwent one THR revision surgery (cases: 8, 9, 10, 12, 14, 15, 16, 17, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29, 31, 34, 35, 36).

The mean time of anaesthesia was 118.47 ± 47.08 minutes (range: 70–80 minutes, median: 100 minutes). The mean time of surgery was 79.58 ± 46.86 minutes (range: 25–240 minutes, median 60 minutes).

The bacterial culture results at the beginning of surgery were positive in 8 cases (cases: 1, 2, 6, 7, 8, 9, 10, 11) and negative in 28 (cases: 3, 4, 5, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36). At the end of the surgery, 7 dogs were still positive for bacterial infection (cases: 1, 2, 3, 4, 5, 6, 7), whereas the other 29 dogs had a negative culture and sensitive test with no postoperative infection developing (cases: 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36).

The bacteria isolated at the beginning of surgery included *S. pseudintermedius* (methicillin-resistant *S. pseudintermedius* [MRSP, n = 5]), *Pseudomonas aeruginosa* (n = 2), and *Enterococcus faecalis* (n = 1). At the end of the surgery, after SPL was applied, the bacteria isolated were *Enterobacter cloacae* (n = 1) *and S. pseudintermedius* MRSP (n = 6).

Regarding the seven dogs with a positive swab at the end of the surgery, three underwent three consecutive revision surgeries (cases: 1, 6, 7), and four dogs underwent two consecutive revision surgeries due to recurrent luxation (cases: 2, 3, 4, 5). Only one case (case 1), which experienced three revision surgeries, required total implant explantation. The bacteria isolated at the end of the first revision surgery was the same as the bacteria isolated at the end of the following revision surgery in all seven cases.

In the other two cases (cases: 6, 7), the implant removal was carried out not for reasons related to infections, as the swabs were negative at the beginning and end of surgery, but due to the recurrence of THR dislocation.

The other dogs showed neither clinical nor radiographic signs of infection, with good outcomes over a long period (**►Table 1**).

In our cases, no complications, such as adverse tissue or bone reactions, delayed bone healing, or surgical wounds, were observed.

The mean duration of the last clinical and radiographic follow-up in cases where implants were not removed (n = 33) was 865.27 days (range: 612–1,557 days).

Of the 36 patients who received treatment, 34 showed no clinical or radiographic signs of infection, with only two cases exhibiting such signs. In one case (case 2) that underwent two surgical revision surgery with a positive swab for *S. pseudintermedius* MRSP at the beginning and end of the surgery, still showed signs of lysis around the stem screws. However, it was not clinically significant, and was managed conservatively.

Contrary to another case (case 1) that underwent three surgical revisions, with a positive swab for *S. pseudinterme-dius* (MRSP) at the beginning and at the end of the surgery for each surgical revision, which instead required explantation.

Discussion

Surgical site infection is an infection that occurs within 30 days of surgery and affects the incision, organs, or spaces in the body where the operation occurred.⁵¹

SSI is particularly challenging to resolve in orthopaedic surgery due to the possibility of bacterial biofilm formation on the implants. Biofilm adheres to implants, resists eradication, and necessitates implant removal to eliminate the infection.^{32,52–54} For this reason, cases of revision surgeries were used to evaluate the effect of SPL on biofilms as an adjuvant to all previous measures.

In our study, only one case (case 8) underwent a positive preoperative culture and sensitivity test before the revision surgery, suggesting the probable presence of biofilm on the implants. This dog had previously experienced unsuccessful antibiotic treatment for the infection. However, using the SPL as an adjuvant during the revision surgery resulted in a successful outcome, effectively resolving the infection without requiring the removal of the prosthesis.

A preoperative culture and sensitivity test were not performed before the surgical revision for the other seven cases. However, the swab taken at the beginning of the

| ID case | Number of revision surgery | Coculture and sensitive test before the use of Simini Protect Lavage | Coculture and sensitive test after the use of Simini Protect Lavage | Outcome |
|---------|-------------------------------|--|---|---------|
| 1 | 3 | Staphylococcus pseudintermedius MRSP | Staphylococcus pseudintermedius MRSP | Explant |
| 2 | 2 | Staphylococcus pseudintermedius MRSP | Staphylococcus pseudintermedius MRSP | Good |
| 3 | 2 | Negative | Staphylococcus pseudintermedius MRSP | Good |
| 4 | 2 | Negative | Enterobacter cloacae | Good |
| 5 | 2 | Negative | Staphylococcus pseudintermedius MRSP | Good |
| 6 | 3 | Staphylococcus pseudintermedius MRSP | Staphylococcus pseudintermedius MRSP | Good |
| 7 | 3 | Staphylococcus pseudintermedius MRSP | Staphylococcus pseudintermedius MRSP | Good |
| 8 | 1 | Pseudomonas aeruginosa | Negative | Good |
| 9 | 1 | Pseudomonas aeruginosa | Negative | Good |
| 10 | 1 | Staphylococcus pseudintermedius MRSP | Negative | Good |
| 11 | 3 | Staphylococcus pseudintermedius MRSP | Negative | Good |
| 12 | 1 | Negative | Negative | Explant |
| 13 | 2 | Negative | Negative | Explant |
| 14 | 1 | Negative | Negative | Good |
| 15 | 1 | Negative | Negative | Good |
| 16 | 1 | Negative | Negative | Good |
| 17 | 1 | Negative | Negative | Good |
| 18 | 1 | Negative | Negative | Good |
| 19 | 1 | Negative | Negative | Good |
| 20 | 1 | Negative | Negative | Good |
| 21 | 3 | Negative | Negative | Good |
| 22 | 1 | Negative | Negative | Good |
| 23 | 1 | Negative | Negative | Good |
| 24 | 1 | Negative | Negative | Good |
| 25 | 1 | Negative | Negative | Good |
| 26 | 1 | Negative | Negative | Good |
| 27 | 1 | Negative | Negative | Good |
| 28 | 1 | Negative | Negative | Good |
| 29 | 1 | Negative | Negative | Good |
| 30 | 2 | Negative | Negative | Good |
| 31 | 1 | Negative | Negative | Good |
| 32 | 2 | Negative | Negative | Good |
| 33 | 2 | Negative | Negative | Good |
| 34 | 1 | Negative | Negative | Good |
| 35 | 1 | Negative | Negative | Good |
| 36 | 1 | Negative | Negative | Good |

Table 1 Summary of the cases for surgical revision of total hip replacement in which Simini Protect Lavage was used

Abbreviation: MRSA, methicillin-resistant Staphylococcus pseudintermedius.

revision surgery tested positive for SSI, which may have compromised the surgical revision. The authors believe that using SPL as an adjuvant helped to avoid or limit the progression of these infections.

In the 28 cases where infection was not present at the beginning of the surgery, the authors suggest that the aseptic

techniques, including the use of SPL, facilitated multiple surgical revisions, up to three in the same patient, without the occurrence of SSIs. Only two cases required implant removal, but this was for reasons unrelated to infection.

Out of the 28 cases, three presented with a negative swab prior to surgery but later exhibited signs of infection with a

positive swab postsurgery. These three cases underwent two surgical revisions, and subsequent swabs consistently returned positive results. Nonetheless, none of the three cases displayed clinical or radiographic signs of infection. The authors believe that the intraoperative technique utilized to disrupt the biofilm played a crucial role in allowing the antibiotics to eliminate the infection. However, it is important to recognize that, at this time, there is a lack of concrete evidence to substantiate this assertion.

Simini Protect Lavage was developed in veterinary medicine to remove planktonic bacteria and biofilms. It destroys the EPS matrix, exposing the bacteria to host defence mechanisms and antibiotics, and promotes eliminating bacteria from the surgical site. Ethanol, a key component, dehydrates and denatures proteins, interferes with the cell plasma membrane, disrupts cell metabolism, and halts bacterial growth.^{55,56}

Specific procedures and patient-related factors can increase the risk of SSI. Operative time is often a predictor factor, with increased times associated with more complex surgical procedures. It is well known that longer operative times predispose patients to increased bacterial contamination. Eugster et al⁵⁷ reported that the risk of infection increased 1.01 times for each additional minute of surgery. Pratesi et al found that the risk of infection in orthopaedic surgeries increased by 2% for each additional minute of anaesthetic time.⁵⁸ That is an important consideration, especially when planning for the revision surgery. The estimated time of any revision surgery is difficult to predict due to the altered anatomy, need for implant removal and its replacement, which can result in longer anaesthetic times and a higher risk of postoperative infection.

In our study, gentamicin-impregnated collagen sponges were also applied, and antibiotic therapy was administered to all patients. Indeed, this represents a limitation of this study, as its purpose was not to compare the different additives but only to report our clinical experience using a new preformulated irrigation solution that allows surgical revisions to be carried out without the need to remove existing implants in which a biofilm had likely formed.

In our study, seven dogs had a positive culture and sensitivity test after using SPL. However, only one case that experienced three consecutive revision surgeries developed a clinical infection requiring THR explantation. The literature describes the possibility of a positive swab at the end of the surgery due to the duration of the surgery and anaesthesia. It has also been described as a positive swab after using antiseptic solution irrigation, particularly in human medicine after using Bactisure Wound Lavage. The authors reported that there was not a statistically significant increase in culturable bacteria after the wash, which may indicate only the liberation of bacteria from the biofilm, thus allowing antibiotic therapy to act.^{13,59} Based on the findings above, it is plausible that the SPL exerted an effect in six out of the seven patients who tested positive for SSI, thereby permitting the antibiotic to function without leading to a long-term infectious condition.

The ideal volume, pressure, and irrigation duration for treatment or prevention of SSI need to be better defined. In human medicine, pulsed lavage with Bactisure Wound Lavage is reported to remove wound debris. Low-pressure lavage may be adequate for SSI with low bacterial load but may not be effective for deep SSI where bacteria and biofilm adhere to the implants and tissues.¹² Our study only applied the SPL to the wound without a pulse lavage. This procedure is based only on our subjective observation and should be verified with further studies.

Another essential aspect of antiseptic lavage is that not all commercially available antiseptic solutions are sterile, leading to nonsterile or multiuse containers harbouring contaminants. An advantage of the SPL is that it is dispensed in single-use 20mL vials, which helps maintain the solution's sterility.

No complications directly related to SPL were observed. Wound healing and implant osseointegration were uneventful, suggesting that SPL was clinically nonirritating and noncytotoxic.

Conclusions

Simini Protect Lavage was used in 36 cases of surgical THR revisions with no signs of SSI based on both radiographic and clinical long-term observation in 35 cases. Only one dog, which underwent three revision surgeries, exhibited clinical and radiographic signs of persistent infection, requiring a THR explant.

In the context of surgical revision procedures, implementing a valid protocol for maintaining a sterile environment is of utmost importance. The incorporation of SPL into the antiseptic protocol in our total hip replacement revision case series seemed safe without causing further disruption to the osteointegration of the preserved implants or without negatively influencing postoperative surgical wound healing. The use of SPL as an adjuvant in the antiseptic protocol can be considered in THR revision surgeries even without a plan for prosthesis explantation. According to the conclusions of the in vitro study by Marquez-Gomez and colleagues, the use of several antiseptic agents in the same operation may be the most effective solution to treat and prevent periprosthetic infections.⁵⁹

This work describes a protocol employed in a case series. One notable limitation of this study pertains to its retrospective nature and the absence of a control group.

Further studies are needed to objectively evaluate the efficacy of SPL in eradicating postoperative infections.

Author's Contribution

A.V. and I.F. contributed to the conception of the study. A.V., I.F., L.V., and M.B. contributed to the study design, acquisition of data, data analysis, and interpretation. All authors drafted, revised, and approved the submitted manuscript.

Funding None.

Conflict of Interest None declared.

Acknowledgments

None.

References

- 1 Vasseur PB, Paul HA, Enos LR, Hirsh DC. Infection rates in clean surgical procedures: a comparison of ampicillin prophylaxis vs a placebo. J Am Vet Med Assoc 1985;187(08):825–827
- 2 Johnston SA, Tobias KM. Veterinary Surgery Small Animal. Elsevier; 2018:148–155
- 3 Turk R, Singh A, Weese JS. Prospective surgical site infection surveillance in dogs. Vet Surg 2015;44(01):2–8
- 4 Nelson LL. Surgical site infections in small animal surgery. Vet Clin North Am Small Anim Pract 2011;41(05):1041–1056, viii
- 5 Weese JS. A review of post-operative infections in veterinary orthopaedic surgery. Vet Comp Orthop Traumatol 2008;21(02): 99–105
- 6 Frey TN, Hoelzler MG, Scavelli TD, Fulcher RP, Bastian RP. Risk factors for surgical site infection-inflammation in dogs undergoing surgery for rupture of the cranial cruciate ligament: 902 cases (2005-2006). J Am Vet Med Assoc 2010;236(01):88–94
- 7 Beal MW, Brown DC, Shofer FS. The effects of perioperative hypothermia and the duration of anesthesia on postoperative wound infection rate in clean wounds: a retrospective study. Vet Surg 2000;29(02):123–127
- 8 Paul HA, Bargar WL. A modified technique for canine total hip replacement. J Am Anim Hosp Assoc 1987;23:13–18
- 9 Olmstead ML. Total hip replacement. Vet Clin North Am Small Anim Pract 1987;17(04):943–955
- 10 Olmstead ML. The canine cemented modular total hip prosthesis. J Am Anim Hosp Assoc 1995;31(02):109–124
- 11 Vezzoni L, Vezzoni A, Boudrieau RJ. Long-term outcome of Zurich cementless total hip arthroplasty in 439 cases. Vet Surg 2015;44 (08):921–929
- 12 Christopher ZK, Deckey DG, Pollock JR, Spangehl MJ. Antiseptic irrigation solutions used in total joint arthroplasty: a critical analysis review. JBJS Rev 2022;10(03):
- 13 Ireifej S, Marino DJ, Loughin CA, Lesser ML, Akerman M. Risk factors and clinical relevance of positive intraoperative bacterial cultures in dogs with total hip replacement. Vet Surg 2012;41 (01):63–68
- 14 Senthi S, Munro JT, Pitto RP. Infection in total hip replacement: meta-analysis. Int Orthop 2011;35(02):253–260
- 15 Dyce J, Olmstead ML. Removal of infected canine cemented total hip prostheses using a femoral window technique. Vet Surg 2002; 31(06):552–560
- 16 Zimmerli W, Sendi P. Orthopaedic biofilm infections. APMIS 2017; 125(04):353–364
- 17 Stoodley P, Ehrlich GD, Sedghizadeh PP, et al. Orthopaedic biofilm infections. Curr Orthop Pract 2011;22(06):558–563
- 18 Davies D. Understanding biofilm resistance to antibacterial agents. Nat Rev Drug Discov 2003;2(02):114–122
- 19 Perencevich EN, Sands KE, Cosgrove SE, Guadagnoli E, Meara E, Platt R. Health and economic impact of surgical site infections diagnosed after hospital discharge. Emerg Infect Dis 2003;9(02): 196–203
- 20 Costerton JW, Geesey GG, Cheng KJ. How bacteria stick. Sci Am 1978;238(01):86–95
- 21 Costerton JW, Lewandowski Z, Caldwell DE, Korber DR, Lappin-Scott HM. Microbial biofilms. Annu Rev Microbiol 1995;49:711–745
- 22 Mack D, Becker P, Chatterjee I, et al. Functional molecules, regulatory circuits, and adaptive responses are mechanisms of biofilm formation in *Staphylococcus epidermidis* and *Staphylococcus aureus*. Int J Med Microbiol 2004;294:203–212
- 23 Ramage G, Mowat E, Jones B, Williams C, Lopez-Ribot J. Our current understanding of fungal biofilms. Crit Rev Microbiol 2009;35(04):340–355

- 24 Wei Q, Ma LZ. Biofilm matrix and its regulation in *Pseudomonas aeruginosa*. Int J Mol Sci 2013;14(10):20983–21005
- 25 Palková Z, Váchová L Life within a community: benefit to yeast long-term survival. FEMS Microbiol Rev 2006;30(05):806–824
- 26 Otter JA, Vickery K, Walker JT, et al. Surface-attached cells, biofilms and biocide susceptibility: implications for hospital cleaning and disinfection. J Hosp Infect 2015;89(01):16–27
- 27 Machado I, Graça J, Lopes H, Lopes S, Pereira MO. Antimicrobial pressure of ciprofloxacin and gentamicin on biofilm development by an endoscope-isolated *Pseudomonas aeruginosa*. ISRN Biotechnol 2012;2013:178646
- 28 Hall-Stoodley L, Costerton JW, Stoodley P. Bacterial biofilms: from the natural environment to infectious diseases. Nat Rev Microbiol 2004;2(02):95–108
- 29 Carpentier B, Cerf O. Biofilms and their consequences, with particular reference to hygiene in the food industry. J Appl Bacteriol 1993;75(06):499–511
- 30 Johnson KA. Osteomyelitis in dogs and cats. J Am Vet Med Assoc 1994;204(12):1882–1887
- 31 Siqueira EG, Rahal SC, Ribeiro MG, Paes AC, Listoni FP, Vassalo FG. Exogenous bacterial osteomyelitis in 52 dogs: a retrospective study of etiology and in vitro antimicrobial susceptibility profile (2000-2013). Vet Q 2014;34(04):201–204
- 32 Savicky R, Beale B, Murtaugh R, Swiderski-Hazlett J, Unis M. Outcome following removal of TPLO implants with surgical site infection. Vet Comp Orthop Traumatol 2013;26(04):260–265
- 33 Nicoll C, Singh A, Weese JS. Economic impact of tibial plateau leveling osteotomy surgical site infection in dogs. Vet Surg 2014; 43(08):899–902
- 34 Nazarali A, Singh A, Moens NMM, et al. Association between methicillin-resistant *Staphylococcus pseudintermedius* carriage and the development of surgical site infections following tibial plateau leveling osteotomy in dogs. J Am Vet Med Assoc 2015;247 (08):909–916
- 35 Heller S, Rezapoor M, Parvizi J, Jefferson T. Minimising the risk of infection: a peri-operative checklist. Bone Joint J 2016;98-B(1, Suppl A):18–22
- 36 Edmiston CE Jr, Leaper DJ. Intra-operative surgical irrigation of the surgical incision: what does the future hold saline, antibiotic agents, or antiseptic agents? Surg Infect (Larchmt) 2016;17(06): 656–664
- 37 Kavolus JJ, Schwarzkopf R, Rajaee SS, Chen AF. Irrigation fluids are used for the prevention and treatment of orthopedic infections. J Bone Joint Surg Am 2020;102(01):76–84
- 38 Siddiqi A, Abdo ZE, Springer BD, Chen AF. Pursuit of the ideal antiseptic irrigation solution in the management of periprosthetic joint infections. J Bone Jt Infect 2021;6(06):189–198
- 39 Siddiqi A, Abdo ZE, Rossman SR, et al. What is the optimal irrigation solution in management of periprosthetic hip and knee joint infections? J Arthroplasty 2021;36(10):3570–3583
- 40 Israel SK, Jaramillo E, Liska WD. Preclosure povidone-iodine lavage in total hip replacement surgery: infection outcomes and cost-benefit analysis. Vet Surg 2023;52(01):33–41
- 41 Brown NM, Cipriano CA, Moric M, Sporer SM, Della Valle CJ. Dilute betadine lavage before closure for the prevention of acute postoperative deep periprosthetic joint infection. J Arthroplasty 2012; 27(01):27–30
- 42 Calkins TE, Culvern C, Nam D, et al. Dilute betadine lavage reduces the risk of acute postoperative peri-prosthetic joint infection in aseptic revision total knee and hip arthroplasty: a randomized controlled trial. J Arthroplasty 2020;35(02):538–543.e1
- 43 van Meurs SJ, Gawlitta D, Heemstra KA, Poolman RW, Vogely HC, Kruyt MC. Selection of an optimal antiseptic solution for intraoperative irrigation: an in vitro study. J Bone Joint Surg Am 2014; 96(04):285–291
- 44 Goswami K, Austin MS. Intraoperative povidone-iodine irrigation for infection prevention. Arthroplast Today 2019b;5(03):306–308

- 45 Gilotra M, Nguyen T, Jaffe D, Sterling R. Dilute betadine lavage reduces implant-related bacterial burden in a rabbit knee prosthetic infection model. Am J Orthop 2015;44(02):E38–E41
- 46 Cichos KH, Andrews RM, Wolschendorf F, Narmore W, Mabry SE, Ghanem ES. Efficacy of intraoperative antiseptic techniques in the prevention of periprosthetic joint infection: superiority of betadine. J Arthroplasty 2019;34(7S):S312–S318
- 47 George J, Klika AK, Higuera CA. Use of chlorhexidine preparations in total joint arthroplasty. J Bone Jt Infect 2017;2(01):15–22
- 48 Penn-Barwell JG, Murray CK, Wenke JC. Comparison of the antimicrobial effect of chlorhexidine and saline for irrigating a contaminated open fracture model. J Orthop Trauma 2012;26 (12):728–732
- 49 Andriollo L, Sangaletti R, Velluto C, Perticarini L, Benazzo F, Rossi SMP. Impact of a novel antiseptic lavage solution on acute periprosthetic joint infection in hip and knee arthroplasty. J Clin Med 2024;13(11):3092
- 50 Premkumar A, Nishtala SN, Nguyen JT, Bostrom MPG, Carli AV. Comparing the efficacy irrigation solutions on Staphylococcal biofilm formed on arthroplasty surface. J Arthroplasty 2021; 36:26–32
- 51 Berríos-Torres SI, Umscheid CA, Bratzler DW, et al; Healthcare Infection Control Practices Advisory Committee. Centers for disease control and prevention guideline for surgical site infection. JAMA Surg 2017;152(08):784–791

- 52 Ernest EP, Machi AS, Karolcik BA, LaSala PR, Dietz MJ. Topical adjuvants incompletely remove adherent *Staphylococcus aureus* from implant materials. J Orthop Res 2018;36(06):1599–1604
- 53 Knecht CS, Moley JP, McGrath MS, Granger JF, Stoodley P, Dusane DH. Antibiotic loaded calcium sulfate bead and pulse lavage eradicates biofilms on metal implant materials in vitro. J Orthop Res 2018;36(09):2349–2354
- 54 Thompson AM, Bergh MS, Wang C, Wells K. Tibial plateau levelling osteotomy implant removal: a retrospective analysis of 129 cases. Vet Comp Orthop Traumatol 2011;24(06):450–456
- 55 Ingram LO. Ethanol tolerance in bacteria. Crit Rev Biotechnol 1990;9(04):305–319
- 56 Elzain AM, Elsanousi SM, Elfatih M, Ibrahim A. Effectiveness of ethanol and methanol alcohols on different isolates of *Staphylococcus* species. J Bacteriol Mycol Open Access. 2019;7:71–73
- 57 Eugster S, Schawalder P, Gaschen F, Boerlin P. A prospective study of postoperative surgical site infections in dogs and cats. Vet Surg 2004;33(05):542–550
- 58 Pratesi A, Moores AP, Downes C, Grierson J, Maddox TW. Efficacy of postoperative antimicrobial use for clean orthopedic implant surgery in dogs: a prospective randomized study in 100 consecutive cases. Vet Surg 2015;44(05):653–660
- 59 Márquez-Gómez M, Díaz-Navarro M, Visedo A, et al. An in vitro study to assess the best strategy for the chemical debridement of periprosthetic joint infection. Antibiotics (Basel) 2023;12(10):1507