N. SHIMADA, Y. ABE, S. YOKOSHIMA, T. FUKUYAMA\* (THE UNIVERSITY OF TOKYO, JAPAN)

Total Synthesis of (-)-Lycoposerramine-S Angew. Chem. Int. Ed. 2012, 51, 11824-11826.

## Total Synthesis of (-)-Lycoposerramine-S

Significance: Fukuyama and co-workers report the first total synthesis of the caged tetracyclic Lycopodium alkaloid (-)-lycoposerramine-S. The enantioselective synthesis is centered around an impressive 1,3-dipolar cycloaddition which diastereoselectively constructs the central pentasubstituted pyrrolidine ring utilizing a chiral morpholinone. A radical cyclization and alkylative ring closure of the nine-membered ring using a 4-nitrobenzenesulfonyl amide leads to the synthesis of the natural product in only 14 steps.

Comment: In a striking intramolecular 1,3-dipolar cycloaddition, condensation of aldehyde **D** with morpholinone E led to the diastereoselective formation of pyrrolidine **G** containing four newly constructed contiguous stereocenters in excellent yield. The formation of the 2,5-cis relationship is thought to arise from preferential formation of Z-azomethine ylide **F**. Exhaustive reduction, selective elimination of the resulting secondary alcohol followed by a radical annulation led to tricycle J. Finally, the medium-sized ring was assembled by use of alkylative nosyl amide chemistry previously developed by the Fukuyama group.

SYNFACTS Contributors: Erick M. Carreira, Simon Breitler Synfacts 2013, 9(1), 0007 Published online: 17.12.2012 DOI: 10.1055/s-0032-1317853; Reg-No.: C02512SF