

L. ZHOU, X. LIU, J. JI, Y. ZHANG, X. HU, L. LIN, X. FENG\* (SICHUAN UNIVERSITY, CHENGDU, P. R. OF CHINA)

Enantioselective Baeyer–Villiger Oxidation: Desymmetrization of *Meso* Cyclic Ketones and Kinetic Resolution of Racemic 2-Arylcyclohexanones

*J. Am. Chem. Soc.* **2012**, *134*, 17023–17026.

## Enantioselective Baeyer–Villiger Oxidation

Category

Metal-Catalyzed  
Asymmetric  
Synthesis and  
Stereoselective  
Reactions




Key words

Baeyer–Villiger  
oxidation


scandium

desymmetrization





### Desymmetrization of *meso*-cyclic ketones:






### Kinetic resolution of racemic cyclohexanones:



### Selected examples of desymmetrization:



### Selected examples of kinetic resolution:



**Significance:** The asymmetric Baeyer–Villiger oxidation of prochiral and racemic cyclic ketones effectively synthesized optically active  $\epsilon$ - and  $\gamma$ -lactones. The desymmetrization of racemic cyclohexanones interestingly showed a reversal of migratory aptitude with high levels of enantioselectivity.

**Comment:** The authors continued their use of chiral  $N,N'$ -dioxide-metal catalysts for the Baeyer–Villiger oxidation reaction. During the desymmetrization of *meso*-cyclohexanones and *meso*-cyclobutanones, the electronic and steric nature of the substituents appeared to have no effect on enantioselectivity; the opposite was true for the kinetic resolution of racemic cyclohexanones.

**SYNFACTS Contributors:** Hisashi Yamamoto, Kimberly Griffin  
Synfacts 2013, 9(1), 0049 Published online: 17.12.2012


DOI: 10.1055/s-0032-1317774; Reg-No.: H15412SF

2013 © THIEME STUTTGART • NEW YORK


D. BEST, S. KUJAWA, H. W. LAM\* (UNIVERSITY OF EDINBURGH, UK)

Diastereo- and Enantioselective Pd(II)-Catalyzed Additions of 2-Alkylazaarenes to *N*-Boc Imines and Nitroalkenes  
*J. Am. Chem. Soc.* **2012**, *134*, 18193–18196.

## Palladium-Catalyzed Asymmetric Addition of Alkylazaarenes to Imines and Nitroalkenes



### Selected examples:




**Significance:** While precedence of the direct addition of alkylazaarenes to imines and nitroalkenes in a racemic manner exists, the authors report the use of a chiral palladium(II)-bis(oxazoline) catalyst that can render this reaction highly diastereo- and enantioselective. The reaction proceeds under practical conditions, employing undried solvent at mild temperatures and under an air atmosphere.

**Comment:** The use of electron-withdrawing groups on the azaarene facilitates the deprotonation of the benzylic position at lower temperatures, which allows the catalyst to exert high stereocontrol. The corollary is, that the scope is limited to electron-poor azaarenes. However, the authors demonstrate the utility of these products with functionalization of the nitro group on the azaarenes. Treatment of the imine-addition products with mild acid readily deprotects the Boc group.


SYNFACTS Contributors: Mark Lautens, Lei Zhang  
 Synfacts 2013, 9(1), 0050 Published online: 17.12.2012  
 DOI: 10.1055/s-0032-1317750; Reg-No.: L15212SF

2013 © THIEME STUTTGART • NEW YORK


## Chiral Cp Ligands in Rhodium-Catalyzed Asymmetric C–H Functionalization



### Selected examples:



### Postulated model for stereoinduction:




**Significance:** A rhodium complex with a chiral Cp ligand that catalyzes an enantioselective synthesis of isoquinolones via a directed C–H bond functionalization is reported. Often, in half-sandwich transition-metal-catalyzed reactions, Cp remains the sole permanent ligand on the metal. Thus, despite the challenges, the development of chiral Cp ligands for inducing enantioselectivity is a powerful approach.

**Comment:** The highly effective Cp ligand reported is postulated to control the spatial orientation of the coupling partners. For instance, the ligand is  $C_2$ -symmetric to avoid diastereomeric coordination of the metal. The benzophenone ketal shields one face of the substrate and the equatorial methyl group pushes the bulky Boc group away. The controlled trajectory of the attacking alkene gives rise to the stereo-configuration of the product.

A. G. SMITH, H. M. L. DAVIES\* (EMORY UNIVERSITY, ATLANTA, USA)  
Rhodium-Catalyzed Enantioselective Vinylogous Addition of Enol Ethers to Vinylidiazooacetates  
*J. Am. Chem. Soc.* **2012**, *134*, 18241–18244.

## Rhodium-Catalyzed Asymmetric Vinylogous Addition to Vinylidiazooacetates




**Significance:** A rhodium-catalyzed asymmetric vinylogous addition of silyl enol ethers to siloxyvinylideneacetates is reported. Depending on the sterics of the substituents on the substrate, this method can access cyclopentenones **2** or alkynoates **3** with high yield and excellent enantioselectivity.

**Comment:** The use of (*Z*)-silyl enol ethers is critical in achieving the observed enantioselectivity. In the proposed mechanism, vinylogous adduct **5** can undergo a stereoselective 1,4-silyoxy shift to form **3**. Bulkier R<sup>1</sup> groups favor the aldol reaction to form formal [3+2] adduct **6**, which in one pot, in acid, can afford **2**.

A. K. MOURAD, J. LEUTZOW, C. CZEKELIUS\* (FREIE UNIVERSITÄT BERLIN, GERMANY AND FAYOUM UNIVERSITY, EGYPT)  
Anion-Induced Enantioselective Cyclization of Diynamides to Pyrrolidines Catalyzed by Cationic Gold Complexes  
*Angew. Chem. Int. Ed.* **2012**, *51*, 11149–11152.

## Cationic Gold-Catalyzed Cyclization of Diynamides



**Significance:** The authors report an enantioselective cycloisomerization of diynamides to methylene pyrrolidines catalyzed by cationic gold with optically active binol phosphates as counterions. This work was inspired by Toste and co-workers' application of chiral counterions in gold-catalyzed functionalization of allenes (*Science* **2007**, *317*, 462). The chiral pyrrolidine products formed are highly valuable as they contain an all-carbon-substituted quaternary stereocenter and are difficult to prepare in enantiomerically pure form by other conventional methods.

**Comment:** Czekelius and co-workers had previously demonstrated that cationic gold complexes cyclize diynols and diynamides to the corresponding unsaturated heterocycles in good yield (*Chem. Eur. J.* **2009**, *15*, 13323). However, optically active phosphine and carbene ligands gave poor enantioselectivity due to the linear coordination geometry in gold(I)-alkyne complexes. The use of commercially available binol phosphates as chiral counterions overcomes this problem and allows for high enantioselectivity in the cyclization. The best results were obtained in chlorinated solvents at low temperatures, which is in line with the contact ion pair model of the cationic gold-alkyne complex and the anionic chiral phosphate.

**SYNFACTS Contributors:** Mark Lautens, Jennifer Tsoung  
Synfacts 2013, 9(1), 0053 Published online: 17.12.2012  
DOI: 10.1055/s-0032-1317753; Reg-No.: L15512SF

2013 © THIEME STUTTGART • NEW YORK

Category

Metal-Catalyzed  
Asymmetric  
Synthesis and  
Stereoselective  
Reactions

Key words

gold




cyclo-isomerization

pyrrolidines








This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.

A. A. MIKHAILINE, M. I. MAISHAN, R. H. MORRIS\* (UNIVERSITY OF TORONTO, CANADA)  
Asymmetric Transfer Hydrogenation of Ketimines Using Well-Defined Iron(II)-Based Precatalysts Containing a PNNP Ligand  
*Org. Lett.* **2012**, *14*, 4638–4641.

## Iron-Catalyzed Asymmetric Transfer Hydrogenation of Ketimines



### Selected examples:



**Significance:** The authors report an iron-catalyzed asymmetric transfer hydrogenation under mild conditions that gives chiral amines with high enantioselectivity (94–99% ee). The system provides a solution to the challenging C=N bond reduction and proceeds with 2-propanol as the reducing agent.

**Comment:** Iron(II)–PNNP complexes that catalyze the asymmetric reduction of *N*-(diphenylphosphino)- and *N*-(4-tolylsulfonyl)ketimines were developed. The  $(R,R)$ -diamine catalyst produces the  $(S)$ -amine.  $(S,S)$ -**3** are found to be the most active and stereoselective catalyst. The reaction outcome is influenced mainly by the sterics around the imine carbon but is insensitive to its electronic character.

**SYNFACTS Contributors:** Hisashi Yamamoto, Lan Luo  
*Synfacts* 2013, 9(1), 0054 Published online: 17.12.2012  
**DOI:** 10.1055/s-0032-1317846; **Reg-No.:** H16512SF

2013 © THIEME STUTTGART • NEW YORK

## Palladium-Catalyzed Enantioselective Arylation of $\alpha$ -Imino Esters

Category

Metal-Catalyzed Asymmetric Synthesis and Stereoselective Reactions



Key words

palladium


asymmetric arylation

arylglycine derivatives


$\alpha$ -amino acids



### Substrate scope:



### Proposed transition state:



**Significance:** This protocol provides a practical and direct route to chiral arylglycines with high enantioselectivity (up to 99% ee). These derivatives can be easily converted into optically active  $\alpha$ -amino acids, which are commonly used as chiral auxiliaries in asymmetric catalysis.








**Comment:** A palladium(II)-catalyzed asymmetric arylation of *N*-aryl- $\alpha$ -imino esters using a chiral BOX ligand was developed. This method is applicable to various aromatic boronic acids. A stereochemical model, consistent with experimental results, suggests a *re*-face attack of the aryl group onto the *N*-arylimine carbon.

S. L. SHI, X. F. WEI, Y. SHIMIZU, M. KANAI\* (THE UNIVERSITY OF TOKYO AND JAPAN SCIENCE AND TECHNOLOGY AGENCY, KAWAGUCHI-SHI, JAPAN)


Copper(I)-Catalyzed Enantioselective Incorporation of Ketones to Cyclic Hemiaminals for the Synthesis of Versatile Alkaloid Precursors

*J. Am. Chem. Soc.* **2012**, *134*, 17019–17022.


## Copper-Catalyzed Enantioselective Incorporation of Ketones to Hemiaminals



### Application:



### Proposed mechanism:



**Significance:** The authors developed a copper-catalyzed enantioselective incorporation of ketones to cyclic hemiaminals. A series of hemiaminals, including five-, six- and seven-membered rings, were applicable to provide versatile alkaloid precursors in high yield with excellent enantioselectivity.

**SYNFACTS Contributors:** Hisashi Yamamoto, Susumu Oda  
Synfacts 2013, 9(1), 0056 Published online: 17.12.2012  
**DOI:** 10.1055/s-0032-1317777; **Reg-No.:** H15712SF

**Comment:** This reaction proceeds through three successive steps: aldol reaction, dehydration and intramolecular enantioselective aza-Michael reaction. Employment of this pathway contributed to improve the reaction conditions and expand the substrate scope. Synthetic utility was demonstrated by the preparation of alkaloid and drug precursors.



M. L. COOKE, K. XU, B. BREIT\* (ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG, GERMANY)  
Enantioselective Rhodium-Catalyzed Synthesis of Branched Allylic Amines by Intermolecular Hydroamination of Terminal Allenes  
*Angew. Chem, Int. Ed.* **2012**, *51*, 10876–10879.


# Rhodium-Catalyzed Enantioselective Hydroamination of Allenes

Category


Metal-Catalyzed  
Asymmetric  
Synthesis and  
Stereoselective  
Reactions

Key words


rhodium  
hydroamination  
allenes



## Selected examples:



## Proposed mechanism:




**Significance:** Despite the versatility of  $\alpha$ -chiral allylic amines, synthetic methods to access them have been underdeveloped. The authors reported the first example of the enantioselective intermolecular hydroamination of mono-substituted allenes.

**Comment:** A variety of substituted anilines, even bearing unprotected alcohol and indole moieties, were employed to give good yields and high enantioselectivities. Further mechanistic study is desirable to explain the regioselectivity of the hydro-metalation step.

**SYNFACTS Contributors:** Hisashi Yamamoto, Susumu Oda  
Synfacts 2013, 9(1), 0057 Published online: 17.12.2012  
**DOI:** 10.1055/s-0032-1317778; **Reg-No.:** H15812SF

K. HYODO, S. NAKAMURA,\* N. SHIBATA\* (NAGOYA INSTITUTE OF TECHNOLOGY, JAPAN)  
Enantioselective Aza-Morita-Baylis-Hillman Reactions of Acrylonitrile Catalyzed by Palladium(II) Pincer  
Complexes having  $C_2$ -Symmetric Chiral Bis(imidazoline) Ligands  
*Angew. Chem. Int. Ed.* **2012**, *51*, 10337–10341.

## Palladium-Catalyzed Enantioselective Aza-Morita-Baylis-Hillman Reaction





**Significance:** This paper describes the palladium-catalyzed enantioselective aza-Morita-Baylis-Hillman reaction of acrylonitriles with imines. The bulky pincer ligand enabled the synthesis of enantioenriched  $\alpha$ -methylene- $\beta$ -aminonitriles in high yield.

**Comment:** The palladium-pincer complex preferably activates acrylonitrile, even in the presence of ethyl acrylate. The palladium ketenimide is a key intermediate for the asymmetric induction. The palladium complex may promote other Lewis acid catalyzed reactions.


**SYNFACTS Contributors:** Hisashi Yamamoto, Susumu Oda  
*Synfacts* 2013, 9(1), 0058 Published online: 17.12.2012  
**DOI:** 10.1055/s-0032-1317779; **Reg-No.:** H15912SF

2013 © THIEME STUTTGART • NEW YORK

## Ruthenium-Catalyzed Asymmetric Hydrogenation of $\beta$ -Ketophosphonates



### Selected examples:




**Significance:** The current work represents an efficient protocol for the enantioselective hydrogenation of  $\beta$ -ketophosphonate derivatives catalyzed by a ruthenium-(S)-Sunphos complex. Good to excellent enantioselectivity and yield were obtained for a variety of substrates.

**Comment:** Hydroxyphosphonate motifs are known to be mimics of hydroxy carboxylic acids or amino acids. Given their medicinal importance, many synthetic methodologies have been developed. The protocol described herein was even used for the reduction of  $\alpha$ -substituted  $\beta$ -ketophosphonates, providing the desired products with good *syn* diastereoselectivity.


M. XU, T.-T. REN, C.-Y. LI\* (ZHEJIANG SCI-TECH UNIVERSITY, HANGZHOU, P. R. OF CHINA)

Gold-Catalyzed Oxidative Rearrangement of Homopropargylic Ether via Oxonium Ylide  
*Org. Lett.* **2012**, *14*, 4902–4905.


## Homopropargylic Ether Rearrangement via Gold Catalysis



**Selected substrate scope:**



**Proposed mechanism:**



**Significance:** Gold catalysis has emerged as a powerful platform to conduct complex organic transformations. Specifically, the implementation of gold carbenoids has shown great promise in synthetic planning. These useful intermediates offer a convenient alternative to generate metal carbenes which are traditionally obtained from diazo compounds. The authors utilize these intermediates to synthesize  $\alpha,\beta$ -unsaturated carbonyl compounds from homopropargylic ethers.


**Comment:** The authors report a silver-assisted gold(I)-catalyzed carbonyl synthesis. In an effort to obtain cyclobutanes **3** via a [1,2]-shift mechanism (path d), the authors unexpectedly obtained the corresponding  $\alpha,\beta$ -unsaturated carbonyl compounds **2**. Control experiments show that neither IMesAuCl, nor AgNTf<sub>2</sub> or HNTf<sub>2</sub> alone could catalyze the reaction. The scope of the reported reaction is quite broad; however, yields are generally moderate to good. In some instances cyclobutanones are obtained as the major product.

**SYNFACTS Contributors:** Mark Lautens, David A. Petrone  
Synfacts 2013, 9(1), 0060 Published online: 17.12.2012


**DOI:** 10.1055/s-0032-1317758; **Reg-No.:** L16012SF

B. M. TROST,\* P. J. MORRIS, S. J. SPRAGUE (STANFORD UNIVERSITY, USA)  
Palladium-Catalyzed Diastereo- and Enantioselective Formal [3 + 2]-Cycloadditions of Substituted  
Vinylcyclopropanes  
*J. Am. Chem. Soc.* **2012**, *134*, 17823–17831.

## Palladium-Catalyzed Asymmetric Formal [3+2] Cycloaddition



Proposed Curtin–Hammett reaction pathway:



**Significance:** A palladium-catalyzed asymmetric formal [3+2] cycloaddition of vinylcyclopropanes to electron-poor olefins is reported using the Trost ligand. The developed method can access highly substituted cyclopentanes with high diastereo- and enantioselectivity with moderate to high yield.

**Comment:** As the vinylcyclopropanes **1** used are racemic, the authors propose that the reaction occurs under Curtin–Hammett conditions for this stereo-convergent reaction. Notably, the effects of  $\pi$ – $\sigma$ – $\pi$  interconversion and the reversibility of the conjugate addition establish pre-equilibria of diastereomeric reactive intermediates **4** and **5**, consequently favoring the formation of **3**.

**SYNFACTS Contributors:** Mark Lautens, Lei Zhang  
*Synfacts* 2013, 9(1), 0061 Published online: 17.12.2012  
DOI: 10.1055/s-0032-1317751; Reg-No.: L15312SF

2013 © THIEME STUTTGART • NEW YORK

Category

Metal-Catalyzed  
Asymmetric  
Synthesis and  
Stereoselective  
Reactions

Key words

palladium

Trost ligand

vinylcyclopropanes


formal [3+2]  
cycloaddition

cyclopentanes


T. C. JOHNSON, W. G. TORRY, M. WILLS\* (THE UNIVERSITY OF WARWICK, COVENTRY, UK)  
Application of Ruthenium Complexes of Triazole-Containing Tridentate Ligands to Asymmetric Transfer  
Hydrogenation of Ketones  
*Org. Lett.* **2012**, *14*, 5230–5233.

## Asymmetric Ruthenium-Catalyzed Transfer Hydrogenation of Ketones

Overall transformation:



Selected substrate scope:



**Significance:** Transition-metal-catalyzed asymmetric transfer hydrogenation (ATH) has become a leading reduction method, which can be credited to its broad scope and relatively mild conditions. Additionally, the development of more general methods to synthesize chiral secondary alcohol is a useful endeavor. Specifically, the ATH reduction of *ortho*-substituted aryl ketones is considered a more challenging transformation than that of related *meta*- and *para*-substituted substrates.


**Comment:** The authors report a ruthenium-catalyzed ATH of substituted aryl methyl ketones using a novel tridentate triazole containing ligand. The scope of this transformation is quite broad, and conversions and enantioselectivities range from moderate to excellent. Notably, tetralone and 4-chromanone can be reduced efficiently with synthetically useful enantioselectivity. The reduction of cyclohexyl methyl ketone proceeds with excellent conversion, yet enantioselectivity remains low (13% ee).

**SYNFACTS Contributors:** Mark Lautens, David A. Petrone  
 Synfacts 2013, 9(1), 0062 Published online: 17.12.2012  
 DOI: 10.1055/s-0032-1317759; Reg-No.: L16112SF

2013 © THIEME STUTTGART • NEW YORK

# Mechanistic Study of Palladium-Catalyzed Wacker-Type Cyclizations

## Stereochemical pathways:




**Significance:** Recently, Stahl and co-workers had shown that a  $Pd(II)$  catalyst with a chiral pyridine–oxazoline (pyrox) ligand allowed preparation of pyrrolidines in high yield and enantioselectivity (*Org. Lett.* 2011, 13, 2830). In the enantioselective cyclization of  $\gamma$ -alkenyl tosylamides, the anionic ligand (TFA vs OAc) was found to have a significant impact on the reaction outcome, where the use of  $[Pd(pyrox)(OAc)_2]$  gave significantly diminished yield and enantioselectivity. Through a series of mechanistic investigations with a chiral, deuterated substrate probe, the authors showed the significant effect the anionic ligand has in selecting the nucleopalladation (NP) pathway of the Wacker-type cyclization, which in course determines the ancillary neutral donor's ability to alter the stereochemical course of the pathway. This data provides the first direct correlation between NP stereoselectivity and the enantioselectivity of the transformation.

**Comment:** By using  $^1H$  NMR spectroscopy and HPLC analyses to determine H/D ratios and enantiomeric excesses, the authors were able to determine the yields of the four possible products from the reaction of a deuterated acyclic substrate under different conditions (see above). They showed that only in the *trans* pathway does the pyrox ligand play a significant role, thus the *trans*-amido-palladation (AP) pathway proceeds with high enantioselectivity, while the *cis*-AP pathway exhibits low enantioselectivity. The authors suggest that the carboxylate ligand acts as a Brønsted base to mediate  $Pd$ -amidate bond formation in the *cis*-AP pathway, whereas the TFA anionic ligand is substituted by the substrate alkene and favors the *trans*-AP pathway.


Y.-L. SHAO, X.-H. ZHANG,\* J.-S. HAN, P. ZHONG\* (WENZHOU UNIVERSITY, P. R. OF CHINA)  
Pd(II)-Catalyzed Dehydrogenative Olefination of Terminal Arylalkynes with Allylic Ethers: General and Selective Access to Linear (*Z*)-1,3-Enynes  
*Org. Lett.* **2012**, *14*, 5242–5245.

## Access to 1,3-Enynes by Pd(II)-Catalyzed Dehydrogenative Olefination

Overall transformation:



Selected substrate scope:



**Significance:** 1,3-enynes are important motifs found in pharmaceutically active compounds and natural products. For this reason, efficient methods which easily access these structures are desirable to synthetic chemists. Despite advances made using copper and iron catalysis, which commonly require alkene pre-activation, palladium-catalyzed dehydrogenative cross-coupling has shown promise as a more benign strategy in this regard.



**Comment:** The authors report the first example of a Pd(OAc)<sub>2</sub>-catalyzed direct dehydrogenative olefination of terminal aryl alkynes and allylic ethers to exclusively access (*Z*)-1,3-enyne derivatives. The reaction exhibits good scope with respect to arylalkynes, however, only allylic ethers and thioethers were used as coupling partners, thus limiting the applicability. Nonetheless, this method appears to be an interesting application of dehydrogenative cross-coupling which accesses these important compounds in a step-efficient manner.

**SYNFACTS Contributors:** Mark Lautens, David A. Petrone  
Synfacts 2013, 9(1), 0064 Published online: 17.12.2012  
**DOI:** 10.1055/s-0032-1317763; **Reg-No.:** L16312SF




2013 © THIEME STUTTGART • NEW YORK




# Artificial Rh(III)–Metalloenzyme-Catalyzed Asymmetric C–H Activation



## Selected examples:



## Streptavidin with engineered carboxylate mutation:



**Significance:** A highly active, artificial rhodium(III) metalloenzyme that catalyzes an asymmetric synthesis of dihydroisoquinolones through C–H activation is reported. A biotinylated rhodium(III) complex is successfully incorporated into streptavidin. With active-site mutagenesis, the engineered enzyme displayed up to 100-fold reaction rate increase compared to the activity of the unbound rhodium complex.


**Comment:** As Cp is the only permanently bound ligand on rhodium in the catalytic cycle, it has been difficult to render this reaction enantioselective until recently. This report provides an alternative solution for this problem. Based on the concerted metalation–deprotonation mechanism, the authors used docking modeling and introduced a basic carboxylate moiety in the active site. With kinetic isotope effect experiments, the importance of this mutation in accelerating the catalysis is demonstrated.

A. R. BURNS, J. S. GONZÁLEZ, H. W. LAM\* (UNIVERSITY OF EDINBURGH, UK)

Enantioselective Copper(I)-Catalyzed Borylative Aldol Cyclizations of Enone Diones

*Angew. Chem. Int. Ed.* **2012**, *51*, 10827–10831.

# Enantioselective Copper-Catalyzed Borylative Aldol Cyclizations



**Significance:** The formation of metal enolates allows for precise enolization, as well as potential enantio- and diastereoselective enolization. In this report, the authors apply this idea to a copper-catalyzed conjugate boration–aldol cyclization sequence to produce enantioenriched decalin-, hydrindane- and diquinone-based products.

**Comment:** The copper–bisphosphine catalyst system developed, produces decalins as well as [5,6]-, [6,5]-, and [5,5]-bicyclic ring products with high levels of diastereo- and enantioselectivity. Kinetic resolution of a racemic chiral enone also afforded the cyclization product with good diastereo- and enantioselectivity.

**SYNFACTS Contributors:** Hisashi Yamamoto, Kimberly Griffin



Synfacts 2013, 9(1), 0066 Published online: 17.12.2012

DOI: 10.1055/s-0032-1317775; Reg-No.: H15512SF


2013 © THIEME STUTTGART • NEW YORK

## In Situ Aldehyde Enolate Formation by Rhodium-Catalyzed Isomerization


### Isomerization–cross-aldol reaction with triallylboranes:



### Isomerization–cross aldol reaction with allylic alcohols:



### Isomerization–cross aldol with homoallyloxyborane and homoallylic alcohol:



**Significance:** Aldol reactions in which the aldol donor is derived from an aldehyde, are particularly challenging. This report describes a strategy in which aldehyde enolates are generated in situ by rhodium-catalyzed isomerization of triallylboroxanes. High *syn*-selectivity is obtained for a variety of aldehyde-donor and -acceptor partners.

**Comment:** Remarkably, the use of triallyloxyboranes is not required; simple primary and secondary allylic alcohols also undergo the isomerization–cross-aldol sequence with similar levels of reactivity and selectivity, presumably through a rhodium-enolate or -enol mechanism.


T. J. HARRISON, P. M. A. RABBAT, J. L. LEIGHTON\* (COLUMBIA UNIVERSITY, NEW YORK, USA)

An 'Aprotic' Tamao Oxidation/*Syn*-Selective Tautomerization Reaction for the Efficient Synthesis of the C(1)–C(9) Fragment of Fludelone


*Org. Lett.* **2012**, *14*, 4890–4893.

## A Rhodium(I)-Catalyzed Silylformylation–Crotosilylation–Tamao Oxidation

Overall transformation:



Reaction sequence:



**Significance:** Access to complex polyketide fragments typically consists of complex stepwise syntheses. Recent advances, including asymmetric crotylation and aldol cascades, have allowed chemists to synthesize extremely complex polyketide fragments with good step- and redox-economy, as well as minimal use of protecting groups. In this regard, silylformylation and silylcrotylation have emerged as complementary methods towards this end.

**SYNFACTS Contributors:** Mark Lautens, David A. Petrone  
Synfacts 2013, 9(1), 0068 Published online: 17.12.2012


**DOI:** 10.1055/s-0032-1317757; **Reg-No.:** L15912SF

**Comment:** The authors report the synthesis of the C1–C9 fragment of fludelone, a polyketide natural product. The authors elegantly utilize their silylformylation–crotosilylation chemistry (*J. Am. Chem. Soc.* **2000**, *122*, 8587) in conjunction with this newly developed aprotic Tamao oxidation–diastereoselective tautomerization methodology to access this ketone containing four stereocenters, three of which are contiguous.


C.-H. WEI, S. MANNATHAN, C.-H. CHENG\* (NATIONAL TSING HUA UNIVERSITY, HSINCHU, TAIWAN)

Regio- and Enantioselective Cobalt-Catalyzed Reductive [3+2] Cycloaddition Reaction of Alkynes with Cyclic Enones: A Route to Bicyclic Tertiary Alcohols  
*Angew. Chem. Int. Ed.* **2012**, *51*, 10592–10595.


## Cobalt-Catalyzed [3+2] Cycloaddition of Alkynes with Cyclic Enones




### Selected examples:




71% yield, 90% ee




76% yield, 77% ee




63% yield, 90% ee




69% yield, 78% ee



54% yield, 93% ee



53% yield, >99% ee



**Significance:** Cheng and co-workers describe a cobalt-catalyzed [3+2]-cycloaddition reaction that provides an atom-economic method for the synthesis of bicyclic tertiary alcohols from alkynes and cyclic enones with regioselectivity. During their previous studies of enantioselective reductive coupling of alkynes with cyclic enones to synthesize  $\beta$ -substituted ketones **1**, they found that the use of a  $\text{CoBr}_2/\text{dppe-Mn-ZnCl}_2$  system gave the bicyclic product **2** instead in high yield. With the use of a chiral ligand such as Duanphos, moderate to high enantioselectivity was also obtained.

**Comment:** This reported system is remarkable in that it allows for the reductive cycloaddition of various alkynes and cyclic enones to occur with good regio- and stereoselectivity using an air-stable cobalt catalyst, a mild reducing agent and water as the hydrogen source. Unsymmetrical alkynes also undergo reductive cycloaddition with good to high regioselectivity, though terminal alkynes and silyl-protected alkynes were unsuitable.

**SYNFACTS Contributors:** Mark Lautens, Jennifer Tsoung  
Synfacts 2013, 9(1), 0069 Published online: 17.12.2012  
**DOI:** 10.1055/s-0032-1317752; **Reg-No.:** L15412SF

2013 © THIEME STUTTGART • NEW YORK

Category

Metal-Catalyzed  
Asymmetric  
Synthesis and  
Stereoselective  
Reactions

Key words

cobalt

cycloaddition



tertiary alcohols

P. HE, X. LI, H. ZHENG, W. LI, L. LIN, X. FENG\* (SICHUAN UNIVERSITY, CHENGDU, P. R OF CHINA)







Asymmetric 1,2-Reduction of Enones with Potassium Borohydride Catalyzed by Chiral *N,N'*-Dioxide–Scandium(III) Complexes

*Org. Lett.* **2012**, *14*, 5134–5137.

## Scandium-Catalyzed Asymmetric Reduction with Potassium Borohydride



### Selected examples:



**Significance:** As an extension on previous work using chiral *N,N'*-dioxide–metal complexes for asymmetric catalysis (see Review), the authors now describe the scandium-catalyzed asymmetric reaction of enones and ketones with  $\text{KBH}_4$ . The resulting chiral alcohols are obtained with good yield and enantioselectivity.

**Review:** X. Liu, L. Lin, X. Feng *Acc. Chem. Res.* **2011**, *44*, 574–587.

**Comment:** Chiral allylic alcohols are important motifs widely present in natural products and biologically active molecules. The enantioselective reduction of enones is known as the most straightforward access to such motifs. Herein, the first example of catalytic enantioselective reduction of enones and ketones by using  $\text{KBH}_4$  is reported. The utilization of an aqueous solution of  $\text{KBH}_4$  was found to be crucial for obtaining high yield and enantioselectivity as the presence of water is believed to benefit proton transfer to accelerate the catalytic cycle. In this case, the reaction was performed in a homogeneous catalyst system. The HRMS spectra experiments indicated that the initial reducing species is  $\text{KBH}_3\text{OH}$ .

**SYNFACTS Contributors:** Hisashi Yamamoto, Jiajing Tan  
Synfacts 2013, 9(1), 0070 Published online: 17.12.2012


**DOI:** 10.1055/s-0032-1317785; **Reg-No.:** H16212SF

2013 © THIEME STUTTGART • NEW YORK


N. GAO, X.-W. GUO, S.-C. ZHENG, W.-K. YANG, X.-M. ZHAO\* (TONGJI UNIVERSITY, SHANGHAI AND DALIAN UNIVERSITY OF TECHNOLOGY, P. R OF CHINA)  
 Iridium-Catalyzed Enantioselective Allylation of Sodium 2-Aminobenzenethiolate: An Access to Chiral Benzo-Fused N,S-Heterocycles  
*Tetrahedron* **2012**, *68*, 9413–9418.

## Enantioselective Iridium(I)-Catalyzed Allylation of Sodium 2-Aminobenzenethiolates

Overall transformation:



Selected substrate scope:



**Significance:** Iridium-catalyzed enantioselective allylation has emerged as a powerful method to synthesize structurally diverse, chiral molecules. Despite much progress in the area of enantioselective carbon–sulfur bond formation using iridium, there have been no reports on the use of sodium 2-aminobenzenethiolate as a nucleophile in this class of reaction. Despite the potential of this substrate class to encounter detrimental ‘*ortho*-substituent effects’ on stereoselectivity, Zhao accomplishes selective and highly enantioselective S-allylation.


**SYNFACTS Contributors:** Mark Lautens, David A. Petrone  
*Synfacts* 2013, 9(1), 0071 Published online: 17.12.2012  
 DOI: 10.1055/s-0032-1317762; Reg-No.: L16212SF

**Comment:** The authors report an iridium-catalyzed asymmetric S-allylation reaction using chiral phosphoramidite ligands. The method is highly regio- and enantioselective for a variety of aryl- and alkyl-substituted allyl carbonates. Yields range from moderate to good with excellent enantiocontrol. In most cases, the authors are able to completely inhibit bisallylation and maintain high levels of branched-to-linear selectivity. The authors use the products to synthesize enantioenriched N,S-heterocycles via an N-allylation/ring-closing metathesis sequence.

G. I. MCGREW, C. STANCIU, J. ZHANG, P. J. CARROLL, S. D. DREHER,\* P. J. WALSH\*  
(UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA AND MERCK AND CO. INC., RAHWAY,  
USA)

Asymmetric Cross-Coupling of Aryl Triflates to the Benzylic Position of Benzylamines  
*Angew. Chem. Int. Ed.* **2012**, *51*, 11510–11513.

## Palladium-Catalyzed Direct Arylation of Chromium-Activated Benzylic C–H Groups



**Significance:** The authors previously described the application of  $(\eta^6\text{-C}_6\text{H}_5\text{CH}_2\text{R})\text{Cr}(\text{CO})_3$  complexes as nucleophile precursors in Pd-catalyzed allylic substitution reactions (*J. Am. Chem. Soc.* **2011**, *133*, 20552). They now report the first catalytic asymmetric cross-coupling of benzyllithiums  $\alpha$  to tertiary amines using  $[\text{Cr}(\text{CO})_3]$  activation of benzylic C–H bonds. The stabilized organolithium undergoes Pd-catalyzed coupling with aryl triflates by dynamic kinetic resolution to yield enantioenriched Cr-coordinated diarylmethylamines in good to high yield, which can be de-complexed by exposure to sunlight and air.

**Comment:** Development of an enantioselective version of the previously reported transformation is challenging as it requires the enantioenriched palladium catalyst to select for one of the chromium adducts faster than the other, and also requires the products to be impervious to racemization. High-throughput screening identified the chiral ligand Cy-Mandyphos, and that the addition of PMDETA and toluene as co-solvents increased the yield. The authors report future plans to close the catalytic cycle by focusing on an arene exchange between the chromium-complexed product and the free arene to liberate the product and regenerate the substrate.

**SYNFACTS Contributors:** Mark Lautens, Jennifer Tsoung  
Synfacts 2013, 9(1), 0072 Published online: 17.12.2012  
**DOI:** 10.1055/s-0032-1317756; **Reg-No.:** L15812SF




## Erratum

## Palladium-Catalyzed Direct Arylation of Chromium-Activated Benzylic C–H Groups

G. I. McGrew, C. Stanciu, J. Zhang, P. J. Carroll, S. D. Dreher,\* P. J. Walsh\* *Synfacts* **2013**, *9*, 72.

The keywords were incorrect. The correct keywords are palladium, enantioselective cross-coupling, diarylmethylamines. In addition, in the proposed mechanism, the two structures on the right should not contain palladium. The correct scheme is shown below. We apologize for this mistake.



J. T. BINDER, C. J. CORDIER, G. C. FU\* (MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE AND CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA, USA)  
 Catalytic Enantioselective Cross-Couplings of Secondary Alkyl Electrophiles with Secondary Alkylmetal Nucleophiles: Negishi Reactions of Racemic Benzylic Bromides with Achiral Alkylzinc Reagents  
*J. Am. Chem. Soc.* **2012**, *134*, 17003–17006.

Category

Metal-Catalyzed Asymmetric Synthesis and Stereoselective Reactions



Key words

nickel


Negishi coupling

oxazoline ligands


## Negishi Reaction of Racemic Benzylic Bromides and Alkylzinc Reagents




Selected examples:



For acyclic alkylzinc reagents:



Proposed mechanism:



**Significance:** Reported here is an enantioselective cross-coupling of racemic benzylic bromides with achiral alkylzinc reagents. A novel bidentate oxazoline-type ligand was developed, leading to the desired products in good yield and enantioselectivity.

**Comment:** It is surprising that both reagents are achiral. For acyclic alkylzinc reagents, an usual isomerization was observed and a substantial amount of a branched product was generated from an unbranched nucleophile.


**SYNFACTS Contributors:** Hisashi Yamamoto, Jiajing Tan  
 Synfacts 2013, 9(1), 0073 Published online: 17.12.2012  
 DOI: 10.1055/s-0032-1317780; Reg-No.: H16012SF

2013 © THIEME STUTTGART • NEW YORK


I. CANO, E. GÓMEZ-BENGOA, A. LANDA, M. MAESTRO, A. MIELGO, I. OLAIZOLA, M. OIARBIDE, C. PALOMO\* (UNIVERSIDAD DEL PAÍS VASCO, SAN SEBASTIÁN AND UNIVERSIDADE DA CORUÑA, SPAIN)

*N*-(Diazoacetyl)oxazolidin-2-thiones as Sulfur-Donor Reagents: Asymmetric Synthesis of Thiiranes from Aldehydes  
*Angew. Chem. Int. Ed.* **2012**, *51*, 10856–10860.


## Asymmetric Synthesis of $\alpha,\beta$ -Thioepoxy Carbonyls by Rhodium Catalysis




Selected examples:



Synthetic transformations of thirane products:



Proposed reaction pathway:




**Significance:** Stereoselective formation of C–S bonds is a difficult yet important challenge. This report describes the use of diazo thianes as intramolecular sulfur-donor reagents. Under rhodium catalysis, reaction with aldehydes forms thiiranes with high selectivity.

**SYNFACTS Contributors:** Hisashi Yamamoto, Patrick Brady  
*Synfacts* 2013, 9(1), 0074 Published online: 17.12.2012  
**DOI:** 10.1055/s-0032-1317773; **Reg-No.:** H15312SF


**Comment:** Computational studies indicate formation of thiocarbonyl ylide intermediate **A**. Reaction with an aldehyde yields a tricyclic adduct, with preferential formation of *anti,exo*-product **B** by 0.8–1.2 kcal/mol, which collapses to the *cis* product by an  $\text{S}_{\text{N}}2$  reaction. However, when the aryl substituent is anisyl, the *trans* product forms by an  $\text{S}_{\text{N}}1$  mechanism.

X. LIN, F. ZHENG, F.-L. QING\* (SHANGHAI INSTITUTE OF ORGANIC CHEMISTRY AND DONGHUA UNIVERSITY, SHANGHAI, P. R. OF CHINA)  
 Regio- and Diastereoselective Nickel-Catalyzed Allylation of Aromatic Aldehydes with  $\alpha$ -Halo- $\beta,\beta$ -difluoropropene Derivatives  
*J. Org. Chem.* **2012**, *77*, 8696–8704.


## Nickel-Catalyzed Synthesis of $\gamma$ -Fluorinated Homoallylic Alcohols




### Selected examples:




X = I, 60% yield, dr = 8:1



X = Br, 54% yield, dr = 3.5:1




X = Br, 66% yield, dr = 11:1



X = Br, 56% yield, dr = 8:1

### Proposed mechanism:



**Significance:** Functionalized fluoro olefins have been synthetic targets due to the ability of fluorine to alter the biological activity of organic compounds. In response to the high demand of fluorinated olefins, the authors developed a nickel-catalyzed reductive coupling of fluorinated dienes and carbonyl compounds to synthesize fluoro olefinic alcohols.




**Comment:** Both electron-rich and electron-deficient aromatic aldehydes undergo allylation, albeit with lower regioselectivity for electron-deficient aldehydes. The authors rationalize the *Z/E*-selectivity by the coordination ability of the aldehyde to  $\text{ZnCl}_2$ : for electron-rich aldehydes, the coupling reaction proceeds faster than diene isomerization, and the *Z/E*-ratio remains unchanged in the product.

**SYNFACTS Contributors:** Hisashi Yamamoto, Kimberly Griffin  
 Synfacts 2013, 9(1), 0075 Published online: 17.12.2012  
 DOI: 10.1055/s-0032-1317776; Reg-No.: H15612SF


S. K. RAY, P. K. SINGH, N. MOLLETI, V. K. SINGH\* (INDIAN INSTITUTE OF TECHNOLOGY KANPUR AND INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH BHOPAL, INDIA)

Enantioselective Synthesis of Coumarin Derivatives by PYBOX–DIPH–Zn(II) Complex Catalyzed Michael Reaction  
*J. Org. Chem.* **2012**, *77*, 8802–8808.


## Zinc-Catalyzed Synthesis of Coumarin Derivatives by Asymmetric Michael Reaction




### Selected examples:




R<sup>2</sup> = Ph      99% yield, 88% ee  
R<sup>2</sup> = 2-O<sub>2</sub>NC<sub>6</sub>H<sub>4</sub>      95% yield, 91% ee  
R<sup>2</sup> = 1-Naph      95% yield, 53% ee




97% yield, 85% ee




93% yield, 80% ee



93% yield, 97% ee



87% yield, 70% ee



90% yield, 96% ee

**Significance:** Coumarin derivatives are a broad class of biologically interesting molecules. The zinc-catalyzed system presented provides an efficient access to the direct precursors of such compounds with excellent yield (up to 99%) and enantioselectivity (up to 97%).

**Comment:** The authors report a PYBOX–DIPH–Zn(II) catalyzed asymmetric Michael reaction and its successful application in the synthesis of coumarin derivatives. This method can tolerate a wide range of cyclic 1,3-dicarbonyl compounds. The resulting products can be easily converted into bioactive molecules such as warfarin and acenocoumarol without loss of enantiopurity.

**SYNFACTS Contributors:** Hisashi Yamamoto, Lan Luo  
Synfacts 2013, 9(1), 0076    Published online: 17.12.2012

**DOI:** 10.1055/s-0032-1317844; **Reg-No.:** H16312SF



2013 © THIEME STUTTGART • NEW YORK

E. HERNANDO, R. G. ARRAYÁS,\* J. C. CARRETERO\* (UNIVERSIDAD AUTÓNOMA DE MADRID, SPAIN)


Catalytic Asymmetric Mannich Reaction of Glycine Schiff Bases with  $\alpha$ -Amido Sulfones as Precursors of Aliphatic Imines

*Chem. Commun.* **2012**, 48, 9622–9624.


## Copper-Catalyzed Asymmetric Mannich Reaction of Glycine Imines



### Selected examples:



### Proposed transition-state model:




**Significance:**  $\alpha,\beta$ -Diamino acids are valuable due to their presence in peptide-based drugs and other bioactive compounds. In this report, the authors have extended their copper-catalyzed Mannich reaction of glycine Schiff bases to imines derived from aliphatic aldehydes, which previously performed poorly.

**Comment:**  $\alpha$ -Amido sulfones are employed as imine precursors, due to the instability of imines derived from aliphatic aldehydes. Excellent enantioselectivity and *syn*-selectivity is obtained for a variety of imines. The products have high synthetic applicability due to the orthogonal protection of the amines.


**SYNFACTS Contributors:** Hisashi Yamamoto, Patrick Brady  
*Synfacts* 2013, 9(1), 0077 Published online: 17.12.2012  
**DOI:** 10.1055/s-0032-1317771; **Reg-No.:** H15112SF

L. NICOLAS, P. ANGIBAUD, I. STANSFIELD, P. BONNET, L. MEERPOEL, S. REYMOND,\*  
J. COSSY\* (ESCI PARISTECH, JANSSEN RESEARCH & DEVELOPMENT, VAL DE REUIL,  
FRANCE AND BEERSE, BELGIUM)  
Diastereoselective Metal-Catalyzed Synthesis of C-Aryl and C-Vinyl Glycosides  
*Angew. Chem. Int. Ed.* **2012**, *51*, 11101–11104.


## Cobalt-Catalyzed Cross-Coupling of 1-Bromo Glycosides and Grignard Reagents



### Selected examples:



### Mechanistic study:



**Significance:** Numerous metal-catalyzed cross-coupling methods to form anomeric C–C bonds exist, which are important for the synthesis of carbohydrate analogues such as C-glycosides (see Review below). However,  $\beta$ -elimination is a major drawback of these reactions. The authors report a new diastereoselective cobalt-catalyzed cross-coupling between 1-bromo glycosides and aryl and alkenyl Grignard reagents with moderate to good  $\alpha$ -selectivity.

**Review:** L. Somsák *Chem. Rev.* **2001**, *101*, 81–136.

**SYNFACTS Contributors:** Mark Lautens, Jennifer Tsoung  
Synfacts 2013, 9(1), 0078 Published online: 17.12.2012  
**DOI:** 10.1055/s-0032-1317754; **Reg-No.:** L15612SF

**Comment:** The authors report that there was good  $\alpha$ -selectivity for the cross-coupling reaction with mannose and galactose derivatives, but lower  $\alpha/\beta$  ratios for glucose derivatives. Like most cobalt-catalyzed cross-coupling reactions, the stereoselectivity of this reaction supports a radical pathway. Treatment of a  $\delta$ -olefinic 1-bromoglycoside produced an epimeric mixture of the bicyclic product, which would result from the formation of an anomeric radical that leads to a 5-*exo*-trig cyclization followed by cross-coupling with PhMgBr.