

Erratum

Risk of bleeding related to selective and non-selective serotonergic antidepressants: a case/non-case approach using data from two pharmacovigilance databases

¹Maximilian Gahr, ¹René Zeiss, ²Dirk Lang, ¹Bernhard J. Connemann, ³Christoph Hiemke, ¹Roland W. Freudenmann, ¹Carlos Schönfeldt-Lecuona

¹University of Ulm, Department of Psychiatry and Psychotherapy III, Leimgrubenweg 12-14, 89075 Ulm, Ulm, Germany

²University of Ulm, Department of Psychosomatic Medicine and Psychotherapy, Albert-Einstein-Allee 23, 89081 Ulm, Germany

³University Medical Center of Mainz, Department of Psychiatry and Psychotherapy, Untere Zahlbacherstr. 8, 55131 Mainz, Mainz, Germany

Pharmacopsychiatry 2014; 47

DOI: 10.1055/s-0034-1394398

Erratum:

In the abstract, wrong results were erroneously presented for ASS and diclofenac. This error only concerns the reporting odds ratios (ROR) and confidence intervals (CI) of ASS and diclofenac calculated for the BfArM/AkdÄ dataset; ROR and CI calculated for the WHO dataset were correctly presented for both substances in the abstract. Correct results (ROR and CI) for both substances are displayed in table 2 of the article.

The correct results for ASS (to be considered in the abstract) are: BfArM/AkdÄ, ROR 13.71 [95% CI 12.82–14.66] and for diclofenac: BfArM/AkdÄ, ROR 2.80 [95% CI 2.59–3.03].