


H.-H. LU, M. D. MARTINEZ, R. A. SHENVI* (THE SCRIPPS RESEARCH INSTITUTE, LA JOLLA, USA)

An Eight-Step Gram-Scale Synthesis of (-)-Jiadifenolide
Nat. Chem. **2015**, *7*, 604–607.

Total Synthesis of (-)-Jiadifenolide



Significance: Jiadifenolide is one of a number of secondary metabolites with neurotrophic activity isolated from the *Illicium* genus. The authors present a short route towards this natural product relying on a diastereoselective coupling of two butenolides. The strategy might allow for the synthesis of structurally related neurotrophic natural products.

Comment: Chiral bicycle **D** was prepared from (+)-citronellal via a short dehydration–ozonolysis–hetero-Pauson–Khand sequence. The lithium dienolate derived from **D** was diastereoselectively coupled with building block **F** to give the entire carbon skeleton **G** upon treatment with excess LDA and Ti(Oi-Pr)_4 . Four more steps allowed the isolation of considerable amounts of jiadifenolide.

SYNFACTS Contributors: Erick M. Carreira, Matthias Westphal
Synfacts 2015, 11(9), 0897 Published online: 18.08.2015
DOI: 10.1055/s-0034-1378755; Reg-No.: C04315SF