

In vivo imaging of duodenal follicular lymphoma with confocal laser endomicroscopy



Fig. 1 Conventional endoscopic image. Multiple small whitish polyps were identified in the descending part of the duodenum in a 70-year-old woman.



Fig. 2 Magnifying endoscopy combined with narrow-band imaging. The villi were fused and enlarged compared to the surrounding normal villi. Curled vascular loops were present in the enlarged villi.




Fig. 3 Probe-based confocal laser endomicroscopy (pCLE) imaging. **a** pCLE revealed the villiform architecture. The epithelium was uniformly bright and goblet cells were dark (arrows). **b** Numerous small cells with a size of about 5–8 μm were present.




Fig. 4 Histological findings of the biopsy specimens. A section stained with hematoxylin–eosin revealed a subepithelial nodular proliferation of small- to medium-sized atypical lymphoid cells (original magnification $\times 40$).

Follicular lymphoma arising in the duodenum is a distinct, relatively rare disease [1,2]. Here, we report our experience with confocal laser endomicroscopy (CLE) for follicular lymphoma of the duodenum with the corresponding histopathologic images, which, to our knowledge, is the first description of this.

The patient was a 70-year-old woman in whom multiple small whitish polyps were noted in the descending part of the duodenum on upper gastrointestinal endoscopic screening (Fig. 1). When the polypoid lesion was observed using narrow-band imaging, the villi were fused and enlarged compared to the surrounding normal villi (Fig. 2). Curled vascular loops were present in the enlarged villi. Subsequently, fluorescein-dripping CLE [3] was performed using probe-based confocal laser endomicroscopy (pCLE) (GastroFlex UHD, Cellvizio; Mauna Kea Technologies, Paris, France).

First, pCLE images of normal mucosa near the lesion were observed as a control, and revealed the villiform architecture with uniformly bright tall-columnar epithelium and dark goblet cells (Fig. 3a, arrows). Then the lesion was observed using pCLE. The width of each villus had increased compared to that of the normal villus structure, and numerous bright small cells were present (Fig. 3b).

Biopsy specimens of the lesion site revealed a subepithelial nodular proliferation of small- to medium-sized atypical lymphoid cells (Fig. 4), which were immunohistochemically positive for CD20, CD79a, CD10, and BCL-2, and negative for CD3, CD5, and cyclin D1, resulting in the pathological diagnosis of follicular lymphoma. The tumor cell size in the histological section was consistent with the size of the bright cells in the pCLE image. On positron emission tomography-computed tomography (PET-CT), no accumulation was noted in any regions of the body including the duodenum, and no involvement of the bone marrow was revealed by biopsy of the bone marrow. Based on these findings, the patient was diagnosed as stage I follicular lymphoma. With sufficient informed consent from the patient, a treatment strategy of “watchful waiting” was adopted.

Endoscopy_UCTN_Code_CCL_1AB_2AZ_3AB

Competing interests: None

**Kouichi Nonaka¹, Ken Ohata¹,
Shinichi Ban², Maiko Takita¹,
Yasushi Matsuyama¹, Tomoaki
Tashima¹, Yohei Minato¹,
Nobuyuki Matsuhashi¹**

¹ Department of Gastroenterology, NTT
Medical Center Tokyo, Tokyo, Japan

² Department of Pathology, Saiseikai
Kawaguchi General Hospital, Kawaguchi,
Japan

References

- 1 *Nadal E, Martinez A, Jimenez M et al.* Primary follicular lymphoma arising in the ampulla of Vater. *Ann Hematol* 2002; 81: 228–231
- 2 *Yoshino T, Miyake K, Ichimura K et al.* Increased incidence of follicular lymphoma in the duodenum. *Am J Surg Pathol* 2000; 24: 688–693
- 3 *Nonaka K, Ohata K, Nakai Y.* Probe-based confocal laser endomicroscopy of the duodenal mucosa with fluorescein dispersion. *Dig Endosc* 2014; 26: 604

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1390731>
Endoscopy 2015; 47: E16–E17
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Ken Ohata

Department of Gastroenterology
NTT Medical Center Tokyo
5-9-22 Higashi-gotanda Shinagawa-ku
Tokyo 141-8625
Japan
Fax: +81-3-34486541
ken.ohata1974@gmail.com