

Grasp-to-retract modification of the tulip-bundle technique in forward and retroflexed position for difficult hemostatic therapy in the sigmoid colon




Fig. 1 Endoscopic mucosal resection of a 7-cm, type 0-Is lesion in the distal sigmoid colon.




Fig. 2 After complete resection, a 4×4 cm mucosal defect over a colonic fold was apparent, with diffuse oozing but no visible vessels. The defect was closed using hemostatic clips, but diffuse oozing persisted between the clips.




Fig. 3 Attempts to place a detachable snare underneath the clips, in order to perform the tulip-bundle technique, were unsuccessful because of the large diameter of the defect with clips and its position over the colonic fold.




Fig. 4 Using a double-channel colonoscope, a grasping forceps was used to retract the defect while a detachable snare was positioned underneath the clips and closed, resulting in immediate hemostasis.




Fig. 5 In a second colonoscopy for hematochezia 4 hours later, oozing was observed from the proximal border of the defect. The same grasp-to-retract and tulip-bundle technique previously described was performed in retroflexed position in the sigmoid colon, resulting in prompt hemostasis.




Fig. 6 Final defect, in forward-viewing position, after the two tulip-bundle procedures, showing definitive hemostasis.

A 60-year-old woman with no relevant medical history underwent endoscopic mucosal resection (EMR) of a 7-cm 0-Is lesion in the distal sigmoid colon. A solution of saline, indigo carmine, and 1/100 000 adrenaline was injected into the submucosa, and piecemeal snare resection was performed (▶ **Fig. 1**). Persistent oozing occurred during EMR and was partially controlled by subsequent submucosal injections and resections. After complete resection, a 4×4 cm mucosal defect over a


colonic fold could be seen, with diffuse oozing but no visible vessels.

The defect was closed using hemostatic clips, but diffuse oozing persisted between the clips (▶ **Fig. 2**). Attempts to place a detachable snare (MAJ-254; Olympus, Tokyo, Japan) underneath the clips, in order to perform the tulip-bundle technique, were unsuccessful because of the large diameter of the defect with clips and its position over the fold (▶ **Fig. 3**, ▶ **Video 1**). Therefore, a double-channel

colonoscope (GIF 2T160I; Olympus) was used, and a grasping forceps was used to retract the defect while the detachable snare was positioned underneath the clips, resulting in immediate hemostasis (▶ **Fig. 4**, ▶ **Video 2**).


Despite initial hemostasis, the patient presented with hematochezia 4 hours later. Recurrent oozing from the proximal border of the mucosal defect, which had not been entrapped by the detachable snare, was observed and could not be

Video 1




Attempts to perform the tulip-bundle technique to treat diffuse oozing from a large endoscopic mucosal resection defect closed with clips were unsuccessful because of the large diameter of the defect with clips and its position over a colonic fold.

Video 2



A double-channel colonoscope was used to retract the defect with a grasping forceps while a detachable snare was positioned underneath the clips and closed, resulting in instant hemostasis.

Video 3



The same grasp-to-retract and tulip-bundle technique was performed in the retroflexed position in the sigmoid colon to treat bleeding that occurred from the proximal border of the defect 4 hours later.

treated with further clipping. With the endoscope in the retroflexed position, the tulip-bundle technique was attempted but was, again, unsuccessful. Using the double-channel colonoscope in the retroflexed position, and the same grasp-to-retract and tulip-bundle technique, definitive hemostasis was achieved (▶ Fig. 5, ▶ Fig. 6, ▶ Video 3).

Histology revealed a tubulovillous adenoma with high grade dysplasia.

Detachable snares have various indications that include assisting polypectomy, resecting submucosal tumors [1], and performing full-thickness resections [2]. The tulip-bundle technique involves the snare entrapping the clips to achieve hemostasis [3] or to close perforations [4]. This grasp-to-retract modification, which has been described previously for other techniques [5], can assist the tulip-bundle technique in difficult procedures.

Endoscopy_UCTN_Code_CPL_1AJ_2AD

Competing interests: None

Rolando Pinho¹, Joana Silva¹, Ana Ponte¹, Jaime Rodrigues¹, Iolanda Ribeiro¹, Maria Conceição Lucas², João Carvalho¹

¹ Gastroenterology Department, Centro Hospitalar de Vila Nova de Gaia, Vila Nova de Gaia, Portugal

² Surgery Department, Centro Hospitalar de Vila Nova de Gaia, Vila Nova de Gaia, Portugal

References

- 1 Veloso R, Pinho R, Rodrigues A et al. Endoloop ligation (“loop-and-let-go”) of a large ileal lipoma by balloon-assisted enteroscopy. *Endoscopy* 2012; 44 (Suppl. 02): E176
- 2 Pinho R, Oliveira M, Mascarenhas-Saraiva M. Endoscopic full-thickness resection of an inverted colonic diverticulum with intraepithelial neoplasia using the ligate-and-let-go technique. *Clin Gastroenterol Hepatol*. In press 2015. DOI: 10.1016/j.cgh.2015.08.007
- 3 Lee JH, Kim BK, Seol DC et al. Rescue endoscopic bleeding control for nonvariceal upper gastrointestinal hemorrhage using clipping and detachable snaring. *Endoscopy* 2013; 45: 489–492

4 Mocciano F, Curcio G, Tarantino I et al. Tulip bundle technique and fibrin glue injection: unusual treatment of colonic perforation. *World J Gastroenterol* 2011; 17: 1088–1090

5 Ponte A, Pinho R, Vale S et al. Resection of a large ileal lipoma exhibiting ball-valve pro-lapse into the cecum with a “grasp-to-retract, ligate, unroof, and let-go” technique. *Endoscopy* 2015; 47 (Suppl. 01): E215–216

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1393393>
 Endoscopy 2015; 47: E554–E555
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Rolando Taveira Pinho, MD

Serviço de Gastroenterologia
 Centro Hospitalar de Vila Nova de Gaia
 Rua Conceição Fernandes
 Vila Nova de Gaia 4434-502
 Portugal
 Fax: +351-227-868369
 rolandopinho@gmail.com