


Y. HATANAKA, T. HIYAMA\* (SAGAMI CHEMICAL RESEARCH CENTER, SAGAMIHARA, JAPAN)


Cross-Coupling of Organosilanes with Organic Halides Mediated by Palladium Catalyst and Tris(dimethylamino)sulfonium Difluorotrimethylsilicate

*J. Org. Chem.* **1988**, *53*, 918–920, DOI: 10.1021/jo00239a056.

## Cross-Coupling of Organosilanes with Organic Halides Mediated by Palladium Catalyst and TASF



### Selected examples:



**Significance:** In 1988, the authors reported a catalytic coupling of simple trimethylsilanes and organohalides in a process that is now known as Hiyama cross-coupling. Organosilanes are stable, easy to handle coupling partners that have low toxicity and good solubility. In addition to trialkylsilanes, trialkoxysilanes and fluoroalkylsilanes are also competent coupling partners. However, a stable C–Si bond requires activation, which is most commonly performed by using a F<sup>−</sup> source.

**Comment:** In this first report coupling proceeded under relatively mild conditions and sensitive products were obtained. However, the scope of organosilanes was very limited. Today, many protocols exist for coupling of organosilanes. Those that are fluoride-free are especially useful and desirable. The commercial availability of organosilanes remains a limiting factor in the widespread use of Hiyama coupling.

**Review:** Y. Nakao, T. Hiyama *Chem. Soc. Rev.* **2011**, *40*, 4893–4901.

**SYNFACTS Contributors:** Paul Knochel, Dimitrije Djukanovic  
*Synfacts* 2020, 16(12), 1451 Published online: 17.11.2020  
DOI: 10.1055/s-0040-1719506; Reg-No.: P14520SF

© 2020, Thieme. All rights reserved.  
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Metals in Synthesis

Key words

organosilanes

palladium catalysis

cross-coupling

Synfact  
Classic