



# Quad Fever in a Case of Cervical Cord Injury—A Rare Case Report

Sheena Ali<sup>1</sup> Duraisamy Ganesan<sup>1</sup> Varun Sundaramoorthy<sup>1</sup>

<sup>1</sup>Kovai Medical Centre and Hospital, Coimbatore, Tamil Nadu, India

AJNS 2022;17:85–87.

Address for correspondence Sheena Ali, MBBS, MS, DrNB SS, Department of Neurosurgery, Kovai Medical Centre and Hospital, 99, Avinashi Road, Coimbatore, 641015, Tamil Nadu, India (e-mail: dr.sheenaali90@gmail.com).

## Abstract

By definition, “quad fever” is an extreme elevation in body core temperature beyond 40.8°C (105.4°F) in a patient with spinal cord injury. This type of central nervous system hyperpyrexia is seen in spinal cord injury patients, particularly those with high cervical spine injury with quadriplegia. However, it has also been described in paraplegics with a mid- or higher level thoracic spine injury. The incidence of “quad fever” is rare, with the highest reported temperature being 44°C (111.2°F) with chronic fatal outcomes. Though the use of antipyretics is generally efficacious, they are considerably ineffective in treating the hyperpyrexia seen in this type of severe central autonomic thermoregulation.

Here, we present a case of high cervical spine injury in a 24-year-old male. The trauma resulted in a C3–5 level cord contusion with incomplete quadriplegia (ASIA [American Spinal Cord Injury Association Impairment Scale] grade B). The patient developed high grade fever of 106°F within a week of admission postoperatively.

Pancultures were negative and the wound was clean. Despite treatment with higher antibiotics and an infection disease specialist’s consult, no obvious etiology was found. Drug-induced fever and thyroid function tests were excluded in other less-common causes.

Based on the diagnosis of exclusion, “quad fever” was inferred as the cause. He had other signs of autonomic instability during the episodes such as bradycardia with hypotension. Our patient showed an almost early response to treatment to betablockers and antipsychotics after failure to respond to antibiotics, mechanical hypothermia, and antipyretics.

## Keywords

- ▶ hyperpyrexia
- ▶ quad fever
- ▶ thermoregulation
- ▶ spinal cord injury
- ▶ neurogenic fever

## Introduction

Hyperpyrexia (temperature > 42.1°C [106.7°F]) is rare in hospital settings. More common causes include malignant neuroleptic syndrome, cholinesterase deficiency, drug fever, and central fevers.<sup>1,2</sup>

Quad fever must be ruled out in a spinal cord injury (SCI) patient. It is an extreme elevation in body core temperature beyond 40.8°C (105.4°F) in a patient with SCI.<sup>3</sup> Those with a high cervical spine injury and paraplegics with a mid- or higher level thoracic spine injury<sup>3,4</sup> have recorded literature of this phenomenon.

published online  
June 28, 2022

DOI <https://doi.org/10.1055/s-0042-1748784>.  
ISSN 2248-9614.

© 2022. Asian Congress of Neurological Surgeons. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India


The highest reported temperature is 44°C (111.2°F)<sup>4</sup> which have all proved fatal. Antipyretics have been deemed ineffective in treating the hyperpyrexia seen in this severe autonomic thermodysgulation.<sup>3,4</sup>

The pathophysiology comes down to the autonomic dysfunction associated with the injury. Besides thermodysgulation, other features of bradycardia, arrhythmias, hypotension and orthostatic hypotension, neurogenic shock, autonomic dysreflexia, and hyperhidrosis are also observed.<sup>3</sup>

## Case Presentation

A 24-year-old male patient presented with history of fall on his neck while jumping on a trampoline. He complained of immediate loss of power and sensations below T4 level and loss of power and grip of both upper limbs with a left-sided predominance. (ASIA [American Spinal Cord Injury Association Impairment Scale] score-B). He presented in a vitally stable condition with a Glasgow coma scale (GCS) of 15 and was normotensive and normovolumic at the time of admission. His magnetic resonance imaging (MRI) cervical spine was suggestive of a flexion extension and rotational injury of the cervical spine with anterior wedge compression fracture with mild reterolisthesis, left lamina, and transverse process fracture of C5 and undisplaced fracture of left lamina and C5 process of C4 (► Fig. 1).

He was posted for an anterior cervical discectomy and fusion with corpectomy and placement of an OCTACAGE after starting him on the National Acute Spinal Cord Injury Study (NACIS) III protocol the following day. He was kept for



**Fig. 1** Anterior wedge compression fracture with mild retrolisthesis of C5 is seen causing mild spinal canal compromise. Undisplaced fracture is noted in left lamina, pedicle and transverse process of C4. Undisplaced fracture is noted in left lamina of C5 vertebra with extension to left transverse process. Mildly displaced vertical fracture is seen in the body of C6 vertebra.

observation in the intensive care unit (ICU), in view of inotropic support which was gradually weaned. However, on postoperative day (POD) 3, 5 days after the injury, he developed new onset hyperpyrexia (106°F) measured via digital thermometer in the axilla.

Pharmacologic efforts included 6 hourly does of paracetamol infusions (1 gm), intravenous steroid, 4-mg dexamethasone along with the use of empirical antibiotics like injectable meropenem and vancomycin. Physical measures of sponging with multiple ice packs to the axillary regions was done. Despite all of the above measures, the patient's body temperature continued to rise, peaking at 42°C (107°F). Vitally his heart rate was 113 beats per minute and blood pressure was 110/70 mm Hg (► Fig. 2).

It was then decided to start him on treatment for neurogenic fever, as all previous treatments were futile. He was started on oral betablockers with strict blood pressure monitoring and Emetil (chlorpromazine), an antipsychotic drug, on postoperative day 5.

His blood chemistry was significant for falling hemoglobin levels with leukocytosis (19,700), with an elevated blood urea nitrogen, and C-reactive protein (CRP) on 1 mg/dL. Creatinine phosphokinase (CPK) levels were 41 μ/L.

Normothermia was successfully achieved 37°C (98.6°F) after 12 hours. The diagnosis of "quad fever" was made as a diagnosis of exclusion retrospectively.

The patient, however, did develop another episode of hyperpyrexia (106°F) on POD 28, secondary to a urinary tract infection which was managed with appropriate antibiotics. The rest of his hospitalization was uneventful with an improving neurological status, good toleration to regular occupational therapy, and physiotherapy.

## Discussion

Patients with high spinal cord lesions and contusions suffer from thermodysgulation due to the alteration of the hypothalamus and other regulatory centers to counteract endogenous heat production.<sup>1,3,5</sup>

Hyperpyrexia as a side effect causes reduced blood flow to the brain, damaging central homeostatic mechanisms, further enhancing this vicious cycle, resulting in increased core body temperature.

At the cellular level, one can expect protein denaturation, alterations in signal transduction<sup>6,7</sup> at these high temperatures of 40–41°C (104–105.8°F).

Other vital systems may be affected in the form of coagulopathy, hypovolemia, acute tubular necrosis, abnormal liver enzymes, progressing to severe liver necrosis, brain edema, extensive neuronal loss and gliosis, and intraparenchymal hemorrhage in the adrenal gland bone marrow suppression resulting in thrombocytopenia, as well as intravascular hemolysis may also be noted.<sup>8,9</sup>

The main purpose of achieving normothermia is to ensure cardio protection and in turn neuroprotection, hence breaking the vicious cycle.

Literature states use of multiple antipyretic drugs, prophylactic antibiotics, invasive intravenous cooling, sponging,

