



# Outcomes of Combined Visual and Auditive Stimulation on Functions of Hand and Grip Strengths in Patients with Hemiplegia

Bhaswati Baishya<sup>1</sup> Kalidasan Varathan<sup>1</sup>

<sup>1</sup> Krupanidhi College of Physiotherapy, Bengaluru, Karnataka, India

Address for correspondence Kalidasan Varathan, MPT, Krupanidhi College of Physiotherapy, Bengaluru, Karnataka, 560 035, India (e-mail: rvkalidasan@gmail.com; physio.kric@krupanidhi.edu.in).

J Health Allied Sci<sup>NU</sup> 2023;13:199–205.

## Abstract

**Background** Stroke is a major ailment that results from hypoxia, ischemia, blockage, or hemorrhage. The recovery of hand functions is an essential goal in stroke patients' recovery.

**Aim** This study aimed to analyze the outcomes of rhythmic auditory stimulation (RAS) and mirror therapy on hand functions, as well as grip strength, in stroke patients.

**Materials and Methods** A quasiexperimental study has been used based on criteria of inclusion, 30 patients were enrolled. The participants were assigned to two groups, 15 patients each. Group A (control group) underwent traditional physiotherapy. Group B (experimental group) underwent RAS and mirror therapy. Both the groups received 20 minutes of treatment, 20 sessions in 1 month. The Action Research Arm Test (ARAT) and hand-held dynamometer were used to evaluate results. Paired *t*-test has been used to analyze the data with the SPSS software tool.

**Results** The paired *t*-test results showed that notable distinctions in ARAT mean pretest scores between two groups were not found. A notable change in ARAT results was present between the two groups with mean posttest score and difference in pretest to posttest. No notable distinctions in grip strength results were seen with mean pretest results. But a notable change was found in grip strength results among two groups with mean posttest scores.

**Conclusion** Combining RAS and mirror therapy programs has shown beneficial effects on hand functions, as well as grip strength of stroke patients and can be used as adjunct interventions along with conventional physiotherapy to encourage restoration of hand functions in hemiparetic patients.

## Keywords

- ▶ rhythmic auditory stimulation
- ▶ mirror therapy
- ▶ stroke
- ▶ the action research arm test
- ▶ grip strength

## Introduction

Stroke can be defined as evolving medical symptoms of principal disruption of cerebral function of which the symptoms can continue for 24 hours or more and can lead to death.<sup>1</sup> The neurological presentations relate to the extent of injury, site of the part engaged, as well as fundamental causes

of brain dysfunction.<sup>2</sup> Globally, stroke results in 12.6 million individuals having a disability of which 8.9 million people belong to lower middle-income countries.<sup>3</sup>

Stroke accounted for 41% of deaths and 72% of disabilities in our country.<sup>4</sup> In India, the second leading cause of death is stroke, the most significant cause of disability.<sup>5,6</sup> Upper extremity function is damaged in approximately 70% of these

article published online  
July 1, 2022

DOI <https://doi.org/10.1055/s-0042-1749422>.  
ISSN 2582-4287.

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

individuals.<sup>7</sup> Damage to the upper extremity function of hemiplegic patients critically disrupts their ability to perform daily tasks independently.<sup>8</sup>

Following the onset of stroke, motor paralysis appears, and the recovery of finger extension function takes the longest period.<sup>9</sup> Given that many daily activities require using movements of fingers and hands, patients, who are not able to use their hands after the onset of stroke, experience a certain level of physical and mental pain.<sup>10</sup>

Mirror therapy is an easy and feasible intervention for upper limb training. The purpose is restoring the upper limb dexterity while asking the patient to emphasis on unininvolved extremity movements.<sup>11</sup>

Rhythmic auditory stimulation (RAS) is another treatment intervention that has shown to be beneficial in enhancing upper limb movements.<sup>12</sup> RAS enhances motor control by promoting organization and accomplishment by powerful entrainment and integrated outcomes of repeated rhythmic sensory signals on the motor system.<sup>13</sup>

Park et al stated stroke survivors accounting for 85%, undergo paralysis, and 69% of patients account for disrupted functioning of upper limbs.<sup>8</sup> The hand and digits dexterity recovery makes a crucial contribution to the rehabilitation of hemiplegic patients.<sup>11</sup> Because of the limited evidence of RAS on hand functions in hemiplegic patients, a need arises for looking into its uses together with mirror therapy in improving the hand functions, as well as strength, of the hemiparetic hand. The framed null hypothesis was that there would be no significant effect of RAS and mirror therapy on the hand functions and grip strength of the stroke patients. The alternate hypothesis was there would be a significant effect of RAS and mirror therapy on the hand functions and grip strength of the stroke patients.

## Objective

This study aimed to evaluate the (1) effect of conventional therapy on hand functions and grip strength of stroke patients, (2) combined effects of RAS and mirror therapy on the hand functions and grip strength of the stroke

patients, and (3) compare the effects of conventional therapy against the combined effects of RAS and mirror therapy on the hand functions and grip strength of the stroke patients

## Materials and Methods

The study design is a pre-post quasiexperimental study involving patients clinically diagnosed with a stroke, duration of 6 months from onset, both the genders, aged 50 to 65 years for 4 weeks.<sup>8,11</sup> A total of 30 patients were conveniently selected as per the inclusion criteria. Patients with anterior cerebral artery (ACA) and middle cerebral artery (MCA) involvement with the affected side being the dominant side and those with Brunnstrom's recovery stages of 3 and 4 took part.<sup>8</sup> Patients with dementia, depression, or productive psychosis were not included.<sup>2</sup> Patients having any visual or auditory impairments were excluded.<sup>14</sup> The patients were grouped into the following: the control group receiving traditional physiotherapy and the experimental group receiving exclusively mirror therapy plus RAS. The purposive sampling technique was used to group the patients into control and experimental group.


### Procedure

►Fig. 1 shows the flowchart for the procedure. Overall, 20 treatment sessions were given for 1 month for both the groups. Assessments were made before and after the treatment sessions using hand-held dynamometer and ARAT. Conventional physiotherapy was provided for 20 minutes to the control group in each session.

### Conventional Physiotherapy

#### Tone Normalization of Spastic Muscles

Slow sustained stretching of the agonist spastic muscles through range of motion (ROM). Active exercises focused on the activation of the weak antagonist muscles using slow and controlled movements were performed. Local facilitation techniques, such as stretching, tapping, and light resistance, were added to enhance the action of the weak



**Fig. 1** Flowchart for the procedure. ARAT, action research arm test.

antagonist muscles.<sup>15</sup> Tapping to the antagonist muscle was given to facilitate voluntary movement out of synergy pattern which caused reciprocal inhibition of the spastic muscle. Tone normalization was provided for 5 minutes.

**To maintain Range of Motion**

ROM exercises were given for 5 minutes.

**Sensory Reeducation**

Stroking through thenar, as well as hypothenar, eminences five times to achieve purposeful muscular contractions. Superficial and deep pressure were applied. Localization to touch was to allowing patients to touch objects of various sizes, shapes, and materials. Sensory re-education was given for 5 minutes.

**Encouraging Voluntary Movement**

Patients were encouraged to make use of their involved hand for performing daily activities such as grooming and dressing. The patients performed all the voluntary activities for 5 minutes. Integrated mirror therapy and RAS for 20 minutes were rendered to the experimental group in each session.

**Rhythmic Auditory Stimulation**

Participants were made to sit in a back-supported chair, with the affected hand placed on the desk.<sup>14</sup> RAS using an electronic smartphone-based metronome was given. Patients performed hand movements such as grasping a ball and releasing it, rolling the ball from the tip of fingers to the palm, pinching a ball, stacking coins, and other physiological movements of the hand. Patients executed the repeated movement sequence with the metronome beats in time.<sup>14</sup> To find out the preferred auditory stimulation frequency, each patient was asked to perform a repetition of each task for 1 minute using the affected hand.<sup>14</sup> The advantage of using a smartphone is that it can be easily operated and is instantly available (►Table 1).<sup>16</sup>

**Mirror Therapy**

A mirror box training session followed this. The involved hand of the patient was asked to place at the back of the box while the uninvolved extremity ahead of the box. Participants performed various physiological forearm and hand movements, viewing the uninvolved extremity image, there-

by watching the reflection of movements of the hand projected over the affected hand.<sup>10,16</sup> The patients performed the same physiological activities simultaneously using the paretic hand (►Table 1).<sup>17</sup>

**Outcome Measures**

- Primary outcome measure was ARAT and secondary outcome measure was
- hand-held dynamometer.

**Statistical Analysis**

Statistical analyses are listed in ►Tables 2–7.

**Results**

Notable changes among the groups were not found: control (57.07 ± 4.079) and experimental (57.87 ± 4.824) groups with the distribution of age and mean age (*p* = 0.666) and gender distribution and mean (*p* = 0.719).

A notable difference in ARAT scores between the groups was not seen: control (27.07 ± 8.852) and experimental (27.33 ± 10.560) groups with mean pretest scores (*p* = 0.937). But notable changes in ARAT results were present in the groups: control (32.20 ± 8.986), and experimental (39.07 ± 8.345) groups with mean posttest scores (*p* = 0.021) and change from pretest to posttest (*p* = 0.0001) scores.

This shows that the experimental group is showing maximum changes in ARAT results compared with the control group.

No notable distinctions in grip strength results were observed among groups: control (14.20 ± 4.329) and experimental (14.67 ± 3.773) groups with mean pretest scores (*p* = 0.753). Notable distinction was observed: control (17.00 ± 4.375) and experimental (20.33 ± 4.923) groups with mean posttest scores (*p* = 0.059); change from pretest to posttest was noted (*p* = 0.0001).

It shows that the experimental group shows maximum changes or improvement in grip strength scores than the control group.

Thus, it shows that the integrated visual and auditory stimulation (RAS and mirror therapy) improved functions of the hand and grip strength among hemiparetic patients. Statistically significant values were observed for the experimental group.

**Table 1** Duration, frequency, and activities for the treatment interventions

Treatment interventions	Duration	Frequency	Activities to be performed
Rhythmic auditory stimulation	10 minutes	5 days/week	<ul style="list-style-type: none"> <li>• Grasping a ball and releasing it</li> <li>• Rolling the ball from the tip of the fingers to the palm</li> <li>• Pinching a ball</li> <li>• Stacking coins</li> <li>• Other physiological movements of the hand</li> </ul>
Mirror therapy	10 minutes	5 days/week	<ul style="list-style-type: none"> <li>• Supination and pronation</li> <li>• Flexion–extension of wrist and finger, finger movements</li> <li>• Sponge squeezing</li> <li>• Swiping a table with a towel etc.</li> </ul>

**Table 2** Comparison of the groups by age

Variable	Paired differences					t	df	Significance (two-tailed)
	Mean	Standard deviation	Standard error mean	95% confidence interval of the difference				
				Lower	Upper			
AGE_E AGE_C	0.800	7.022	1.813	-3.089	4.689	0.441	14	0.666

Note: Notable distinctions were not present among the two groups (experimental group and control group) in age distribution and mean age ( $t = 0.441$  and  $p = 0.666$ ; ► **Table 2**).

**Table 3** Comparison of the groups by gender

Variable	Paired differences					t	df	Significance (two-tailed)
	Mean	Standard deviation	Standard error mean	95% confidence interval of the difference				
				Lower	Upper			
Gender	0.067	0.704	0.182	-0.323	0.456	0.367	14	0.719

Note: Notable distinctions were not present among the two groups (experimental and control group) in gender distribution and mean age ( $t = 0.367$  and  $p = 0.719$ ; ► **Table 3**).

**Table 4** Pre-post action research arm test (ARAT) scores

Variable		Paired differences					t	df	Significance (two-tailed)
		Mean	Standard deviation	Standard error mean	95% confidence interval of the difference				
					Lower	Upper			
Pair 1: experimental group	Pre-post scores of ARAT	-11.73	4.35	1.123	-14.14	-9.324	-10.45	14	0.000
Pair 2: control group	Pretest scores of ARAT, posttest scores of ARAT	-5.133	1.685	0.435	-6.066	-4.2	-11.8	14	0.000

Note: The mean of paired difference of pre-post ARAT scores in the control group was -5.133 with standard deviation of 1.685, which is statistically significant ( $p < 0.0001$ ; ► **Table 4**).

**Table 5** Pre-post grip strength scores

Variable		Paired differences					t	df	Significance (two-tailed)
		Mean	Standard deviation	Standard error mean	95% confidence interval of the difference				
					Lower	Upper			
Experimental group	Pretest scores of grip strength (kg), posttest scores of grip strength	-5.667	2.289	0.591	-6.934	-4.399	-9.589	14	0.000
Control group	Pretest scores of grip strength, posttest scores of grip strength	-2.8	1.612	0.416	-3.693	-1.907	-6.725	14	0.000

Note: The mean of paired difference of pre-post grip strength scores in the control group was -2.880 with standard deviation of 1.612 which is statistically significant ( $p < 0.0001$ ; ► **Table 5**).

**Table 6** Paired differences of pretest scores of action research arm test (ARAT) and grip strength

Variable	Paired differences					t	df	Significance (two-tailed)
	Mean	Standard deviation	Standard error mean	95% confidence interval of the difference				
				Lower	Upper			
Pretest scores of ARAT between the experimental and control group	0.267	12.87	3.323	-6.861	7.394	0.08	14	0.937
Pretest scores of grip strength (kg) between the experimental and control group	0.467	5.643	1.457	-2.658	3.591	0.32	14	0.753

Note: The values obtained show that the subjects were homogenous in both groups (► **Table 6**).

**Table 7** Paired differences of posttest scores of ARAT and grip strength

Variable	Paired differences					t	df	Significance (two-tailed)
	Mean	SD	Standard error mean	95% confidence interval of the difference				
				Lower	Upper			
Posttest scores of ARAT between the experimental and control group	6.867	10.19	2.631	1.224	12.51	2.61	14	0.021
Posttest scores of grip strength between the experimental and control group	3.333	6.275	1.62	-0.142	6.809	2.057	14	0.059

Abbreviations: ARAT, action research arm test; SD, standard deviation.

Note: The mean of paired differences of scores of ARAT posttest was 6.867 with SD = 10.19 ( $p = 0.021$ ). The mean of paired differences in grip strength score was 3.333 with SD = 6.275 ( $p = 0.059$ ), which is statistically significant (► **Table 7**).

### Discussion

Stroke often results in a paretic hand. After 6 months from the duration of stroke, patients regaining dexterity in manipulative tasks accounted for 38%, while only 11.6% achieved complete functional recovery in hand dexterity.<sup>18</sup>

Differences in hand functions and grip strengths were evaluated and compared. The application of RAS by a smart-phone-based metronome application showed improvements in grip strength, as well as various physiological hand functions, while mirror therapy provided the appropriate visual feedback.

The results obtained after the paired *t*-test analysis showed that there was a recommendable improvement in both the outcome measures in the experimental group: ARAT score ( $39.07 \pm 8.345$ ) and grip strength score ( $20.33 \pm 4.923$ ). The findings matched the results of Street et al where significant mean values of ARAT ( $29.80 \pm 18.75$ ) were observed on the application of auditory stimulation which was designed for home-purpose for stroke patients with hemiparesis of the arm.<sup>13,19</sup> Raglio et al found recommendable recovery in grip strength ( $24.91 \pm 11.55$ ) in their study.<sup>20</sup> The improvements in the strength of grip functions observed are similar to the improvements observed in El Shemy and Abd El-Maksoud in which strength of grip

functions improved profoundly ( $10.63 \pm 1.49$  vs.  $5.7 \pm 1.42$ ,  $p < 0.05$ ) were observed.<sup>21</sup> The obtained results also come in agreement with Sathian et al.<sup>22</sup>

After the analysis of the paired *t*-test, the results showed improvement in both the outcome measures in the control group: ARAT score ( $32.20 \pm 8.986$ ) and grip strength score ( $17.00 \pm 4.375$ ).

Recommendable improvement in ARAT scores postassessment was noted. A significantly higher effect in ARAT scores was seen among the experimental group. The positive results in the ARAT scores were due to the improvement of considerable dexterity which emphasized four aspects, that is, pinch, grip, grasp, and gross movements of the hand.

A significantly higher effect in grip strength scores was seen in the experimental group than in the control group. Grip strength improvement is related to improvements in complicated motor functions which shows that grip strength can be used as an indicator of the functional recovery of the hand and arm.

Conventional physiotherapeutic approaches are influenced by the ability of central nervous system (CNS) to reorganize itself to relearn and perform various cognitive and motor functions. Hence, the results can be ascribed to repetitions of specific tasks and exercises, local facilitation techniques, and motor relearning strategies.<sup>23</sup>

The outcomes show the improvement in quality of movement and enhanced motor control using RAS in stroke patients. The programmed movements produced by RAS were smooth, effective, and brought about wider ROM. The study results show that rhythmic cueing improves the quality of movement and motor control in stroke patients. RAS allowed for a programmed movement that was more efficient and smoother and had wider ROM. The results are likely the result of auditory rhythms transmitted to the motor system which provided constant feedback while performing the various exercises. The auditory and motor system have rich connectivity across various cortical, sub-cortical, and spinal levels. The auditory system projects into motor structures in the brain, creating entrainment between the rhythmic signal and the motor response.

Rhythmic auditory cueing has three advantages. First, holding frequency constant ensures that the same movement is actually repeated. The auditory cueing may entrain the motor system to its beat. Second, matching the sound with full extension or flexion of fingers provides an attentional goal for the patient. Goal setting is also known to promote motor learning. Third, receiving feedback is fundamental to motor learning. In this study, sensory information from the audio cues and visual and somatosensory sources provided intrinsic feedback to the patient regarding the movement goal.<sup>14</sup> The profound effects of RAS on movement have been explained by the auditory-motor entrainment which means innate synchronization of movement rhythm to the regular beat of music/sounds.<sup>24</sup>

The experimental group demonstrated better results in accomplishing the daily tasks of living compared with the control group. These results support the study of Stevens and Stoykov<sup>25</sup> study, stating that observing movements through mirror therapy generates positive feedback. Observation of normal movement provided positive visual feedback and improved the function of their affected limb without it being moved. Normal movement is thought to be induced due to activating the premotor cortex by recalling the proprioception (individual perception) that was reduced or removed when normal visual feedback was provided.<sup>19</sup> Mirror therapy, when used with daily functional activities, enhanced the motor recovery of the paretic upper extremity in stroke patients. Mirror therapy provides sustained regular visual input of movement which may better promote central brain remodeling.<sup>26</sup>

## Limitations and Recommendations

Stroke patients within 6 months from onset were taken into account, hence the homogeneity among the patients hence the homogeneity among the patients were maintained. The long-term effect of the interventions is not known as the research was conducted only for a month without follow-ups. Also, patients with Brunnstrom's stages 3 and 4 were included in the study, leading to nonuniformity in the pre-post test scores between and among the groups. Interventions according to various Brunnstrom's stages and individualized to the patients' functional needs should be developed.

## Conclusion

The study can be concluded by stating that integrated visual and auditory stimulation has beneficial effects on hand functions and strength in hemiparetic stroke patients. However, when compared with the conventional approach, the results of research conclude that combined effects of RAS and mirror therapy have beneficial effects on restoration and improvement of the hand functions. Both these treatment interventions can be used as suitable adjuncts along with conventional physiotherapy to encourage and facilitate the restoration of hand functions in hemiparetic patients.

### Conflict of Interest

None declared.

### Acknowledgment

The authors extend their heartfelt gratitude to the Management of Krupanidhi College of Physiotherapy and Krupanidhi Research and Incubation Center (KRIC) for laying down immense guidance and support throughout.

## References

- 1 Thorvaldsen P, Asplund K, Kuulasmaa K, Rajakangas AM, Schroll MWorld Health Organization Monitoring Trends and Determinants in Cardiovascular Disease. Stroke incidence, case fatality, and mortality in the WHO MONICA project. *Stroke* 1995;26(03): 361–367
- 2 Diamond PT, Gale SD, Evans BA. Relationship of initial hematocrit level to discharge destination and resource utilization after ischemic stroke: a pilot study. *Arch Phys Med Rehabil* 2003;84 (07):964–967
- 3 Feigin VL, Norrving B, Mensah GA. Global burden of stroke. *Circ Res* 2017;120(03):439–448
- 4 Taylor FC, Suresh Kumar K. Stroke in India fact sheet (updated 2012). Accessed April 28, 2022 at: [https://www.researchgate.net/profile/Suresh-Kumar-85/publication/264116605\\_Stroke-in-India\\_-Fact-sheet\\_Updated\\_2012/links/550bdcfb0cf265693cefb81a/Stroke-in-India-Fact-sheet-Updated-2012.pdf](https://www.researchgate.net/profile/Suresh-Kumar-85/publication/264116605_Stroke-in-India_-Fact-sheet_Updated_2012/links/550bdcfb0cf265693cefb81a/Stroke-in-India-Fact-sheet-Updated-2012.pdf).
- 5 Gobindram A. Stroke care: experiences and clinical research in stroke units in Chennai. *Ann Indian Acad Neurol* 2006;9(04): 193
- 6 Yan T, Hui-Chan CW, Li LS. Functional electrical stimulation improves motor recovery of the lower extremity and walking ability of subjects with first acute stroke: a randomized placebo-controlled trial. *Stroke* 2005;36(01):80–85
- 7 Luke C, Dodd KJ, Brock K. Outcomes of the Bobath concept on upper limb recovery following stroke. *Clin Rehabil* 2004;18(08): 888–898
- 8 Park JY, Chang M, Kim KM, Kim HJ. The effect of mirror therapy on upper-extremity function and activities of daily living in stroke patients. *J Phys Ther Sci* 2015;27(06):1681–1683
- 9 Trombly CA, Quintana LA. The effects of exercise on finger extension of CVA patients. *Am J Occup Ther* 1983;37(03):195–202
- 10 Kim JH, Lee BH. Mirror therapy combined with biofeedback functional electrical stimulation for motor recovery of upper extremities after stroke: a pilot randomized controlled trial. *Occup Ther Int* 2015;22(02):51–60
- 11 Hatem SM, Saussez G, Della Faille M, et al. Rehabilitation of motor function after stroke: a multiple systematic review focused on techniques to stimulate upper extremity recovery. *Front Hum Neurosci* 2016;10:442

- 12 Ahn JY, Kim H, Park CB. Effects of whole-body vibration on upper extremity function and grip strength in patients with subacute stroke: a randomised single-blind controlled trial. *Occup Ther Int* 2019;2019:5820952
- 13 Street AJ, Magee WL, Bateman A, Parker M, Odell-Miller H, Fachner J. Home-based neurologic music therapy for arm hemiparesis following stroke: results from a pilot, feasibility randomized controlled trial. *Clin Rehabil* 2018;32(01):18–28
- 14 Kim JR, Jung MY, Yoo EY, Park JH, Kim SH, Lee J. Effects of rhythmic auditory stimulation during hemiplegic arm reaching in individuals with stroke: an exploratory study. *Hong Kong J Occup Ther* 2014;24(02):64–71
- 15 Sullivan O'SB, Schmitz TJ, Fulk GD, eds. *Physical Rehabilitation. Stroke*. 6th edition. Philadelphia, PA: F.A. Davis Company; 2014:685
- 16 Jung-Hee KI, Sung-Gook PA, Hyun-Jung LI, Gyung-Choon PA, Moon-Hyung KI, Byoung-Hee LE. Effects of the combination of rhythmic auditory stimulation and task-oriented training on functional recovery of subacute stroke patients. *J Phys Ther Sci* 2012;24(12):1307–1313
- 17 Da Silva Mendes JC, Freysteinson WM, de Melo Gamboa R, Pereira V. Mirror Therapy: can be this technique used on facial disfigurement? *Revista E-Psi*. 2017;7(01):25–38
- 18 Kim JH, Lee B. The effect of mirror therapy on functional recovery of upper extremity after stroke: a randomized pilot study. *J Exp Stroke Transl Med* 2017;10(01):1–7
- 19 Paik YR, Kim SK, Lee JS, Jeon BJ. Simple and task-oriented mirror therapy for upper extremity function in stroke patients: a pilot study. *Hong Kong J Occup Ther* 2014;24(01):6–12
- 20 Raglio A, Zaliani A, Baiardi P, et al. Active music therapy approach for stroke patients in the post-acute rehabilitation. *Neurol Sci* 2017;38(05):893–897
- 21 El Shemy SA, Abd El-Maksoud GM. Mirror therapy versus cross education on wrist extension and hand grip strength in children with hemiparesis. *Bull Fac Phys Ther* 2009;14(02):
- 22 Sathian K, Greenspan AI, Wolf SL. Doing it with mirrors: a case study of a novel approach to neurorehabilitation. *Neurorehabil Neural Repair* 2000;14(01):73–76
- 23 Dobkin BH, Dorsch A. New evidence for therapies in stroke rehabilitation. *Curr Atheroscler Rep* 2013;15(06):331
- 24 Wang SM, Lin CY, Tse TH, et al. Effects of rhythmic auditory stimulation on upper-limb movement speed in patients with schizophrenia spectrum disorders. *Eur Arch Psychiatry Clin Neurosci* 2021;271(08):1445–1453
- 25 Jennifer A Stevens, Mary Ellen Phillips Stoykov Using motor imagery in the rehabilitation of Hemiparesis. *Arch Phys Med Rehabil* 2003;84(07):1090–1092
- 26 Zhuang JY, Ding L, Shu BB, Chen D, Jia J. Associated mirror therapy enhances motor recovery of the upper extremity and daily function after stroke: a randomized control study. *Neural Plast* 2021;2021:7266263