M. J. ODDY, D. A. KUSZA, R. G. EPTON, J. M. LYNAM, W. P. UNSWORTH, W. F. PETERSEN* (UNIVERSITY OF CAPE TOWN, SOUTH AFRICA)

Visible-Light-Mediated Energy Transfer Enables the Synthesis of β-Lactams via Intramolecular Hydrogen Atom Transfer *Angew. Chem. Int. Ed.* **2022**, *61*, e202213086 DOI: 10.1002/anie.202213086.

Visible Light-Mediated β -Lactam Synthesis from Acryl Amide Precursors

Selected examples

- Proposed mechanism

Significance: Petersen and co-workers report a visible light-mediated β -lactam synthesis from acryl amide precursors catalyzed by 2-chlorothioxanthone (2-CTX). The corresponding 2-azetidinones are obtained in moderate to excellent yields with poor to moderate diastereoselectivities. Mechanistic investigations including DFT calculations, deuterium labeling studies and competition experiments support a variation of the Norrish-Yang type II photocyclization being the dominant mechanistic pathway.

Comment: Supported by DFT calculations, an ionic Staudinger-like pathway is ruled out by the authors. Instead, a reaction mechanism is proposed that is initiated by an energy transfer (EnT) from the excited photosensitizer to the acryl amide, furnishing its triplet excited state. A 1,5-hydrogen atom transfer (HAT) is followed by an intersystem crossing (ISC) to its singlet state. Radical-radical coupling ultimately leads to the β -lactam formation. Notably, the lack of diastereoselectivity that is expected from this radical pathway, is mostly in agreement with the experimental results.

SYNFACTS Contributors: Benjamin List, Wencke Leinung Synfacts 2023, 19(01), 0077 Published online: 16.12.2022 **DOI:** 10.1055/s-0042-1751785; **Reg-No.:** B00723SF

Organo- and Biocatalysis

Key words

β-lactams
photochemistry
energy transfer
hydrogen atom
transfer

